ON ALBUMINURIA.

*Read in the Section of Medicine.*BY SIR ANDREW CLARK, BART., M.D.,
Senior Physician to the London Hospital.

ON this occasion, it is not my intention to deal with the complex problems raised by questions concerning that disturbance of physiological conditions which issues in the appearance of albumen in the urine. I shall confine myself to a much humbler task; and, regarding the subject from a merely clinical stand-point, and avoiding those aspects of it which have been already much discussed, I shall pass beyond empirical experience no farther than may be necessary to make plain a few of the problems ripening for physiological pathology to solve.

In all preceding discussions about albuminuria, the speakers seem to me, if I am not in error, to deal with the presence of albumen in the urine as if it were of renal origin; or, at any rate, they neither sufficiently distinguish renal from non-renal albuminuria, nor explicitly set forth, if they are known, the grounds of discriminating between them.

To avoid confusion in the course of the discussion, it would, therefore, seem necessary to recognise the existence of non-renal albuminuria, to specify in cases of this sort the sources of the albumen, and to set forth, so far as our present knowledge will permit, the means whereby renal may be distinguished from non-renal albuminuria.

In a good many women, for a few days before and for a few days after menstruation, the urine, free from blood-discs, leucocytes, or pus, contains, sometimes continuously, sometimes intermittently, small quantities of albumen. In women guilty of habits of secret personal impurity, a serous fluid is sometimes secreted into the vagina; and afterwards, mixing with the urine, is found therein, responding in the usual manner to the tests for serum-albumen.

In some young men, excited by sexual desire and denying it indulgence, there is secreted from the urethra or its adjacent glands a fluid which, mixing with the urine, yields to the application of the ordinary tests evidence of the presence of serum-albumen.

In eczema of the bladder, in the early stages of villous tumour, and in the venous congestions of aged men, albumen transudes into the vesical cavity, mixes with the urine, and may be readily mistaken for albumen of renal origin.

In the case of the late E. D., about whom I was consulted for renal disease, the albumen in the urine was proved to be independent of renal disorder, and to have its origin in a small delicate villous tumour of the bladder.

I am sure that illustrations of non-renal albuminuria could be added to by others present at the discussion better than by me; but I have adduced from my own experience a number of examples, sufficient to warrant further investigation, and to justify me in bringing the subject to the notice of the Section.

And now arises the question: In what way can we distinguish the non-renal from the renal albumen? I suppose that we shall be very near the truth if, in a given case of albuminous urine, where there are no constitutional evidences of renal disease, where the urine is of normal constitution *plus* albumen, and where some local cause adequate to its production exists, we say that the albumen is non-renal. But we shall not be certain. For any one of the specified local causes may exist without albumen necessarily passing into the urine; and the kidney may yield albumen to the urine without any material alteration of that secretion. In such possible, but rare, cases, one must critically watch the state of the constitution, which rarely fails to give early information of renal degeneration.

Of the non-renal forms of albuminuria occurring in leucorrhoea, cystitis, catarrh of the ureters, and other affections in which histological elements are always present in the urine, it may be said that their diagnosis is attended with no serious difficulty; but, in such cases, one may easily forget that the existence of an albumen of non-renal does not exclude the existence of an albumen of renal origin; and, indeed, that not unfrequently they concur.

I pass now to the subject of cases of renal albuminuria, and divide them provisionally into those of functional and those of structural origin. So much has been already said, and so much will be again said, at the meeting concerning structural albuminuria, that I will leave its discussion entirely in the hands of others more competent, and also more eager, to deal with it than I am. For, to say the truth, I think that, in the present state of knowledge, the discussion of structural is of less interest and of less importance than the discussion of

functional albuminuria, to which I shall confine the few observations I have to offer.

To the use of the term "functional," many general and some just objections will be raised. It will be urged that every lesion of function must have its correlative lesion of structure; and that, to use any term which appears to deny this fact, or fails to give it at least implicit recognition, is at variance with the principles of science, and constitutes a backward instead of a forward movement in knowledge. To the logical coherency and force of this argument, I make no objection; but I contend, as often before I have contended, that mere logical integrity is not, in such matters, the final test or measure of truth, which, in the present state of knowledge, can sometimes be reached only along lines which seem, as we understand them, at present to be illogical. And in no way could this paradox be better illustrated, or more strongly enforced, than by a critical study of what is called functional albuminuria. It is, I venture to say, certain that, in the kidney giving rise to albumen in the urine, there occur states—mechanical, physical, chemical, and, in a provisional sense, vital—which are neither tangible nor visible, which not only cannot be estimated, but are even, by the most delicate instruments of research, incapable of recognition; states which may often come and go, disordering function and disturbing health, and yet leave no abiding marks of their presence and actions. Such states are different in manifold ways from organic states, and must be so named as to ensure recognition of their fundamental differences. For this is not all that has to be said concerning these functional states. We are so much concerned with anatomical changes; we have given so much time to their evolutions, differentiations, and relations; we are so much dominated by the idea that, in dealing with them, we are dealing with disease in itself, that we have overlooked the fundamental truth, that these anatomical changes are but secondary, and sometimes the least important, expressions or manifestations of states which underlie them. It is to these dynamic states that our thoughts and inquiries should be turned; they precede, underlie, and originate structural changes; they determine their character, course, and issues; in them is the secret of disease; and, if our control of it is ever to become greater and better, it is upon them that our experiments must be made.

I say, therefore, that those functional affections of the kidneys, attended by the presence of albumen in the urine, are of the utmost importance to the better understanding of organic disease, and deserve a more prolonged and critical study than they have yet received.

Of the forms of functional albuminuria with which I am practically acquainted, I shall mention four as worthy of further consideration. They are the nervous, the oxaluric, the hepatic, and the gouty. I leave on one side the functional albuminuria of cold, and the various forms of peptonuria connected with indigestion.

The first two forms occur chiefly among adolescents; the latter two are found for the most part among elderly people.

I am as sure as I can be about anything incapable of demonstration, that all strain of nervous system, especially under emotional excitement, is capable of producing functional albuminuria. Among twenty men entering a competitive examination, no one is albuminuric; at the close of the examination, lasting a week, three are found to have albumen in the urine. A gouty man with moderately healthy urine attends a political meeting, and delivers an exciting speech. Soon afterwards, a little albumen is found in the urine; but in a day or two it disappears. In a fortnight, he delivers another speech in circumstances of great excitement, and the albumen reappears in the urine.

The most numerous illustrations which I have met of functional albuminuria have occurred in young men aged from eighteen to thirty, whose urine was of high density, and loaded with oxalate of lime. To put the matter in another way, I have not very often followed carefully a case of "oxaluria" with high density and an excess of urea, without finding sooner or later traces of albumen in the urine. This is by far the most interesting and instructive of cases, and I regret that the time allotted to such communications as this will not permit me to consider them in detail.

Illustrations of the hepatic group occur for the most part in middle-aged men. There is congestive enlargement of the liver with catarrh; the portal system becomes loaded; the skin is dry and icteric; and then, without any appreciable change in the urine beyond the presence of a little bile, albumen appears in it. With the subsidence of the hepatic trouble, the albumen disappears from the urine.

Temporary albuminuria in gouty persons whose kidneys are as yet structurally unaffected is common enough in certain conditions. When the balance between ingoing and outgoing is disturbed, when the blood becomes loaded with excrementitious stuffs, when there is increasing vascular tension, with restlessness, feverishness, dry skin, and headache, then, as a common rule, traces of albumen appear in the urine.