movable medium ; that every movement of the body would keep this in continual motion, and as this pouch was thus increased, and the weighty mercury constantly exerting pressure and a dragging, now on this side now on that, and more and more still as it became more and more distended, and the tendency of all this to pull at the upper descending part of intestine, and drag it over the reflected brim of the detained gut, ultimately we might hope to attain the end we had in view. For this purpose I rather encouraged his journey in his springless cart homewards on the rough road. The result was soon seen. He had not gone two miles when there was an alvine discharge of satisfactory amount ; but as to its colour, or the presence of mercury, there was no ascertaining, it being dusk. However, the perusal of the accounts received leaves no room to doubt that my design in using the very unusual quantity of quicksilver was accomplished. That this was a true invagination there can be no doubt ; the different discharges at uncertain times, with varying quantities of the metal, until the channel of the intestine became sufficiently open, and then followed by the enormous discharges, from the doughy beds of a retained faeces ; the perfect health the man enjoyed for weeks ; the recurrence of his familiar pain in the abdomen excited by violent exertion, and the perfect occlusion of the bowel until he died, all prove this to be a true case of intussusception.

* * Should difficulty be experienced in tracing the reflections of intestines as above narrated, a glance at plate 6 in Dr. Brinton's excellent work on this accident will at once render my meaning plain.

A NEW SYMPTOM AND A NEW THEORY OF LOCOMOTOR ATAXY.

Read in the Section of Medicine.

By JULIUS ALTHAUS, M.D., M.R.C.P.,

Senior Physician to the Hospital for Epilepsy and Paralysis, Regent's Park.

THERE is no disease in the whole range of pathology which may commence in such extremely different ways as *tabes spinalis*, or *locomotor ataxy* ; and it is undoubtedly this circumstance which accounts for the mistakes which have been, and are even now, so frequently made in the diagnosis of the earlier stages of that disease. We may, at the same time, have a number of patients under care who are all in the first stage of *tabes*, and who have yet hardly a single symptom in common. One of them may complain of loss of sexual power ; another of hyperæsthesia in the back ; a third of a troublesome form of indigestion, and a sensation of tightness round the stomach ; a fourth of rheumatic or neuralgic pains ; a fifth of failure of sight, and double vision ; a sixth of a feeling of intense lassitude and exhaustion. This shows the great importance of being intimately acquainted with all the truly protean forms which this extraordinary malady may assume ; an importance which is by no means only theoretical, but chiefly practical ; for it is in the first stage of *tabes* that our therapeutical efforts are more likely to be crowned with success than at any later period in the evolution of the disease.

The principal new symptom which appears in the second stage of *tabes*, and which imparts an entirely new aspect to the malady, is that of *ataxy*. From this time forward, *ataxy* overshadows more or less all the other symptoms, with the result that patients in the second stage of *tabes* resemble one another a great deal more than in the first. The extreme differences which exist in the first or pre-ataxic stage begin now gradually to fade away, and the diagnosis is rendered proportionately easier.

I have been led, by long-continued observations on numerous cases of this disease which have happened to come under my care, to distinguish three different epochs or periods in the development of the ataxic stage of *tabes*. Such a distinction has not hitherto been made, but I believe that it will be found to considerably facilitate the study and comprehension of the different aspects which this important symptom is apt to assume at different times in the progress of the disease. These periods I will characterise as follows.

1. The initial period, in which *ataxy* is so slightly marked, that a skilled exploration is required to discover the symptom.
2. The truly ataxic stage, in which the peculiar walk known as the ataxic gait is observed ; and,
3. The period of muscular madness, in which even the typical ataxic gait is no longer possible, and muscular action, as far as it still exists, is in utter confusion.

The transition from the first to the second period of the disease is often so slow and gradual as to be almost imperceptible. Indeed, the

muscular disorder may at first be so trifling, that the patient succeeds, by slight unconscious efforts, in obviating any actual inconvenience.

At this initial period of the second stage of *tabes*, a skilled objective exploration of the patient's condition is of paramount importance. The subjective symptoms of which he most complains are often misleading and comparatively insignificant ; while the principal objective sign is often so concealed that only the specially trained observer is able to recognise and appreciate it. At this time, a patient may still be able to walk four or five miles at a time without much fatigue, and often scouts the idea that there is anything wrong with his walking powers. It is, therefore, necessary to make him go through a certain number of tests, some or all of which, when *ataxy* is present, will infallibly reveal it. The more important of them are, to watch the way in which the patient gets up from a chair or a couch ; to make him stand with his feet close together, or on one leg, and with his eyes closed ; to make him turn round quickly ; and to go downstairs ; when certain peculiarities will, in a general way, be noticed, which will betray the *ataxy*.

In addition to these tests, I will now describe another symptom, which I have not seen mentioned in any previous treatise on this disease, and which is, that the patient has a considerable difficulty in walking backwards. This faculty, which is chiefly practised and valued by courtiers, is nevertheless possessed by all ordinary mortals as long as they are in good health. For the *tabid* patient, however, it is mostly very difficult to walk backwards, at a time when he may have very little or no trouble in walking forward. His heels seem to catch the ground ; he dare not move, for fear of falling ; and, if he succeed at all in walking backwards, it is in a peculiarly halting and odd fashion, which at once attracts attention.

I noticed this symptom for the first time in the case of a gentleman, aged 24, who consulted me in February 1882, and who was, in consequence of certain official duties, obliged to walk backwards a good deal. For years this had been the easiest thing in the world for him ; but during the last six months he had been mortified to find that he experienced considerable difficulty in accomplishing this feat. He told me that his health was otherwise excellent, with the exception of "rheumatic pains" in the limbs, to which he had been subject, off and on, during the last three years. I then examined him for the patellar tendon-reflex, and found it absent in both sides. In the further course of the interview, I elicited that the patient had had syphilis ten years ago ; that he had a temporary attack of double vision three years ago ; that there was, occasionally, incontinence of urine in the morning, and habitually a feeling of numbness in the soles of the feet. The patient was still able to walk exceedingly well in the daytime and on a level road, but had found difficulty in walking in the dark. He went satisfactorily through most of the other tests, showing that he was just in the commencement of the second stage of *tabes*. The walking backwards was, indeed, the most troublesome thing he had to contend with ; and the contrast between the clumsy and awkward way in which he did this, and the apparent ease with which he walked forward, was indeed striking. When attempting to walk backwards, his heels seemed to become entangled in the carpet ; he evidently did not know how to raise his feet properly from the ground, and on one occasion he would certainly have fallen unless I had supported him. All the muscles of the thighs and legs seemed to become rigid the instant he attempted to walk backwards, while in walking forward he appeared to have no difficulty whatever in bending his knees to the proper degree.

Since then, I have made it a point to inquire about this symptom in patients suffering from *tabes*, and have found it present in the majority of cases. In an artist who is now under my care, this difficulty is particularly annoying, because it prevents him from taking a perspective of his pictures by walking backwards from his easel. In this case, there are no symptoms of *tabes* above the waist, so that the patient is able to paint as well as ever.

In difficult or doubtful cases, therefore, more especially in those at the very threshold of the ataxic stage of *tabes*, the symptom I have mentioned may put us on the proper track, and lead us to examine the patient who shows it for other symptoms of the malady. As *tabes* is still frequently confounded with gout, rheumatism, neuralgia, dyspepsia, idiopathic amaurosis, and other conditions, any addition to our means of diagnosis for that stage where the malady is not yet fully developed must be welcome, and this has been my reason for submitting it to your notice.

I now proceed to the second part of my paper, namely, to develop, as concisely as possible, the theory of the symptom of locomotor and static *ataxy* which I have, after careful consideration, been led to adopt.

Time prevents me from entering fully into the other theories which

have been put forward, and I will therefore only mention that the two principal ones which count most adherents are one first propounded by Leyden, and another given by Erb.

Leyden's theory, which has also been advocated and developed by Pierret, is that all motor disturbance which takes place in tabes, is to be explained by the influence which sensation is known to have on motion, and by the intimate relation existing between the motor and sensory tracts. He points to the fact that that portion of the cord which suffers in tabes is notoriously concerned with sensation; that the latter function always suffers in tabes, and that the affection of sensibility is, if not absolutely, at least tolerably proportionate to the degree of ataxy which may be present.

Erb, on the other hand, considers that ataxy is a true motor symptom, and not in any way connected with or dependent upon the anæsthesia, which is generally present at the same time. He finds that frequently the degree of ataxy is by no means proportionate to the degree of anæsthesia which may be found, and that complete anæsthesia, at least on one side of the body, has been seen without any ataxy at all. For these and other reasons, Erb assumes the existence in the cord of special co-ordinating centrifugal fibres, and thinks that these may be situated either in the central grey matter, or in a portion of the lateral columns, and that disease of these centrifugal fibres is followed by the appearance of locomotor ataxy.

It thus appears that the two principal representatives of German pathology of the present day differ *toto cælo* in their views of this subject, neither view seeming to explain to complete satisfaction the phenomena which are observed. A careful critical consideration of the different points in dispute seems, therefore, necessary in order to enable us to arrive at a satisfactory conclusion.

Let us, then, first, shortly consider the facts of normal and morbid anatomy and physiology which are at our disposal. We know that the posterior columns consist of two different systems, firstly, Burdach's columns, which have to be looked upon as short conducting paths; and secondly, Goll's columns, which system of fibres is made up of long conducting paths. Burdach's columns are direct continuations of posterior root-fibres, and connect the cord with peripheral parts, and thereby with external influences. They also send out numerous fibres, which proceed in various directions into the central grey matter, and which are evidently intended to connect the different segments of grey matter with each other, while others proceed upwards into the medulla oblongata, where they terminate. Goll's columns, on the other hand, are long conducting paths, which proceed from the central grey matter up to the medulla oblongata, and appear, from their anatomical peculiarities, intended to connect extra-medullary centres in the brain and cerebellum with physiologically identical fibre-systems at different levels of the cord.

It seems, *primâ facie*, reasonable to assume that the various commissures which establish a connection between extra- and intra-medullary centres, are intended to put all of these into mutual functional relations, and to form a path for producing a physiological consensus between them. Destruction of the anatomical commissures by disease would, therefore, naturally be expected to lead to a cessation of at least some degree of functional harmony.

Experimental physiology, on the other hand, has hitherto given somewhat ambiguous results. If it could be proved that the posterior columns contain centres of co-ordination, or if not centres, at least paths for the same, everything would be comfortably settled. Goltz, however, who has more particularly studied this question, has come to the conclusion that no centres for the co-ordination of complex movements exist, either in the posterior columns, nor in fact anywhere throughout the entire extent of the cord; but that such centres are to be found in the brain, more particularly in the corpora quadrigemina, the optic thalamus, and the cerebellum. Woroschiloff's experiments on the paths by which the co-ordinating impulses travel from the brain to the muscles, tend to show that, at least in the rabbit, these paths are situated in the middle third of the lateral columns of the cord, and do not touch the posterior columns at all.

The physiology of the posterior roots is somewhat better known than that of the posterior columns. If the posterior roots for one hind-leg be divided in a frog, the movements of that leg are seen to be out of harmony with the other, whether for jumping, swimming, or other modes of locomotion; and when such an animal is held between the fingers, the affected leg will not carry out the movements intended for the purpose of escaping, as is done by the non-operated leg. After experimental division of the whole of the posterior roots, a frog when put into water, is seen to be unable to swim, and to give himself up to ataxic movements.

It is important to notice that impressions which are solely furnished by the skin, are not indispensable for locomotion. Thus, Claude

Bernard has shown that a frog whose skin was entirely removed, could still swim well; while destruction of the posterior roots at once abolished that power. This fact has a distinct bearing on those cases where ataxy has been observed, together with normal sensibility in the skin. Our notions of the position of our limbs are likewise determined by the more deeply situated structures, such as muscles, ligaments, cartilages, and bones, which may lose their sensibility independently of that of the skin. It is true that in many cases of hysterical hemianæsthesia the movements of prehension, locomotion, etc., are reported to have been normal; but there are other cases in which unquestionably ataxy of movements existed under such circumstances. In a girl, aged 11, who was a short time ago under my care at the hospital, there was, on admission, hemianæsthesia of the entire left side of the body, which had apparently come on through hæmorrhage into the most posterior portions of the internal capsule at the time of birth. The affection had continued unaltered all her life, until she came under my care, but yielded nevertheless to a single application of electricity. There had been no sign of paralysis at any time; and the girl was found, on examination, to be well able to carry out all simple movements with the anæsthetic limbs which I asked her to perform. At the same time there was decided ataxy in the left hand. The girl was able to play the piano with the right hand, but could not do so with the left; she had great difficulty in picking up a pin with her left hand, and in doing any useful complex actions; yet the muscular force, as tested with the dynamometer, was quite normal. The use of the hand for finer movements improved only gradually after sensibility had been established.

The anterior cornua of the central grey matter of the cord contain the centre of muscular power and tonicity, while the volitional impulses of motion originate in the central convolutions bordering the fissure of Rolando. Co-ordination of the action of the motor nerves and muscles is, however, not affected either in the Rolandic convolutions or in the grey centre of the cord; but in the central ganglia of the brain, namely, the corpus striatum and thalamus opticus, which communicate through the white internal capsule with the higher motor centres above, and the lower centres further down. The central ganglia have the special function of rendering movements which are intimately connected with sensations, and which are, in the first instance, only excited by conscious volitional effort, mechanical and automatic. Walking and other complex movements have to be learnt early in life by countless conscious efforts on the parts of the hemispheres; and full attention is necessary in the beginning to enable us to carry out such movements in a proper manner. Eventually, however, less and less effort is required for executing such movements, which at last are performed without any attention to them on the part of the grey surface of the brain. A man who is in the habit of writing much, never thinks of the way in which he forms letters on the paper, over which his pen seems to fly quite mechanically. If each time we carry out complex movements, a special conscious effort were necessary, the time at our disposal would not suffice for a hundredth part of the work which we actually get through in life; and some forms of activity, such as finished piano- and violin-playing, would be utterly impossible.

In order, however, that the central ganglia shall be able to thus minimise the work which has to be done in life, it is necessary that they should constantly receive accurate information of the position of our limbs, and the nature of the obstacles with which the latter come in contact. We may be able to walk fast enough on a smooth level road without thinking about it; but if the pavement has been taken up, or if we have to walk across a newly ploughed field, or on the edge of a precipice, or on a narrow bridge or plank thrown across a stream, or in a crowded thoroughfare where hansom-cabs, omnibuses, perambulators, tricycles, and foot-passengers jostle each other, or in the dark on a staircase with which we are not acquainted, then a considerable amount of attention is required for overcoming obstacles in our way, and guiding ourselves along with safety. The mere impressions conveyed to the central ganglia by the posterior columns are then no longer sufficient; but the aid of the eyes, or, in the case of the dark staircase, of the hands and arms, is instinctively called in, in order to supplement the ordinary sensitive impressions by special information and manœuvring. We therefore, under those circumstances, behave like the ataxic does habitually, that is to say, we use our eyes as crutches, and manœuvre with our hands and arms to assist us; and even then we do not walk as well as we do on a smooth level road, where there are no impediments of any kind to overcome. The ataxic, therefore, is habitually in the condition in which we are under such special circumstances only as I have just mentioned. The information habitually given to the central ganglia by the posterior columns, is not available for him, because those columns have ceased

to exist, and the various groups of ganglionic cells can therefore no longer be combined for synergic, orderly, or purposive action. There is no longer any harmony between the muscles which act, and their antagonists which regulate the action; wrong groups of muscles are called into play, which impair the action instead of facilitating it; and there is therefore useless expenditure of nerve-force, causing fatigue.

The mode of production of locomotor ataxy thus appears satisfactorily explained; it only remains for us to account for the phenomena of static ataxy which are generally associated with the former in tabes.

The cerebellum, which was once believed to be the seat of the reproductive faculty and desire, is now known to be the centre of equilibration of the body. Removal of this organ in an animal causes static ataxy; the animal cannot keep steady on its legs, but staggers about as if drunk. It is not paralysed, and endeavours to carry out certain movements; but there is an utter want of precision, and even the most desperate efforts do not succeed in steadying it. The behaviour of animals deprived of their cerebellum, in fact, resembles, in the closest possible manner, that which takes place when the ataxic attempts to stand on one leg, or on both, with his eyes closed, etc. The erroneous information which the cerebellum receives from the diseased posterior columns may, however, be, to some extent, corrected by sight; and this accounts for the fact of standing being so much more difficult when the eyes are closed (Romberg's symptom), as well as for the other phenomena of static ataxy. Whether the paths through which information is given to the cerebellum are situated in Goll's columns, in which case the road would be somewhat more indirect, or in the direct cerebellar strands of the cord, which would carry information in a straight line to the cerebellum, we are at present not in a position to determine. We may, however, take it as an indisputable fact that locomotor ataxy is caused by an interruption of the paths between the posterior roots and the central ganglia of the brain, through sclerosis of the posterior columns, and that static ataxy is, in its turn, brought about by an interruption of the paths between the posterior roots and the cerebellum, through sclerosis either of Goll's columns, or of the direct cerebellar strands.

CLINICAL NOTE ON THE TREATMENT OF BURNS AND SCALDS.

Read in the Section of Surgery.

By J. WALTON BROWNE, B.A., M.D.,
Surgeon to the Belfast Royal Hospital.

DURING the past ten years, I have had a very varied experience in the treatment of burns and scalds, in dispensary, hospital, and private practice, and may say that I have tried almost every known remedy, including continuous warm baths, warm water baths with sulphate of iron in solution, white paint, in the form of carbonate of lead, the immediate application of saturated solution of sodabicharbonate, carron-oil, oakum, and the use of many ointments; deriving much benefit from an ointment composed of iodoform, extract of conium, carbolic acid, and spermacetic. From the use of these various remedies I have obtained more or less success; but the local treatment which I wish especially to bring before the notice of the Section is the application of a solution of chlorate of potash. In doing so I do not imagine I am communicating any new or original method of treatment, but merely wish to urge the use of a remedy which has proved highly beneficial in many dozen of cases.

I know that the treatment which I advocate has been tried by some of my medical friends and found wanting, but I feel confident the non-success arose simply from the fact that the lotion was not applied of the proper strength, and with sufficient frequency.

This treatment was first recommended to me by my friend Dr. Harkin of Belfast, who for upwards of thirty years has carefully studied the therapeutical effects of chlorate of potash, and who read a valuable paper upon its uses at our annual meeting in 1879. Dr. Harkin has shown that the salt exercises a most marked influence on all maladies dependent upon defective nutrition, secretion, excretion, aëration, and molecular metamorphosis. We all know its great value in diphtheritic exudation of the throat, gangrenous stomatitis, etc.; but, perhaps, among its very valuable qualities, none is more remarkable than the rapidity with which its application as a lotion repairs the injuries produced by burns and scalds. It is really marvellous how

soon the healing process commences after the application of the lotion, and how rapidly the raw surface is covered by the formation of new skin.

In the treatment of burns and scalds of the second and third degree the lotion is especially useful; in many of these cases perfect healing has taken place in four to five days. I could give the histories of many such cases.

In severe burns of the fourth or fifth degree it is also useful, seeming to favour the formation of granulations after the separation of the sloughs, and certainly no remedy which I have tried so thoroughly keeps the wounds clean and sweet.

In superficial forms, after a few applications, a zone of new skin appears at the margins of the broken surface, and proceeds towards the centre until healing is perfect. It is quite common to see a border of newly formed cuticle in the short space of twenty-four hours. The method I adopt is the following. In superficial burns, I direct the blisters to be punctured, and a bread and water poultice to be applied over all the injured surface every fourth hour, until the cuticle has become detached; you generally find this accomplished upon the removal of the second poultice. I then commence the application of the solution of chlorate of potash, five grains to the ounce; pieces of lint of the required size, saturated with the lotion, are applied four times a day. The lint is covered over with gutta-percha tissue or oiled silk; all dressings being retained *in situ* by a bandage. The lotion must not be of greater strength than five grains to the ounce; of this strength it acts as a sedative, although at the same time stimulating the capillaries to the formation of new skin. If applied much stronger it causes pain; in fact, it acts as a caustic. In some cases, where the lint occasionally adheres to the wound, and is difficult of removal, without injuring the capillaries and causing bleeding; I add to the chlorate of potash solution some glycerine, in the proportion of two ounces to the pint. To prevent the lint from adhering to the wound, Dr. Harkin has for years used an ointment composed of chlorate of potash and lard, of the strength of five grains to the ounce.

In the treatment of very deep burns I advise the application of poultices until all sloughs are separated; granulations then spring up, and nothing seems to increase their activity and vitality equal to the potash solution, of the strength of five grains to the ounce; should the granulations become weak, flabby, or too exuberant, we then use a potash-lotion of the strength of ten grains to the ounce, and at the same time prescribe a mixture of chlorate of potash and tincture of perchloride of iron, to be taken three times a day. At any time, should there be offensive discharges, one can combine the potash-solution with some permanganate of potash.

In the treatment of burns and scalds, the indications are, after promoting reaction and calming the system: 1, to limit resulting inflammation; 2, to promote the sloughing process; 3, to favour the development of granulations; 4, to moderate contraction; and 5, to sustain the strength of the patient during the wasting effects consequent upon the protracted suffering which so often occurs.

I submit that all these indications are fulfilled by the local and internal uses of the salt, the beneficial effects of which, in the treatment of burns and scalds, I have had the honour of bringing before your notice.

ON THE METHODS OF DEMONSTRATING THE PHYSIOLOGICAL ACTIONS OF DRUGS ON THE FROG'S HEART.

Read in the Section of Pharmacology and Therapeutics.

By DUDLEY W. BUXTON, M.D., B.S., M.R.C.P.,
Assistant to the Professor of Medicine in University College, London.

It is my duty to-day to bring before you some of the many striking points which the field of practical therapeutics brings before the explorer in physiological research. As the time allotted to my demonstration is necessarily short, I have decided to bring under your notice only a few experiments, and explain briefly the method I propose to employ. It has appeared to me most consonant with your interest that I should select illustrations of the methods from most striking instances of antagonism and mutual co-operation of salts, instances which will evolve themselves in a few moments, rather than to attempt to introduce to you, in the short space of time at my disposal, any original and so disputatious communication.

Dr. Sydney Ringer, in the *Journal of Physiology*, has brought before the world the remarkable behaviour of the various saline constituents