Dr. McVAIL, in reply, said that, at that advanced hour of the meeting, he would say only a very few words. He in no sense, at that meeting, sought to controvert the view that, under certain circumstances, the strong expiratory act of coughing might give rise to emphysematous dilatation of the cells, and he would therefore pass over all that had been said by speakers on that point. Also he had distinctly guarded himself from any remarks on the nutritional causes of emphysema, as time had not permitted the extended explanation of their share in the production of emphysema, that was desirable. As, however, nearly all the speakers had alluded to this side of the question, he would say that he was convinced that nutritional, or trophic emphysema, apart altogether from bronchitis, was exceedingly common. He found that young men, who came to the dispensary of the Western Infirmary of Glasgow, were peculiarly subject to it: young men from ship-building yards, and other public works, in Glasgow, who had begun, at a very early age, to drink whisky on pay-nights, and who, at a very little over the age of twenty years, had, in many cases, already lost as much as a quarter, or a third, of their breath-reserve space, and whose arteries also were becoming zigzagged from impairment of their elastic tissue. In fact, this question of trophic emphysema was one largely affecting life-assurance, and all questions of health-prognosis; and its full significance had yet to be wrought out by the aid of such instruments as the spiograph that he showed them. It was because this question was so vast, that he had almost entirely left it out of view in the remarks with which he opened the discussion. He hoped, however, shortly to bring a more extended account of it before the profession. He could not understand how Dr. Gairdner could object to the view that prolonged breathing, at a high respiratory level, would give rise to vesicular dilatation. If the collapse of, for example, four cubic inches of lung-tissue, must necessarily, in Dr. Gairdner's opinion, cause a corresponding dilatation, distributed less or more evenly among the other cells of the lung, surely an extra dilatation of from fifty to two hundred cubic inches, brought about and maintained by any cause, no matter what, would much more give rise to cellular emphysema. Dr. Ewart had objected to the explanation given of the mode in which the dilating force acted on the air-cells; but, as to that, there could be no doubt at all; the distending force was from without, and, for every successive layer of cells acted upon, the outer layers must successively undergo an extra unit of strain, just as the addition of an extra wagon to a railway-train, going up a hill, added an unit of tension to all the couplings between it and the engine.

ON THE OPERATION (ALEXANDER - ADAMS) OF SHORTENING THE ROUND LIGAMENTS FOR UTERINE DISPLACEMENTS, WITH THREE CASES.

Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association.

By W. L. REID, M.D.,

Physician to the Dispensary for Diseases of Women, Western Infirmary;
Physician to the Maternity Hospital, Glasgow.

FROM my friend Dr. James A. Adams, of Glasgow, I had heard of his operation for replacing displaced uteri a considerable time before I knew it had been proposed and practised by Dr. William Alexander, of Liverpool; but I did not attempt the operation until I had seen the latter gentleman perform it on the dead body, and had heard particularly of his results in the living subject. Although, from the position I occupy, I had abundant opportunities, it seemed right to wait until I met with what appeared to be test-cases, in women who were willing to have practised upon them an avowedly new and so far untried operation; and it is to the histories and results of three such cases that I now direct your attention. Possibly, some little extra interest may attach to them, in so far as, to the best of my knowledge, they are the three first successful cases operated on in Scotland, speaking of them simply as operations for replacing displaced uteri.

CASE I.—Mrs. McG., aged 33, began to menstruate at 16, and was regular till pregnancy, two months after marriage, at 20. She had one child naturally, and had never been well since. The uterus was found much hypertrophied and retroflexed. The left ovary was prolapsed. She complained greatly of backache, pain in the left ovarian region, and inability to walk any distance. There was great and continuous leucorrhœa; and intercourse was almost unbearable. Dysmenorrhœa was severe, and there was frequent micturition, especially during the night. These symptoms had continued more or less

steadily for ten years. She was treated by me at the Western Infirmary for nearly two years, during which time she obtained temporary relief from pessaries, glycerine tampons, scarification of the cervix, and intra-uterine applications of iodine, and of nitric and carbolic acids.

On November 1st, 1883, assisted by Drs. James A. Adams, George A. Turner, and James Ross, I performed the operation of shortening the ligaments. The pubes was shaved and washed with carbolic water, and carbolic spray was used during the operation. An incision was made on the left side from the pubic spine upwards and outwards for two inches, and carried down to the tendon of the external oblique muscle. The external abdominal ring was felt, and the tissues coming out of it were gathered up by an aneurysm-needle, and seized with the fingers. They could not be drawn out, even by the exercise of considerable force. They were then grasped by a pair of broad-pointed dressing-forceps, and freed from the surrounding tissues with the help of the scalpel; and, on continued pulling, they very gradually came out, until the firm round ligament appeared. The same procedure was repeated on the opposite side. A sound was then passed into the uterus, and the organ braced well up in its normal position, and the ligaments were pulled on until they were found to be fixing it there, when fully two inches were found drawn out on each side. Three strong chromicised catgut stitches were passed through the pillars of the ring, each transfixing the cord. These were not tied very tightly. The ends of the cord were folded up in the wound, and the edges of the skin brought together by gut-stitches over a rubber drainage-tube. To take the strain off the ligaments, a pessary which the patient had worn before was introduced. A dressing of gauze and cotton-wool was applied, and kept in place by an elastic bandage. In the evening, the pulse was 90, and the temperature 100.2°. The patient complained only of something being "very tight" in the lower part of the abdomen. The wounds were dressed without spray, with the idea that the scar-tissue would be firmer if suppuration were allowed to take place. It would serve no good purpose to give careful details of her progress; suffice it to say, that the wounds were dressed twice a day for four days, and once a day thereafter, until she was dismissed on the twenty-sixth day, there being then only a corner of one of the wounds unhealed. Her highest temperature was on the second day, namely, 101.4°; on the third day, it fell to 98.6°.

At the time of her dismissal, she complained of frequent desire to pass urine when standing or walking, but this soon passed off. The pessary was left in position, and was not removed until a month afterwards.

On April 21st, six months after the operation, I saw and examined her. The cervix was found high up, and normal in position. The fundus could not be felt in the posterior vaginal fornix, but the point of the sound passed backwards, showing that the organ still remained flexed.

On July 23rd, when asked to what extent she had improved since the operation, the patient mentioned the following points. The dysmenorrhœa and dyspareunia were both gone; she did not require to micturate during the night; defæcation was much more easy; and the weakness of her legs was so much improved, that she could walk two miles without special fatigue. On the other hand, she still complained of backache and ovarian pain; the uterus was still three and a quarter inches in depth; there was erosion of the cervix, and a mucopurulent discharge from the os uteri. I may add that, since the operation, the ovary has never come within reach.

CASE II.—Mrs. H., aged 31, was first seen by me on September 4th, 1880, when the following note was taken. She began to menstruate at 10, and was regular, with some dysmenorrhœa on the first day, until pregnancy, two months after marriage, at 21. She menstruated one month after her labour, which was normal, except that the second stage was long. Milk died away when menstruation appeared. She had been quite regular, but never felt well since her confinement. She complained of backache, leucorrhœa, frequent micturition, and of a sharp attack of diarrhœa, which came on every morning immediately after rising, and did not return during the rest of the twenty-four hours. She had no ordinary dysmenorrhœa, but suffered severely from headache before and during each menstrual period. This symptom was often so severe as to be accompanied by delirium, and to confine her to bed for three or four days. The patient was stout and florid. The uterus was found retroverted, three and a half inches in depth, very heavy. The cervix was hypertrophied, and patulous. The perineum was partially destroyed, and much relaxed.

Her history for the next three years may be summed up in saying that, by the use of the means mentioned in connection with the case already detailed, she obtained at various times great relief, which, however, never lasted more than a few weeks; and, on November 3rd,

1883, her ligaments were shortened. The operation was like the former, except that the left ligament broke while being pulled out, and the canal had to be laid open to the extent of about half an inch before it could be again caught up. In this case, except during the operation, the spray was not used. Pus appeared on the fourth day; and by degrees the slack of the ligaments sloughed away, as, indeed, they also did in the first and third cases. The discharge was never great; and the patient was sent home on the twenty-third day after operation, she having been more or less out of bed for some days previously.

The note taken a fortnight afterwards is as follows. The patient had been going about freely for the past week, and had menstruated without headache, or any disturbance whatever. She stated that she felt perfectly well. The uterus was found somewhat anteverted; the cervix high up, and the whole organ lying nearly horizontally across the cavity of the pelvis, yet easily movable. This condition of comfort continued for nearly three months, when many of her old symptoms reappeared. On July 2nd the following note was taken. The uterus was in the position formerly described as existing after the operation; the cervix was smaller and softer; the cavity was three inches in depth, and considerable leucorrhœa existed. The patient complained of backache, and pain in the left ovarian region, on much exertion, and had occasional attacks of morning diarrhœa. When questioned as to improvement, she unwillingly admitted that she had less diarrhœa, much less headache, was much more able to walk freely, and the bladder-trouble was gone. She had become very stout, but thought herself far from being well.

CASE III.—Mrs. M., aged 45, began to menstruate at 18, and was regular and natural in her periods, until pregnancy, one month after marriage, at 25. She had four children naturally, the last thirteen years ago. She did not recover well from this confinement, and ever since suffered from profuse leucorrhœa, backache, bearing down, and inability to walk. Ten years ago retroversion was diagnosed, and all sorts of pessaries and other forms of treatment were tried by one medical man after another, without any permanent relief. For the past year she had been almost entirely confined to bed. I saw her first on October 20th, 1883, and made the following note of the condition of the pelvic organs. The whole uterus was much hypertrophied, its position was that of exaggerated retroversion, the fundus lying, as it were, on the perineum. The patient complained of great backache, and bearing down, whenever she was more than a few minutes in the erect attitude, and of great difficulty and pain during defecation. The uterus was 3½ inches in depth, and was restored to the normal position with difficulty and pain. There were no bladder-symptoms. The patient was very thin and haggard, and took little or no food. She was watched and treated for some months by Dr. Turner and myself, all the means we could think of being tried without more than passing benefit. Most pessaries were gradually borne down by the heavy uterus; those which continued to sustain it becoming unbearable by the patient. Means used to lessen the bulk and congestion of the organ had no permanently good effect. She was operated on on February 20th last; and, although the spray was used during the operation and at every subsequent dressing, and the wounds remained quite sweet, the slack of the ligaments sloughed on both sides, preventing the wounds from healing for nearly six weeks. Otherwise, a good recovery was made. After being six weeks in the country, she came back to town, and was examined on June 25th. She was then able to walk pretty freely and be out of bed all day, was looking much better, and had little or none of the great mental depression from which she formerly suffered. Defecation was quite normal; but she had a rather frequent desire to micturate; not, however, accompanied by pain during the act. The uterus was normal in position, with a tendency to anteversion, 3 inches in depth, not tender, and moderately movable. There was a fold of loose tissue in the posterior vaginal fornix. No menstruation had occurred since two months before the operation. It may be mentioned that this patient continued to wear a large Hodge's pessary for three months after the replacement, owing to a fear on her part that, without it, her old symptoms might return.

It is somewhat difficult to arrive at a sound conclusion in regard to new operative means for the relief of disease. Men are apt to range themselves for and against a new method, and give their experiences shaded or brightened by preconceived notions on the subject. In giving the details of the above cases, I have tried to make plain unvarnished statements, and leave you to draw your own conclusions from them. Permit me, in closing, to advert to one or two points in connection with the operation, drawn from my personal experience, and state my own belief in regard to its results.

1. It is better to use the spray, and to keep the wound aseptic until it has wholly or nearly healed. There is hardly any hope of its healing

by first intention; but, if kept aseptic, it heals more quickly than it would otherwise do.

2. The loose part of the cord should not be folded up in the wound, but the greater part of it cut off, so as to avoid the presence of sloughing tissue in the wound. Properly secured by sound stitches, and with the uterus supported by a good pessary, there is no danger of the cut ends being drawn into the abdomen, they being soon secured by adhesion.

3. The ends of the ligaments can be most easily reached by standing on the side on which you are operating, as the outer edge of the ring is more easily felt by the finger when one stands on the same side.

4. Contrary to Dr. Alexander's recommendation, I have found that the end of the round ligament is more easily isolated by grasping it with a broad pair of dressing-forceps. The cross fibres which prevent it from running out can then be cut or torn gently away, one by one, until it begins to come out freely. In my second case, where the fingers were used, the ligament broke, and the canal had to be opened up before it could be again seized.

5. In this region, an elastic bandage put round the waist and thighs will be found more effective in keeping the wound aseptic than a large dressing with an ordinary or flannel bandage, as the former adapts itself to the varying position of the limbs, and prevents air from being drawn under the dressings.

Referring to the results of these cases only, it seems to me that we cannot promise much in the way of certain and immediate relief from this operation. It remedies the position, but not the condition, of the uterus. I caught at it because, although often using severe means, I have been very unsuccessful in curing chronic hypertrophy of the womb, especially when accompanied, as it so often is, by some form of displacement. The results of this operation, as I have given them, although far from brilliant, are such as to lead me to continue practising it in obstinate cases, in the hope that, the malposition being remedied, the disordered condition may be more readily subdued.

A CASE OF ALEXANDER'S OPERATION OF SHORTENING THE ROUND LIGAMENTS IN A CASE OF AGGRAVATED PROLAPSE AND RETROFLEXION.

Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association.

By GEORGE ELDER, M.D.,

Surgeon to the Hospital for Women, Nottingham.

THE patient upon whom this operation was performed, has been known to me, since 1872, as an almost constant attendant at one or other of our local medical charities; and latterly, as spending most of her time either as an in-patient of one of these institutions, or at home, confined to bed.

Her history is briefly as follows. Prior to the age of 36 years, her health had always been good, and her ability to do her share of hard work unimpaired. Of this she had more than usually falls to the lot of working men's wives, as her husband's drinking habits compelled her to work enough for both.

Whilst lifting a heavy weight, she felt something in "her inside give way;" and, from that time till date of operation, her life had been spent as previously mentioned. In addition to the ordinary discomforts arising from a heavy retroflected and prolapsed uterus, aggravated by the effects of a badly ruptured perineum, her most urgent complaint was one of incontinence of urine, amounting to constant dribbling when in the erect position. Her life was rendered miserable by this state of matters, and to remedy it many and varied forms of treatment were used.

For nine months at a stretch, she lay in bed in one of our hospitals; so that it cannot be said, at least, in this instance, that whatever good the operation has done, might fairly be attributed to the necessary lying in bed.

Hot-water injections, position, pessaries of all kinds, intra-uterine stems, medicated tampons, with the administration, internally, of appropriate remedies, were, singly and in combination, tried, with little permanent good effect; and it was when almost despairing of anything short of extirpation of the peccant organ being of any service, that she came under my notice again last February. My attention had been drawn to Alexander's operation by reading a paper by Mr. Lediard, of Carlisle, and the case under notice seemed a crucial one to test it.

On March 15th I cut down upon both round ligaments, finding no difficulty in reaching and separating them from surrounding tissue, and,