of his admission, there was no discharge from the nostrils. The whole nose had been destroyed, and the nasal passages were converted into a wide chasm exposing the pharynx. There was, in addition, a large perforation of the hard palate, through which the food escaped during mastication. The patient positively stated that he had never had syphilis.

The treatment commenced by closing the hard palate with an obturator and plate of black rubber; and I am much indebted to Mr. W. H. Kirton, the surgeon-dentist of the hospital, for the great care and skill he bestowed upon this important preliminary proceeding. The first operation was done on April 4th, and, three weeks afterwards, the new nose was completed by the addition of the columna. The large wound on the forehead healed rapidly in a month. The engravings—taken from photographs—represent the patient before and after restoration of the nose.

REMARKS.—Rhino-plastic surgery often demands some modification of detail to suit the special features of the case. In this instance the entire nose had been destroyed, and in the place of the organ there existed only a depressed and ugly scar. The first step in the operation consisted in paring the edges of the old stump, and in forming a deep groove for the flap, the nasal cavity being carefully plugged with sponge, to prevent the entrance of blood into the air-passages. By proceeding in this way, much time was saved, for the free bleeding gradually ceased, and the raw surfaces were effectually glazed over during the interval occupied with the dissection of the flap. The incision over the nasal bones extended upwards to the exposed surface on the forehead, and, without any trimming, the soft parts were freely separated on each side. The flap, four inches in diameter, was taken obliquely from the centre and left part of the forehead, and on the right a broad root was left to support it. The twist of the pedicle was then very gradually made, and its neck fitted into the groove over the nasal bones. To prevent manipulation with the fingers, two sutures were at once inserted into the corners of the flap, and, by the aid of these threads, it was readily turned into its new position. Text-books generally recommend the division of the root of the nose, as soon as the adhesions are firm and the vascularity of the part established. In this case, however, there was no root to divide; this arose partly from the method of operating, and partly from the cicatricial depression caused by the destruction of the organ.

PRESENTATION.—Dr. Dodds, who has been appointed Deputy-Superintendent of the Montrose Asylum, was presented by the officers and staff of the Birmingham Borough Asylum, on the 28th ult., with a handsome timepiece. The presentation was made, on behalf of the subscribers, by the Superintendent, who spoke in the highest terms of Dr. Dodds. The timepiece was of black marble, in the shape of a temple, striking hours and half hours on a gong.

EXTENSIVE BURN INVOLVING THE CAVITY OF THE KNEE-JOINT.

Read in the Section of Surgery at the Annual Meeting of the British Medical Association.

By W. H. DALY, M.D., Pittsburgh, Pennsylvania, U.S.A.

THERE are isolated cases in surgery, occurring now and then, which, from their peculiar history, furnish us striking proof that the time-honoured indications for amputation may be ignored, and the patient still recover with a limb nearly as useful as before it was injured.

The case upon which my remarks will be made, is one that I now have much pleasure in presenting to you, after the lapse of five years, which has alike tested the usefulness of the limb and the satisfaction of its possessor. The history is briefly as follows.

H. L. K., aged 41, married, native of Pennsylvania; occupation, a locomotive engineer; weight, 212 pounds; and powerfully built.

On the night of October 21st, 1879, while he was driving his engine at a high rate of speed, with a heavily loaded train of freight cars, on a down grade, a cow was struck, derailing the engine, precipitating it, and many of the cars, down over a high embankment into a narrow ravine, the freight-cars being piled promiscuously upon and about the engine to a considerable number.

One of the brakemen was killed outright, and the other trainmen were more or less injured. When aid arrived at the scene of accident, the cries of the driver were heard beneath the mass of ruined cars, where he was fastened beneath the heated fire-box of the engine. It was found impossible to extricate him from his perilous and agonising situation for three hours and twenty minutes; and, during this time, the fire was fiercely burning under the boiler of the engine, and the iron box containing it was resting, with all the superincumbent weight, upon his limbs, fastening them firmly in the soft earth upon which they rested. It can easily be imagined that the cooking process had ample time to be complete throughout its extent.

Some of the wrecking gang let down stimulants to him, of which he drank judiciously, though freely, preventing collapse. When the mass of wrecked cars was cleared away, the engine was jacked up, and the unfortunate man released.

As the company's general surgeon, I was dispatched, with some of the staff, on a special engine to the scene of accident, and found the patient, with all the anterior and lateral surface of the left leg, about and below the knee, a mass of charred black, grey, and brownish tissue. With a probe, the tibia and fibula could be felt, surrounded by the mass of cooked flesh. The patellar ligament was completely burned from its attachment, allowing the patella to be raised up, and the knee-joint cavity looked into, the full width of this bone.

The superficial area of destroyed tissue was estimated at from 85 to 90 square inches about and below the left knee, with a depth nearly equal to half the antero-posterior diameter of the leg at the knee; from this point of greatest depth, it shaded off gradually downwards anteriorly, to a mere superficial lesion, eleven inches below the knee.

The patient was removed to his home, thirty miles distant, having positively declined hospital care and treatment.

The leg was placed in a fracture-box, and the wound well saturated with carbolic linseed-oil, in the proportion of one drachm of carbolic acid to the pint; a warm flaxseed-poultice covered this and the leg, kept *in situ* by being surrounded within the fracture-box with bran.

The dressings were changed twice in the twenty-four hours, and all kept as sweet and clean as possible. Good nourishing diet, with whisky, and occasionally a dose of morphia, was given him.

Traumatic fever ran high during the first six weeks. The dead soft tissues separating, left the bones (tibia and fibula) standing out in bold relief upon the copiously suppurating surface, and the knee-joint cavity open in its full extent anteriorly, from side to side.

The poultices were continued, but the fracture-box was superseded by the use of a sheet-iron splint to the posterior aspect of the leg; the splint was made to fit accurately, from the glutei to the foot, and was well padded; in this, the limb was suspended at a comfortable elevation.

The question of amputation was freely discussed and advocated by about twenty or more surgeons who were called in consultation during the progress of the case. Not considering it prudent to risk an amputation, I dissented persistently, in the belief that a fairly useful limb could be secured, all of the strongly expressed differences of opinion in nowise involving any breach of the most friendly relations of all concerned.

The dead bones not separating as rapidly as I desired, on January 15th, 1879, with mallet and very sharp chisels, I cut away, by piece-meal, a segment equal to half the antero-posterior diameter of the head of the tibia, carrying the removal of all the bone down the shaft to a point about eight inches below the joint; the head of the fibula, and nearly all of its upper third, I took away *en masse*. I took away a part of the first metatarsal and internal cuneiform bones of the right foot, which had also been burned at the time of the accident.

At the expiration of ten months after the accident, the lesion was nearly healed, being much facilitated by skin-grafting, as many as 50 to 100 grafts being put in at once, at intervals. A few weeks later, he rode out, and soon began to walk with the leg supported in a per-

fectly straight position by a splint, as giving the greatest amount of coaptating surface to become ankylosed, this being our only hope for an useful limb, since all the extensor muscles had been burned away.

The man is before you now, in a photograph, taken twenty-seven months after the injury. He is still, as you see him represented there, hale and hearty; and, while the case, which must be regarded as unique, could have been presented to the profession at any time during these four years past, I desired to wait a reasonable period, to

ascertain what were to be the permanent results as to an useful leg. He has a better limb than any artificial one that can be made and adjusted to a stump by anyone, and, to use his own words, he "would not give his leg for all the artificial legs in the world." He can go up and down stairs with ease, carrying anything that ordinary men can carry. He can walk unaided by either crutch or cane. The knee-joint is solidly ankylosed, and there is not the slightest doubt that this patient will, for many years to come, possess capabilities as a pedestrian that will put to the test the best walker among the twenty and more professional friends who saw him in consultation with me during the course of treatment of the injury, and who honestly desired, without a single exception, to give him a good stump for an artificial limb.

Be that as it may, the patient is satisfied, and I am quite well satisfied, with the results of the most persistent piece of work that I have ever done in connection with conservative surgery; and, for the satisfaction of the large number of my friends who so honestly differed from me, and expressed the opinion that amputation should be performed in accordance with all known rules of surgery, I quote Bryant (vol. ii, see American ed., p. 316), who says: "Extensive wounds of large joints, complicated with other injuries, had better be treated by amputation or excision;" and there is no lack of further high authority for the position my friends assumed in this case with reference to the question of amputation.

Druitt says that extensive burns, even of small severity, are always dangerous; and Gross says with reference to burns that, where extent of surface and depth are combined, death will occur without reaction.

In an interview with this patient two months ago, he informed me he was in the habit of walking over to the city of South Pittsburgh and back to his home, a distance of three miles, and that his leg felt better for the exercise.

A short time ago, I found him engaged in painting the front of his three story house, climbing in and out of windows; and he is about to take a situation again as locomotive engineer.

He has an useful leg that is ankylosed in the straight position, and which is capable of carrying him anywhere, and of enabling him to earn a livelihood in nearly any avocation in life.

Altogether, gentlemen, this is one of those rare instances in surgery when all the established rules for amputation have been ignored, and the best of consequences have resulted, notwithstanding.

KNEE-JERK: ITS CAUSE, NOT REFLEX. ✓

By JAMES OLIVER, M.B. Edin., M.R.C.P. Lond.

THE causation of knee-jerk, and of the allied phenomena produced by a sharp tap upon certain tendinous expansions, is still a debatable question. Numerically, the supporters of the reflex theory outweigh those who resent this as a favourable explanation.

Presupposing the reflex hypothesis feasible, every one will admit that integrity of the arc is essential for the production of the phenomenon, any breach occurring in the continuity of the circle causing inevitable failure. To a certain extent I admit a subsidiary influence of the arc, but cannot believe the condition to be one truly reflex in character.

Three factors are apparently necessary for the production of the tendon-phenomenon: 1, a certain amount of true muscular fibre; 2, muscular harmony; 3, muscular tone.

If there be not sufficient true muscular fibre existing, no jerk will be obtainable. Mechanical stimulation may cause the muscular fibres irritated to contract; the patient even may have sufficient power over the vasti muscles to be able to maintain the leg and thigh fully extended for a very short period; and yet the knee-phenomenon may be in abeyance, and this simply because there is deficient true muscular structure. It does not, however, always follow that a defective state of the muscle causes abolition of the tendon-phenomenon; for, associated with this deficiency, there may be an increased tonicity and irritability sufficient to counteract, more or less, the muscle-loss.

It has been adduced in favour of the reflex theory, that loss of knee-jerk follows division of the corresponding motor or sensory roots. The explanation, however, may be sought elsewhere than under the cloak of a reflex action. If the posterior roots alone be divided at any level in the spinal cord, certain definite changes are noticeable in the movements of the muscles in the area supplied by the severed nerve; they become less orderly; that is, there is interference with muscle-harmony. Now, knee-jerk is more or less dependent upon the harmonious action of the muscle; interference, therefore, with the latter may cause cessa-