ADDRESS IN SURGERY.

Delivered at the Annual Meeting of the British Medical Association in Cardiff.

BY

J. MARSHALL, F.R.S., F.R.C.S.,

Emeritus Professor of Surgery, University College, London; Consulting Surgeon to University College Hospital.

A FORTY YEARS' RETROSPECT.

MR. PRESIDENT AND GENTLEMEN,—Assuming that on such an occasion as the present my discretion is entirely unfettered, and feeling anxious to avoid wearying my audience with a formal essay, I have chosen for the subject of the Address in Surgery, which I have the honour of delivering here to-day, a comparative view of surgical practice, as I myself observed it, when acting as dresser under Robert Liston in the wards of University College Hospital, and of such as now prevails amongst us, particularly as exemplified in my own wards in that institution.

I at first contemplated, for the purpose of this comparison, the selection of the years 1843 and 1883, moving backwards from the last named date, because it coincides with that of the latest of our surgical registrar's reports, compiled by my former house-surgeon, Mr. Victor Horsley. But, on searching the old case-books, I found that, in the medical year 1844-5, I was myself one of Liston's dressers, and accordingly I resolved to compare that year with 1883, which two years, in fact, constitute the first and last of a series of forty. I soon found, also, this advantage, that the perusal of my own descriptive notes, and those of my contemporaries, brought back vivid remembrances of the more important cases then under treatment.

It gave me great pleasure to discover that two of my old and yet surviving friends, James Hakes, of Liverpool, and George Yeoman Heath, of Newcastle, were acting as Liston's house-surgeons in the year selected; and that amongst my fellow-dressers were Henry Wiglesworth, of Ashford; John Newton, of Liverpool; the late Thomas Atchison, of the Indian Medical Service; F. W. Marshall, of Brundall, near Norwich; and Wm. Cadge, of that city. It is with equal pleasure that I mention here the excellence and value of their reports.

Whilst thus limiting my comparison as to period and locality, and to events more or less within my own cognisance, I hope to be able to direct your attention to numerous points of practical interest, whilst occasional generalisations need not be excluded.

I propose to take up, in order, the following subjects: 1. The numbers of the cases, and the character of the diseases and injuries treated in the two selected periods; 2. The methods and means of investigation employed, whether clinical or pathological, for the purposes of diagnosis; 3. The treatment of the cases, understood in its widest sense, hygienic, dietetic, and curative; 4. The results obtained, as indicated by such factors as the duration of the treatment in the wards, the progress of the cases, and the relative mortality.

In dealing with these questions, either exact figures are given, or, when general statements are introduced, they are founded on the best attainable data.

1. *The Numbers of Cases, and the Character of the Diseases and Injuries to be Compared.*—In the year 1844-5, 284 patients were admitted into Liston's wards (excluding eye-cases); of these, 196 were males, and 88 were females. In 1883, there entered my wards 396 patients, of whom 223 were males, and 171 females. Liston's cases constituted about one-half, whilst my own formed only about one-third of the total number of patients received into the surgical wards.

In both sets of cases, fractures of various kinds, chiefly of the lower limbs, reached a very high number; next to these, especially in the earlier series, were wounds of the soft parts, chiefly affecting the head, neck, and upper limb. Burns and scalds were few in each period. Malformations of the lips, palate, fingers, and foot, were more numerous at the earlier date; but, as might be expected, genu valgum is not mentioned. New growths, though actually more numerous, were not relatively so in the later period, the proportions of these to the total number of cases being about $7\frac{1}{2}$ per cent. in 1844-5, and only 7 per cent. in 1883. There is, however, a marked increase in the epitheliomata, though scirrhus and sarcoma remain the same; whilst fatty tumours are more numerous in the earlier period. Of course, the numbers contrasted, 21 and 26, are too small to furnish evidence either way as to the commonly supposed increased prevalence of cancer amongst our existing population. The widespread and well

[1284]

grounded belief in the value of the early removal of all new growths, and the more ready submission of patients to operations now rendered painless by anesthetics, must here be taken into account, as tending to swell the later numbers. Cases of venereal disease were formerly much more freely admitted into the wards of a general hospital than now, and, accordingly, are more numerous at the earlier date. Ordinary abscesses and ulcers are equally represented at the two periods. Gangrene was, and is, rare. Diseases of the joints, and also diseases of bone, are abundantly represented in each set of records; but, in the later period, cases of caries particularly abound, probably owing to the greater success of modern operative interference. *Fistula in ano* and stricture of the rectum are mentioned in both the early and the later lists; but it is remarkable that not a single instance of operation for hemorrhoids occurred in Liston's wards in the year in question. Diseases of the rectum, however, were at that time marked subjects of the specialism founded by Van Butchell. Strangulated hernia, calculus, stricture of the urethra, and affections of the testicle, are met with, though in varying proportions in both lists; but cases of stricture especially are now more frequently admitted as in-patients for the purposes of the so-called radical cure. A single case of ovarian tumour, one of vesico-vaginal fistula, and one of aneurysm, occur in Liston's set of cases; but the last named disease is now comparatively rare, and the two former evils are usually relegated to special hospitals. The eye cases, which were received into Liston's wards, included strabismus, staphyloma, and cataract, for all of which he operated successfully. It would be a mere parade of knowledge to enumerate the numerous modern operations, such as osteotomy, colectomy, nephrectomy, and many others, which, of course, did not occur in Liston's experience.

As to the various forms of specific inflammations, and their consequences, which are now classed together as septic diseases, and which have long been the bane of surgical, and especially of hospital surgical practice, it is well known that they were not so clearly differentiated in 1844-5 as they are at the present time; but it is evident, from a study of the hospital records of the older date, that they were then extremely prevalent. Indeed, the chief lessons to be drawn from the facts I have undertaken to review relate to this class of diseases.

2. *Comparison of the Methods and Means of Investigating Surgical Disease Forty Years since and at the Present Date.*—In 1844-5, the finger, the educated finger, passed down the throat, or up behind the soft palate, was used to explore parts now examined by the laryngoscope or pharyngoscope. The convenient specula, which we now possess for various internal examinations, were then but rudely represented. There was no endoscope, and, above all, no ophthalmoscope. The place of the modern aspirating syringe was but imperfectly supplied by the exploring needle or trocar. Means were not then devised for measuring variations in the corpuscular and chemical elements of the blood in disease and in convalescent states. The presence of albumen and sugar in urine was, no doubt, easily and commonly detected; but quantitative determinations of those substances were troublesome and difficult to make, or were left unattempted; and the same is true in regard to such normal urinary constituents as urea and the chlorides, sulphates, and phosphates.

The clinical thermometer was not in use; yet, truth to say, it had been suggested as far back as the 17th century by Sanctorius, and was actually employed as a measurer of the strength of fever, by De Haen and Dr. James Currie, at the end of the last century. But, though the thermometer was not employed in Liston's time for ascertaining the temperature of the body generally, it is interesting to note that, in a case in which he tied the external iliac artery for aneurysm, exact observations on the temperature of the two lower limbs were made by aid of that instrument.

In the absence of this now ubiquitous clinical registrar, on the indications of which we justly set so high an estimate, great care was then taken to note other signs of the febrile condition. Hence, in Liston's case-books, are to be found constant and minute records of the state of the skin and tongue; of that of the pulse and breathing; of the occurrence of rigors and sweatings; of the supervention of delirium; of the quantity, character, and specific gravity of the urine; and of the occurrence of albumen in that fluid. In these respects, the notes upon the cases are for the most part perfect. There are also occasional stethoscopic observations; but there are no sphygmographic pulse-tracings. The elaborate temperature-charts and diagrams of the present case-books are necessarily absent; so, likewise, the electric and other tests, applicable to the detection of diseased conditions of the muscular and nervous systems, are entirely omitted.

Forty years ago, the microscope had just come into use for pathological, and therefore for clinical, observation. Liston's intimacy with Dalrymple, Gulliver, and Kiernan had its scientific as well as its

social aide, and served to stimulate his interest in microscopic investigations. He possessed one of the best and most powerful instruments of that day; and, as I know from having had the privilege of assisting him, he was particularly interested in the character and mode of growth of new vessels in granulations and elsewhere, in the structure of tumours, the constituents of cancer-juice, the seat of the pigment-granules in melanotic tumours, the presence of spermatozoa in the fluid of certain hydroceles, and in many other facts of micro-pathology. The researches of Müller were then becoming familiar; but the further labours of Schwann and Schleiden, of Addison and Augustus Waller, of Virchow and Cohnheim, and of many others, had to be made generally known before the grand microscopes of the few were supplemented by the clinical microscopes of the many, which now, indeed, are in daily use in every hospital-ward. Liston's *Elements of Surgery* (last edition, 1840), though beautifully illustrated by Wm. Bagge, contains only one solitary woodcut devoted to a microscopic object—namely, a cluster of red blood-corpuscles, represented upon a scale of squares, without a single white blood-corpuscle amongst them. What a contrast to the countless illustrations of minute structures, organisms, and crystals, scattered through the pages of the modern textbook!

It is not surprising that the older case-books contain no account of the microscopic structure of morbid growths, or of the minute constituents and deposits of diseased fluids—much less any illustrative sketches of either; whereas, in the modern case-records of 1883, pen-and-ink drawings constantly occur, more especially, it must be admitted, representative of the larger facts of size, shape, and other obvious characters. Formerly, we find, in the notes, a mere statement that such and such a growth “exhibited the usual characters” of a fatty, sarcomatous, or scirrhus tumour; now, those characters, when necessary, are fully described, or, it may be, actually figured. Permit me here to emphasise the importance to the medical student of the art of drawing accurately, and to express my gratification at the contrast between the useful sketches which serve here and there to illustrate the pages of my own case-books, and the funny caricatures of the features of my old surgical Master which decorate the covers of his.

As a natural consequence of the improved methods and means of pathological and clinical observation which we now possess, the notes by the clinical clerks have become more ample. Indeed, I find that the average space now allotted to each patient's history and record is twice what it formerly was; and this is independent of the temperature chart or charts now affixed to every case, of occasional diagrammatic curves, and of certain hygienic tables, in which are recorded the septic or aseptic condition of every wound in a ward, and the concurrence or absence of serious infective disease. Nevertheless, as I have already hinted, the older notes are usually full and intelligent in all particulars to which they relate; and, as the duties of dresser and clerk, now assigned to different students, were, forty years since, performed by the same person, the observation and the description of characters and symptoms were the work of the same mind, and thus they probably gained in exactitude and completeness.

At the same time, the effect of the non-use of the microscope for purposes of diagnosis may perhaps be recognised in the application of the terms “ichthyosis” and “wart growth” to what would seem to have been, the one a dry epithelioma of the lower lip, and the other a papillary epithelioma on the margin of the tongue, both of which, however, were successfully excised. Of six cases of “lupus,” so-called, it is almost certain that two at least were carcinomatous. Again, the obvious appearances of a growth named “medullary sarcoma” of the tibia, and those of a “fungus hæmatodes” springing from the skin of the forearm, are admirably depicted in words; but there is, of course, no recognition of the modern nice discriminations between the different forms of sarcoma, or between a soft sarcoma and an encephaloid cancer. Different morbid conditions of the kidney could not then be indicated by the various kinds of casts or other deposits in the urine; nor, lastly, were the numerous phenomena of disturbed sensation and motion, which enable us to determine the nature, and even the locality, of certain lesions of the brain and spinal cord, capable of being recorded. It would, indeed, become tedious, were I to recount all the references in the case-books of 1883, which serve to show the great advance which has been made in the methods and accuracy of surgical diagnosis since 1844-5. But it is in regard to the questions which next arise, namely, those which relate to the treatment followed in the two periods, that the contrast I have attempted to institute deepens in interest and in force.

3. *Comparison of the Treatment adopted in University College Hospital, at the two selected Periods, considered in reference to Hygienic Conditions, General Comforts, Diet, Medicines, and Surgical Appliances.*

—A.—*Hygienic Conditions.*—In 1844-5, the north wing of the Hospital not having been built, the surgical wards were much crowded; whilst, in 1883, in spite of an increase in the total aggregate of patients, the numbers in each ward were diminished, and the cubic space per bed proportionately increased. In the presanitary period of forty years since, sculleries and water-closets were but imperfectly separated from the wards; whereas now their separation is complete. Formerly, the doors, windows, and chimney-flues furnished but intermittent means of ventilation; now suitable channels, both inlets and outlets, are provided for the constant renewal of the air in the wards. Besides this, a laundry has been expelled from the basement, whilst that and all the corridors are now efficiently ventilated. The traps and drains are also maintained in good order. It is true that, in 1844-5, the Hospital had been built only ten years, and so far was relatively free from the dangers of so-called hospitalism than now; but, to meet such contingencies, the walls of the wards have been recently lined with an impervious cement, which is regularly painted; whilst the floors are kept carefully purified. Lastly, in the good old times, there were no special isolation-wards for infectious diseases; whilst now there are special erysipelas-wards for both sexes.

B. *Diet and General Comforts.*—It cannot be doubted that quite as great attention is now paid to the question of diet as was formerly the practice; and, as all hospital committees well know, the expenses of general maintenance have increased. In the year 1844, for example, with a total number of 1,410 medical and surgical in-patients, the total general expenditure of the University College Hospital was £4,976, whereas in 1883, with 2,849 in-patients, it was £19,822; so that, whilst the number of patients is doubled, the total expenses of the establishment are multiplied fourfold. We may safely conclude that a certain share of this large increased expenditure arises from a bountiful attention to the dietary of the in-patients. As regards the personal care and comforts bestowed upon these patients, it may be pointed out that the salaries of nurses and domestics, which were somewhat more than £1,600 in 1844, amounted to £2,300 in 1883. Finally, it is with sincere expressions of gratitude to the Sisterhood who now so solicitously watch over the nursing and general economy of the wards, that one alludes to those numerous manifest improvements, which conduce not only to the comfort and cheerfulness, but to the well-doing and speedier convalescence of the inmates.

C. *Medical and Surgical Appliances.*—In 1844, the total expenditure on medical and surgical stores for the hospital, was £643, but in 1883 it amounted to £3,239; that is, the number of in-patients having become double, the cost of their medical and surgical treatment has been multiplied five times. The payments for medical stores alone amounted, in the two periods, to £437 and £1,474, showing an increase in the ratio of 3.3 to 1; but those for the surgical stores (exclusive of instruments) were, at the two periods in question, £206 and £1,765, making an increase of eight to one. Of this last named large sum, no less than £1,273 is specially set down to payments for “antiseptic dressings,” of which, of necessity, by far the larger proportion was employed on in-patients. This great cost of the antiseptic system, as it is carried out at the University College Hospital, is a startling financial fact; but, if the results can be shown to justify the outlay, such expenditure is in itself highly to be approved. It is obvious, moreover, that the public, which provides the needful funds, ultimately reaps the benefit of that, as of all other improvements. Wealthy donors should remember this, in meting out their contributions to hospital resources.

The constant use of anæsthetics, not merely in operations, but for the purpose of more thorough, and otherwise painful, examinations in cases of injury and disease, greatly increases the relative cost of hospital maintenance at the present day. In 1844-5, these were yet unemployed; although it was Liston himself who, two years later, was the first in this country to perform an important operation under their influence. For it was in 1846 that he amputated through the thigh the lower limb of a man, put into a condition of insensibility by the vapour of ether, administered by the late Mr. Peter Squire, in the presence of the late Sir John Forbes, the then editor of the *British and Foreign Medical Review*. Liston's surprise and gratification when the patient, having recovered consciousness, refused to believe that his limb was off until it was shown to him, is still well-remembered by spectators of the scene.

Beside antiseptics and anæsthetics, to which we may add hypodermic injections, many new and expensive medicinal preparations are now ungrudgingly used in every hospital. Improved instruments for the performance of long-established operations, and innumerable and ingenious clamps, cauteries, and other novelties for the accomplishment of new and difficult surgical proceedings, are also freely pro-

vided for the modern hospital-surgeon and his patients. The cost of these at University College Hospital has doubled since 1844. In a word, it may be unhesitatingly affirmed, that the means, as well as the methods of surgical treatment, whether hygienic or curative, were, in 1883, far in advance of those which were available 40 years before that date.

Let us next proceed to consider the ultimate and practical question of results.

4. *Comparative View of the Results of Treatment at the Two Selected Periods.*—Under imperfect sanitary conditions, without special isolation wards, with less attention to the details of ward-management and to their cheering influences, with fewer scientific means of diagnosis, with no chloroform or ether, and no hypodermic method of quelling pain or other nervous disturbance, with simpler instruments, and less recondite dressings and appliances, were the results obtained in 1844-5 inferior to those realised in the same hospital, under other conditions, in 1883? The reply is "Yes," in certain particulars, but not in all.

In the first place, I find that, as well as I can determine, the average period of stay of a surgical patient in the hospital, in 1844-5, was 28½ days in the female wards, and 29½ days in the male wards; whereas, in 1883, it was about 26 days for the female, and 21 days for the male patients, this latter smaller ratio being due to the disproportionate number of simple fractures occurring in the male sex. Too much stress, however, must not be placed on these figures; since the total number of cases is not sufficient to neutralise the effect of important differences in the severity of the cases, whether in kind or in degree.

Partly on a similar ground, but also on account of the different value assigned by different persons to the terms "cured," "relieved," and "discharged as incurable," I refrain from recording, in regard to such general results, figures which would be unreliable, or actually misleading. For example, in the older records, cases of caries of bone, stricture of the urethra, epithelioma, and scirrhus, are entered as having been "cured," and in the later set, as being only "relieved," by operative treatment.

Even the ratio of mortality is deceptive, although it shows an advantage on the side of the later period, the percentage of deaths in 1844-5 being 7.5 for the males, and 6.5 for the females; whilst in 1883, these were about 5.75 and 5.74 respectively. But here, the fatal cases of the two periods are not commensurable, and a single accidental death, as for example from a burn, completely vitiates the result. Hence, a comparison of the treatment and its consequences, in distinct classes of injuries and diseases, will be found much more just and more instructive.

A. The chief point to be noted in regard to differences in the management of simple fractures consists in this: that those of the leg and thigh were formerly retained longer in McIntyre's or Desault's splints, before they were put up in starched apparatus, of which Liston was a great advocate. Patients so injured were, therefore, detained longer in the hospital than now. The introduction of the plaster-of-Paris treatment has still further abridged the time during which a bed is occupied with a case of broken leg, patella or femur. Wiring of bones was not attempted. Compound fractures proved to be very prolonged cases, and those of the lower limb were often fatal, with or without amputation.

B. Injuries not involving a breach of the surface of the body, simple inflammations consecutive to these, or so-called idiopathic local inflammations which come under the care of the surgeon, such as sprains, simple dislocations, synovitis, orchitis, and other cases, were not less satisfactorily treated in 1844-5 than in 1883, and the progress of the patients towards recovery was quite as rapid. Venesection is not mentioned in Liston's case-books, but cupping and leeching were in the ascendant. A patient with disease of the hip-joint, requiring at the time only the application of Liston's famous leather splint, informed her dresser that she had previously had 141 leeches applied around her hip in the course of three months. A man with acute synovitis of the knee was ordered 18 leeches over the joint on the day of his admission, 18 more on the day after, eight on the tenth day, and 10 on the twentieth day, and a week afterwards was discharged convalescent. For a condition described as "mania" from head-injury, a cupping to 12 ounces was ordered to the back of the neck, followed by two sets of leeches, 20 each time, to the temples, and, five days later, by a blister and the administration of calomel, and so was cured. Inflammatory urethral stricture, prostatitis, and supposed cystitis, were relieved by free cupping on the perineum, and so in many other diseases. Whether these sanguinary proceedings were necessary or even beneficial, and whether the local abstraction of blood is now too much neglected in cases of acute inflammation,

especially of important organs, are problems which I cannot here discuss.

C. In comparing the older and the more recent management of inflammations which end in suppuration, ulceration, or gangrene, or in combinations of these morbid processes, we meet with facts more or less unfavourable to the older practice, and unmistakably cumulative in support of the advantages of modern, that is, of aseptic, surgery.

Acute abscesses were treated by very free incisions with ultimate success; but suppuration continued for many days, and the abscess-cavity was filled up slowly. Nor were there wanting instances of more serious evils. Erysipelas and further abscesses often supervened. A patient with successive abscesses in the abdominal wall, due presumably to caries of the ilium, is recorded to have left the hospital on the thirty-second day, and to have died a few days afterwards. An acute abscess in the head of the tibia, opened with the trephine, was 77 days in healing. Abscesses and sinuses, around but not communicating with the hip-joint, are mentioned as not having become closed when the patient was discharged from the wards, after the expiration of 83 days. Lastly, an acute abscess in the knee-joint proved fatal, from distinct pyemia, on the forty-seventh day, purulent deposits being found in the lungs after death. It is hardly necessary to state that poultices, and not antiseptic dressings, were employed; and, although counter-openings were freely practised, no drainage-tube was in use; and the abortive treatment of moderate-sized abscesses by aspiration and the injection of morphia, carbolic acid, or iodoform, was unknown.

Ulcers of the integuments, which usually are not prone to allow infective absorption, were very well managed in Liston's wards. Rest, elevation, and his favourite water-dressing, consisting of wetted lint, covered with oiled silk, cured most cases. All greasy applications were rigorously forbidden, for against these Liston waged an angry war. A spreading ulcer with fetid discharge was treated with a lotion of chlorinated soda, or a weak solution of iodine, both strongly antiseptic and germicidal agents. Red wash, which probably acts in both these ways, as well as a local stimulant, nitrate of silver, and sulphate of copper, were used as now. Strong nitric acid was applied against phagedæna; solutions of chlorine and iodine against slight sloughings and actual gangrene.

Periostitis, osteitis, caries of bone, and necrosis, were treated successfully, as now, by subcutaneous incisions, trephining, the extraction of sequestra, and scooping. In one report, the dresser mentions an instrument under the name of a "proper scoop," which, I assume, was really a "sharp scoop."

Veneral diseases were treated without mercury, or with as small doses as possible, in accordance with views which had then been ably propounded; and, as no distinction was made between the hard and the soft sores, so far as their possible consequences were concerned, whilst the majority of those taken into hospital were evidently soft chancres, followed by bubo, the non-specific or almost non-specific treatment was very successful. Iodide of potassium was becoming a favourite remedy; but it is curious to observe that a common formula in the notes was one grain of blue pill and one grain of iodide of potassium, combined in one mass, to be taken three times a day. When the iodide was administered alone, the usual dose was three grains three times daily; very seldom did it reach to four or five grains. In deep syphilitic ulcerations, and in sloughing gummata, a weak solution of iodine was commonly ordered as a lotion. Numerous cases of syphilitic caries and necrosis were admitted, and one of extensive disease of the parietal bone ended fatally, with abscess in the corresponding part of the brain.

D. Wounds.—Passing by burns, scalds, and injuries from caustics, I proceed to consider the cases of wounds of greater or less severity, 28 in number, which were received into Liston's wards in the year in question. Four of these occurred in women. One only healed within a week, and, as stated in the notes, "by the first intention"—namely, a wound in the palm, involving the superficialis volæ and another small artery. A cut down to the patella granulated and suppurated, and healed favourably in 17 days; an abrasion over the shoulder led to erysipelas and axillary abscess, the patient leaving the hospital on the eighteenth day; the fourth and last case, a punctured wound in the foot, became complicated with abscesses in the foot, leg, and thigh, and was detained in the ward for 55 days. Of the 24 examples of wounds happening to men, four only healed, presumably by the first intention, in from five to nine days; these were wounds affecting, respectively, the ear, eyelid, scalp, and scrotum, the last one laying bare the tunica vaginalis. In six other cases, very free suppuration, with accompanying fever, ensued—namely, in a wound of the scalp, a contused wound of the eyelid and eyebrow, a glass-cut of the forearm, a long split in the perineum, a lacerated wound of the foot, and, lastly, a

scalp-injury requiring counter-openings; in these cases, healing was deferred to from 15 to 28 days. In the remaining patients, still graver complications ensued; they may be briefly summarised thus: Punctured wound of thumb, erysipelas; the patient, being in good circumstances, was sent out on the eighth day; bite on the back of knuckle, erysipelas, abscess; discharged on the twenty-first day; slight wound in butcher's hand, cellulitis, abscess in forearm and arm, 22 days; contused wound of leg, rigors, abscesses, 30 days; wound of ulnar and another smaller artery above the wrist, ligature, cellulitis, sloughing of fascia up the forearm, 32 days; lacerated wound of palm, abscess, sloughing, much fever, 34 days; punctured wound of thumb, suppuration, frequent rigors, 39 days; contused wound of ear and side of head, so-called "erythema," suppuration, 42 days; punctured wound of thigh, profuse hemorrhage, suppuration, burrowing of pus, 45 days; attempted suicide, division of brachial artery at bend of elbow, cellulitis, sloughing of areolar tissue, severe fever, 54 days; gunshot wound of forearm without fracture, deep seated suppuration, with great fever, 53 days; lastly, two fatal cases, namely, a lacerated wound of the leg, erysipelas, rapidly spreading diffuse cellulitis, gangrene, and death on the eighth day; and a cut-throat dividing the trachea, followed by bronchitis and infective pulmonary abscesses, proving fatal after 61 days.

These cases, the only ones admitted, are surely sufficiently striking; but we have yet to consider the results of the cutting operations performed in the hospital during the twelve months of 1844-5.

Operations.—Five cases of lithotomy, four of herniotomy, and one of perineal section for laceration of the urethra, all successful, afford evidence of Liston's skill as an operator; and, as regards the duration of the cases, three of the lithotomy patients were cured within the month, one on the thirty-third day, and the remaining one, an elderly patient with large stone, on the fifty-third day. One of the hernia cases was discharged cured on the ninth day, one on the twentieth, another on the thirtieth, but one not until the forty-seventh day. The perineal section was closed on the twenty-eighth day. A case of phimosis, in a boy, required twenty-four days to cicatrise; and another, in an adult, was followed by erysipelas, severe rigors, and bubo, and was healed only after thirty-nine days. Fistulae in ano behaved as now; but one patient thus afflicted died, on the seventh day after the operation, from pulmonary complications. Two cases of extravasation of urine were unavoidably fatal, in one of which minute abscesses were found in both kidneys.

For the removal or excision of new growths, twenty-one operations were performed in the year 1844-5—eight upon men, and thirteen on women. They may be thus briefly detailed. Male cases: three, namely, a small epithelioma of the tongue, epithelioma of the lower lip, and a small thyroid cyst, healed by first intention by the end of a week; two, namely, a larger thyroid cyst, and a broad epithelioma of the lower lip, healed with granulation in nineteen and twenty days; three, namely, a parotid tumour, a cystic sarcoma of the testis, and a large lymph-cyst of the neck, all attacked by erysipelas, were cured in twenty-nine, thirty-four, and fifty days respectively. Female cases: five fatty tumours, the wound, in four, healed by granulation on the twelfth, eighteenth, twenty-fourth, and fifty-seventh days; in the fifth, a succession of rigors on the first, third, and fourth days, with offensive discharges, the patient was sent out of the hospital on the fourteenth day, and died seven weeks later; five mammary tumours, one a cystic tumour, patient left the hospital doing well on the fourth day; the others, namely, a painful tubercle, two sarcomas with cysts, and a scirrhus, healed by granulation and suppuration, twenty-one to twenty-eight days; a large parotid tumour, followed by abscess and profuse suppuration, thirty-two days; and, lastly, a large thyroid body, seton inserted for ten days, suppuration, intense fever, fifty-seven days. To these cases, I must now add that of a very large ovarian tumour, of three years' duration, tapped once before; operated on, in the words of the case-book, "in Mr. Liston's usual manner," that is, "by a previous incision into the abdomen," and "the introduction into the cyst of a large trocar and cannula;" "thirteen quarts of a dark-coloured glutinous fluid" evacuated, some more afterwards squeezed out, wound closed; next day pain and vomiting; on third day, decided peritonitis; treatment by leeches, fomentations, opium, calomel, hydrocyanic acid, and creasote; incessant vomiting, exhaustion, and death on the twenty-second day. *Post mortem*: peritoneal cavity contained "three pints of purulent-looking fluid, with portions of badly formed lymph floating in it;" "cyst partly refilled," with "soft adhesions to adjacent parts." In Liston's *Elements of Surgery* (edition 1840) ovariectomy is denounced, from contemporary experience, as an "unjustifiable piece of butchery!" How all this has been changed by cautious, and, if I may coin a word, *precautious* surgery!

In two subcutaneous operations, one for a contracted toe, and

another for displacing a loose body from the knee-joint into the areolar tissue outside the capsule, the wounds healed at once, without suppuration; but in a third, for dividing the proper extensor tendon and the long flexor tendon of the great toe, abscesses and ulceration detained the patient 69 days in the ward. Of plastic operations, Liston's great pride, a divided hare-lip united before the end of a week, and an example of webbed fingers healed in a fortnight; but four operations on the face, for pendulous upper eyelids, cancrum oris, partial destruction of nose and upper lip, and complete loss of the nose, required for the healing process 33, 42, 43, and 66 days respectively; an operation on the scrotum, designed to cover a testis exposed by gangrene, necessitated treatment for 50 days. Ten amputations and one excision of a joint are also recorded. First, four female cases: compound dislocation of finger, erysipelas, abscess in axilla, 30 days; diseased phalanges and joint of finger, erysipelas, abscess in axilla, 29 days; excision of elbow-joint, ends of humerus and ulna removed, abscesses and extensive burrowing of pus up and down the limb, eventual good result, 93 days; large "bleeding sarcoma," or "fungus hæmatodes," springing from the integuments of the forearm, three or four years' duration, lately rapid growth, axillary glands sound; amputation above elbow, rigors on seventh day, discharges very offensive, abscess in axilla with similarly offensive pus, sweatings, emaciation, jaundice, made out-patient on sixty-ninth day, death one month afterwards. Male cases, seven: compound fracture of finger, healed in 16 days; badly crushed finger, so-called "erythema," abscess in palm, patient discharged on twentieth day; comminuted fracture of finger, cellulitis, deep subfascial suppuration along the forearm, 59 days; disease of tarso-metatarsal joint of great toe, removal with internal cuneiform bone, prolonged suppuration in foot, 92 days. There remain three major amputations, all fatal. First case: a buffer-accident, compound fracture of tibia and fracture of head of fibula into the knee-joint on the right limb, fracture of tibia with lacerated wound of leg on left limb, primary amputation of right limb above the knee; progress favourable for three days, then some hemorrhage, left limb became much discoloured, rapid exhaustion, followed by death on the eighth day; *post mortem*, collection of pus around cut end of femur, and extending as high as great trochanter, yellow line of demarcation on left leg, internal organs not examined. Second case: sarcoma of leg, originating in a cicatrix from burn; amputation above knee, profuse suppuration, separation of flaps, hemorrhage, general oozing, death on the sixth day. Third case: sarcoma springing from front of tibia, amputation through knee-joint, patella and cartilage on end of femur left, good progress for five days; on the sixth day, a slight rigor; on the ninth day, severe rigors; on the eleventh day, another rigor, with profuse sweatings, discharges fetid, flakes of loose cartilage coming away; on the thirteenth day, another rigor, subsequently several others less severe, and death on the seventeenth day. No *post mortem* examination, but the case was typically pyæmic.

In the case of aneurysm of the femoral artery, in which the temperature of the two limbs was registered by aid of a thermometer, as the tumour reached above Poupart's ligament, it was necessary to tie the external iliac artery. On the third day, swelling, and so-called "erythema," appeared around the wound; the patient had dyspnoea, a rapid pulse, restlessness, and delirium; the temperature between the toes of the affected limb, which became dusky, was gradually lowered to 84°, 72°, and 64°, as compared with 93°, 97°, and 78°, between the toes of the sound limb, the temperature of which, be it observed, also fell; the patient succumbed, apparently septiciæd, on the sixth day.

The picture thus finished of the results of the surgical practice in Liston's wards in 1844-5 is full of instruction. The separate mention of individual cases, all of which I have read from beginning to end, serves to enforce the lessons they teach. In the light of present knowledge, the frequency of grave inflammatory complications must be attributed to removable or corrigible conditions, which would have been prevented or minimised by more effectual sanitary arrangements, by the isolation of infective disease, and by strict antiseptic precautions.

It will have been noticed that, in cases of injury or disease not involving a breach of surface of the body, and in cases in which, as in operations in the perineum or ischio-rectal fossa, antiseptic dressings are not available, the results of the older practice are quite upon a level with those of modern experience. Even in regard to ulcers, this is also true. But in the case of recently opened abscesses, and especially of open wounds, whether accidental or intentionally inflicted by the surgeon in his operations, in which the delicate walls of divided and exposed blood-vessels, the cut or lacerated surfaces of the interstices of the areolar tissue, and the torn and irritated lymphatics are helpless to resist the entrance into the system of infective fluids and

organisms, we find that prolonged suppuration, erysipelas, cellulitis, lymphangitis, pyæmia, and septicæmia, as it were, dogged the footsteps and foiled the handiwork of the most skilful operator of his time.

We have seen that all abscesses, when opened, continued to suppurate freely; whilst in four cases (14 being the total number), grave complications arose, one case ending fatally. Again, of 27 cases of accidental wounds, only five healed quickly by the first intention, seven by granulation and free suppuration, and 13 after prolonged suppuration, cellulitis, or erysipelas, whilst two terminated in death. Lastly, of 54 wounds made in operations, eight only united by the first intention, 26 closed after granulation and free suppuration, 14 were cured after the occurrence of erysipelas or cellulitis, with secondary abscesses, and, excluding two unavoidable deaths from extravasation of urine, four died of septicæmia or pyæmia.

It must be within the experience of every living hospital surgeon that no such calamitous results follow his own practice. It is unnecessary to quote here, in detail, the number and kinds of cases treated in my wards in the year 1883; but the facts relating to those cases, which are carefully reported, show indisputably that abscesses, both acute and chronic, treated by incision, proper drainage, and antiseptic dressings, heal with much less subsequent suppuration, and in less time, than under the open treatment; that accidental wounds, maintained in an aseptic condition, are far less frequently followed by serious complications than formerly happened; and that operation-wounds of all kinds heal more quickly and kindly under antiseptic treatment. By way of illustration, it may be mentioned that in 24 operations for the removal of new growths, the wounds were healed in five cases between the seventh and tenth days, in seven cases between the fourteenth and twentieth days, in seven cases between the twenty-fifth and thirty-fifth days, in four cases between the forty-third and fifty-fifth days, and in one only, a case of a double operation on a cancerous penis, on the seventy-third day. It must also be added that in no instance was there more than a show of suppuration, and that no case of erysipelas, infective cellulitis, septicæmia, or pyæmia, arose, during the year in question, in the wards under my care; yet these diseases were not absent from the hospital, for three cases of pyæmia, one of septicæmia, and five of erysipelas, originated in other wards, whilst 16 cases of erysipelas or infective cellulitis were admitted from outside into the proper erysipelas-wards. It must, in justice to the older period when there were no isolation-wards, be granted that some of the cases of injury complicated with erysipelas were already so affected on their admission; but most of those cases, and all those following on operation-wounds, arose within the hospital, making a total of at least 24 cases of septic disease arising in Liston's wards in 1844-5, against none arising in my wards in 1883. It is easy to trace, in these older records, the frequent and long acknowledged co-existence of endemic and hospital erysipelas, to which a recent joint Franco-German Surgical Commission of inquiry into the effects of that disease has directed attention; and, indeed, it may be said that in Liston's time not only did inflammation, which, like fire, may be a good servant but is a bad master, attack every wound, and free suppuration, which if not a combusive is an exhaustive process, run riot in the wards, but the demon of erysipelas, which so often gained an entrance therein, if it did not strike all the patients, at least hovered over every bed. Its vagaries were well understood; its advent took no one by surprise, and there was one bed in particular, in the corner of a certain ward, in which it most frequently seized its victim. This familiarity with it in hospital practice explains perhaps the promptitude, and, judging from the number of recoveries which followed it, the success with which it was treated when it occurred. But neither it, nor other septic diseases, nor the tendency to undue suppurative inflammation, were efficiently prevented.

On general principles, every surgical patient was subjected, almost *in limine*, to the inevitable purge, consisting of the nocturnal dose of blue pill or calomel, with colocynth and henbane, and a choice between castor-oil or a black or white saline draught in the morning. Subsequent fever was treated by salines, often combined, even in the later stages, with antimonial wine, and sometimes with the then new remedy, aconite; rigors, sweatings, and exhaustion, were combated with ammonia or mineral acids, administered in decoction of bark.¹ In the meantime, complete purity of the surroundings, absolute surgical cleanliness, and the exclusion of all noxious substances, were not strictly aimed at, or attained. Sanitary vigilance was dormant, or did not exist; and the dressings, the famous water-dressings, alas! were doubtless the constant innocent, but not innocuous vehicles of infection. Liston, as is well known, denounced all resinous plasters as

"dirty applications," and all ointments as "greasy abominations," prefixing to these appellations a strong characteristic epithet. Iodine-glass plaster, wet lint, and oiled silk, were no doubt neat, and to the eye cleanly; but they were not aseptic. Lint "stuffed" into open abscesses, sinuses, and carious cavities, or placed for a few hours between the flaps, after an amputation, would certainly contain organised or unorganised extraneous particles; and the water with which it was moistened, surely not previously boiled, would also contain them. The heat and moisture of the living tissues, and of the exudations from them, would favour the development of germs, as well as the formation of lifeless but injurious chemical compounds. To us, the consequences are clear. No change of dressings, however frequent, and no occasional use of chlorinated lotions, or iodised solutions, would cleanse effectually the microscopic but Augean recesses of the injured tissues. Hence, more or less suppurative action ensued, and more or less deadly agents of infection gained an entrance into the vessels and the blood. Asepticised air, hands, instruments and sponges, together with all the varieties of shortened aseptic ligatures, sutures, and dressings, form a remarkable contrast to Liston's means of treatment. It would be untrue, and a calumny against Nature, to say that these and other antiseptic agencies are everywhere and at all times indispensable necessities of surgical practice. As narrated to me by a French surgeon, an eye-witness to the facts of an African campaign, an Arab's cheek, all but sliced off and hanging down, may be stitched up in its place, and bandaged with strips from his own cotton garments, and may then unite by the first intention, with scarcely a drop of exudation; and I have been told of a gentleman colonist in Australia, who cut open his knee-joint in felling a gum-tree, sewed up the wound himself, and put on a back-splint, before the nearest surgeon, who resided 60 miles off, could arrive, and was cured "straight off" without suppuration, preserving a perfectly useful limb. Union by the first intention occurred long before the antiseptic system was devised; and it occurs still without its aid. But in formidable operations anywhere, and in most operations in hospitals, it is a safe and therefore an incumbent provision to adopt. In this view, adequate hospital expenditure for antiseptics is fully justified. Their employment shortens the period of cure, limits the amount of suppuration, diminishes the drain upon the bodily strength, lessens the risk of complications and the chances of future organic degenerations, and thus contributes to a more rapid, complete, and lasting recovery.

As anesthetics encourage patients to submit more frequently, willingly, and hopefully to more and more serious operations, so antiseptics have emboldened surgeons to attempt, and enabled them to succeed in, enterprises formerly considered hopeless and unjustifiable. When I remember Liston's grave and despairing looks at his too frequent disappointments, I welcome the confident glance and triumphant expression of many a modern operator.

Of the three great cavities of the body, the abdomen, with its membranous coverings, the least protected and most accessible, containing parts less immediately essential to life, and yet, from its size and the extent of its lining membrane, the most prone to a rapidly fatal inflammation, was the first to be entered by the properly aseptic hand; next, the thorax, with its half-bony and half-membranous walls, and therefore better protected and less open to access, containing parts more immediately necessary to life, but lined by a membrane of smaller extent, and subject to less formidable inflammatory reaction, yielded entrance to the antiseptic method of procedure; and, lastly, the cranio-spinal cavity, the smallest, the most perfectly enclosed, the best protected, enclosing organs of vital essentiality, and yet lined by a membrane of moderate proclivity to disease, has also been invaded by the new surgery, under the guidance of an improved diagnosis, in search of removable disease. Even the cervical region, which, though not designated a cavity, is, nevertheless, a specially adapted part of the general cavity of the body, has become the field for the employment of aseptic methods.

It might thus seem as if surgical enterprise and skill, applied to the trunk as contrasted with the limbs, had reached their utmost attainable limits; but it cannot be so. Already, the cry is for still greater security against the consequences of irritant and septic influences. That this will be obtained in the future, no one can doubt.

Forty years since, Liston was deprived of our present advantages, and suffered accordingly; but his dexterity and fame as an operator, his desire to simplify all surgical proceedings, his lasting devotion to and sound knowledge of anatomy, and his interest in the then opening science of pathology, require no vindication at my hands. May the next forty years witness as great advances as have been realised since his death; and, in that future, may the reputation of British surgeons, and the triumphs of British surgery, be maintained by our successors with undiminished lustre!

¹ "If morbid spot of septic sore invade,
By heaven-sent bark the morbid spot is stayed."

— From Rev. Sydney Smith's *Poetical Medicine Chest*.