This theory involves two assumptions: (1) that destruction of a bilaterally associated nerve-centre gives rise to constitutional disturbance, and (2) that the lesion causing herpes zoster causes destruction of the nerve-centre. No proof is afforded of either; and, so far as I know, there are no facts in pathology to warrant them. On the contrary, apart from the pathological process involved in its general destruction, general constitutional symptoms do not follow from the suppression of the function of even large parts of the nervous system. It is, therefore, unreasonable to attribute so much to the "loss of a definite entity" of microscopic dimensions only. If there were actual destruction of the nerve-centre in zoster, we should expect some paralysis of motion or sensation; but the commonest permanent result is chronic painfulness, generally attributed to a lesion of the nerve-trunk, but for the recognition of which there must be a percipient centre.

3. Having assumed that he has met with a case of bilateral zoster giving rise to constitutional symptoms, having further assumed that such an association is usual, and stands in need of explanation, and having formulated the explanation already quoted, he announces the discovery of a "natural law governing the action of the nervous system in disease." This "law" is not formulated; but it appears to be, that neuroses do not attain their acme until both sides of the body are affected. Hemicrania is said to afford illustrations of it; but the "law" needs to be formulated a little more precisely before any opinion of its value can be formed.

4. Finally, he suggests that this case affords an example of the similarity of plant and animal pathology.

He quotes a passage from Sir James Paget's address at Cambridge, which appears to be a sufficient answer to such a suggestion, as in it Sir James points out the fundamental distinction between the elementary pathological processes of animals and plants, caused by the influences of the nervous system. Plants have no differentiated nervous system; and, therefore, they are the last place in which to seek analogies with neuro-pathology.

He thinks that the death of a branch from frost is analogous to this attack of herpes affecting a definite nerve-area after a chill; but the comparison is strained. In the first case, we have to do with the direct effect of cold, as in gangrene after frost-bite; in the second, with the indirect and obscure action of a chill of the surface, determining changes in subjacent structures.

We want thoughtful men to help to organise facts; and I am, I trust, always grateful for well considered speculations, adapted to the outline of our present knowledge. It is from no disapproval of speculation generally, but from doubt of the method adopted in the present instance, that I have ventured to point out what appear to me to be objections to it.

CIRRHOTIC ENLARGEMENT OF THE LIVER IN A FEMALE.

By JAMES OLIVER, M.B. Edin., M.R.C.P. Lond.,
Assistant-Physician to the Hospital for Women, Soho Square.

AN interstitial hepatitis resulting in cirrhotic enlargement of the liver is, by some authorities, denied; whilst others, admitting its existence, express the opinion that the affection is one totally distinct from that of cirrhotic contraction. The active congestion, going on to actual inflammation, and produced by the more or less constant irritation of the organ by the imbibition of alcohol, results in an exudation and proliferation of cells which, in the early stage, are seen to be oval, fusiform, and round in shape. Later on, these so called embryonic cells undergo gradual change, and develop fibrous tissue, which will be found along the course of the blood-vessels, and penetrating between the lobules.

In this stage, the liver is usually enlarged. The newly formed fibrous tissue, however, tends to retract, and cause wasting of the lobules, thereby developing a cirrhotic contraction, and producing the well known hob-nail liver.

The symptoms associated with cirrhotic enlargement, it is to be remembered, are not solely dependent upon the structural changes occurring in the liver, but are rather the result of the cause of the irritation in this all important organ. The alcohol, prior to its passing to the liver through the medium of the portal circulation, exerts its deleterious influence on the mucous membrane and other structures of the gastro-intestinal tract, creating a chronic hyperæmia, which

eventually results in impairment of function, not only of the secreting, but of the absorbing, tissues of the canal. Hence the reason that, even in the early stage, we may have loss of flesh, a marked symptom of the disease.

It is possible—nay, likely—that interstitial hepatitis may result from causes other than alcohol; regarding these, however, little or nothing is known.

CASE.—E. H., aged 28, married ten years, no children, and no miscarriages, four months ago began to notice a slight increase in the size of the stomach, and ever since then there had been complete amenorrhœa. The catamenia appeared at the age of 11, and, hitherto, she had been always perfectly regular, the hæmorrhagic discharge recurring every twenty-eight days, and continuing three or four. For the last three months, she had suffered from sickness, vomiting food at all times soon after its ingestion. The patient had a feeling of intense lassitude, and was constantly more or less drowsy. Two years ago, the appetite began to fail; and, during the last twelve months, she had lost flesh. At times she had suffered pain in the region of the liver, and under both shoulder-blades. During the last two months, headache had been a troublesome symptom, and there had been occasional attacks of profuse epistaxis. Lately there had been diarrhœa, not specially in the morning. Her habits had been intemperate; she had taken two pints of beer daily, besides some spirits. The liver was uniformly enlarged, measuring nine inches and a half in the nipple-line, and five below the ribs; the lower margin, hard and regular, was felt below the umbilicus. The slightest pressure anywhere in the region of the epigastrium caused a feeling of sickness. The urine was thick, and of a deep colour; it contained lithates, bile-pigment, and a small quantity of albumen. For two months, the patient had to get up at least once every night to pass urine. There was slight œdema of the ankles; no ascites.

BRITISH MEDICAL ASSOCIATION.

FIFTY-THIRD ANNUAL MEETING.

PROCEEDINGS OF SECTIONS.

THE MECHANISM AND MANAGEMENT OF THE THIRD STAGE OF LABOUR.

Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association in Cardiff.

By D. BERRY HART, M.D., F.R.C.P.E.,

Lecturer on Midwifery and Diseases of Women in the School of Medicine, Edinburgh.

It may seem, on first sight, that the subject chosen for discussion is too elementary and well threshed out to claim much of the attention of our present meeting; but, although no process has been so much and so often studied practically by the vast majority of our profession, it is yet true that we know only too little of its real mechanism, and are in great variance in regard to its treatment; and this in spite of the fact that the expulsion of the placental membranes entire and without any internal manipulation, in a natural labour, is of the highest importance in relation to after-septic processes, whether the graver septicæmia of puerperal fever, or the less severe and yet troublesome attacks which often leave chronic endometritis, cellulitis, and peritonitis as their sequelæ.

Apart from all this, however, the present time seems to me opportune for considering the subject in our country. The attention now paid to abdominal palpation before and during labour, the increase in our knowledge of the exact relations of the membranes in the third stage—due mainly to the splendid work of my friend Dr. Freeland Barbour, and the application to obstetrics of the principles of antiseptics imperishably connected with the name of Lister—enable us to consider the whole matter from new and commanding stand-points.

In the following remarks, I purpose briefly to relate the facts known to us in relation to the natural and morbid mechanism of the third stage, pointing out what has yet to be discovered in order to make our knowledge complete. The management which seems to me to favour the natural process, or to be necessary when the natural becomes morbid, will also be briefly considered.