even to delay it, and thus save the perinæum. This is a use of the forceps which I think has not been sufficiently dwelt upon, and which can only be satisfactorily accomplished by the long curved forceps.

I repudiate the dogma, that force is never to be used in artificial delivery; in employing it judiciously we imitate nature herself, and it is only when abused that it becomes dangerous. I have over and over again verified in practice that, with proper precaution, and with appropriate manipulation, delivery may be accomplished with safety to both mother and child, even when considerable force has been necessary to effect the extraction of the head.

I regard the use of the forceps as more appropriate in those cases in which many obstetricians recommend the ergot of rye, or other oxytoxics, for the following reasons: it gives the accoucheur a more certain, regular, and complete power over the course of labour, especially during the exit of the head, enabling him either to hasten or retard its progress as circumstances may require.

Their use does not involve certain risks which are admitted to attach to the action of ergot, even by those who advocate its employment in tedious labour. Dr. Johnson, for instance, says, "That when ergot is given it brings matters to a crisis, for if the child be not delivered within a certain time it will in all probability be destroyed." Drs. Hardy and McClintock state they "have on many occasions observed the ergot of rye to exercise a very depressing influence upon the maternal circulation." More recently Dr. McClintock has advocated the view, taught by Dr. Robert Lee when I was a student, that danger to the child arises from the incessant uterine action, and the consequent pressure, which the ergot is so liable to occasion. Dr. Churchill attributes to its use "retention of the placenta, from irregular uterine action, after the birth of the child." (*Theory and Practice of Midwifery*, 1866, p. 264.)

In my own practice I rarely use it in actual labour, preferring it as a means of completing abortion, when all hope of saving the ovum is abandoned, or of producing contraction in *post partum* hæmorrhage. It is my firm conviction that the more the forceps come into favour, the more rarely will ergot of rye be used, and that ultimately it will be altogether set aside as a substitute for what I cannot but regard as a safer and more efficient means of affording assistance in tedious labour.

In conclusion, I must beg your indulgence for many imperfections in this mere sketch of a very large and comprehensive subject. Those who desire to follow it out more completely than my limits on this occasion will permit, I would refer to the detailed observations of the authors I have quoted, and especially to two admirable papers in the first volume of the *Transactions of the Obstetrical Society of London*, by Dr. Tyler Smith and Mr. Harper.

I have purposely abstained from wearying you by a detail of cases; I may, however, observe that the views which I have brought forward are founded not upon theory, or the teachings of others alone, but are an exposition of the opinions I have formed during a somewhat lengthened and extensive course of practical experience as one of the accoucheurs to the Ladies' Charity of this town; an institution which comprises the results of about two thousand cases of labour annually, and those chiefly in a class of women especially liable to various forms of dystocia and other complications of parturition.

[ERRATA. At page 480, line 9 of second column, for "posterior", read "anterior"; and in the succeeding line, after the word "temple", insert "the other the posterior over the side of the occiput."]

REPORT OF

TWENTY-FOUR YEARS' VACCINATION IN THE MOLD DISTRICT OF THE HOLYWELL UNION.

By EDWARD T. HUGHES, M.D., etc.,
Public Vaccinator.

DURING the last twenty-four years I have reported to the Holywell Board of Guardians 10,344 successful vaccinations, and noted their influence in checking the spread of small-pox.

For the first five years, the Mold district consisted of three parishes; population 10,851; acreage 21,505. It was also the Mold Registration District. The registered births were 1535, and vaccinations 1340. Allowing for deaths previous to vaccination, this would have been a satisfactory number, had not a great majority of the children been above a year old, when vaccinated. Small-pox prevailed in some part of the district every year, and eighteen children born therein died of it, nine of whom were under one year, and only one above six years of age, all unvaccinated.

At the end of 1847, the parish of Northop was added to the Mold district, its population according to the census of 1851 being 14,577, and acreage 29,505. From January 1st, 1848, to December 31st, 1853, (when the Compulsory Vaccination Act had just come into force,) the number of registered births was about 2530, and my vaccinations 2609, of these 1303 were in children above a year old; so that, although the district had been on three occasions during the six years vaccinated almost from house to house, the arrears were considerable, as six hundred and forty-six of the children vaccinated were born before January 1st, 1848. These were well accounted for in subsequent vaccinations.

Twenty deaths from small-pox were registered during the same period; twelve of these occurred in the Mold Registration District, in which there had been but one fatal case, and that an imported one, from August 24th, 1847, to September 16th, 1852, when it was brought into Mold by the family of a knife-grinder. Five children died of it, four the children of tramps lately come into the town, and the fifth a sickly child sixteen months old, who had been unfit for vaccination from birth, and whose parents lived next to the house in which it first broke out. The other seven cases were in persons from seventeen to thirty years of age, all unvaccinated but one, who was said to have been vaccinated by her father, a village shopkeeper in Carnarvonshire, of which there was but faint proof. The other eight deaths in the parish of Northop, which forms part of the Flint Registration District, were all in young unvaccinated children.

My vaccinations during the remaining thirteen years have been 6395, and of these 403 were in children above a year old. The registered births must have been 7000 or thereabouts, but of these I have no trustworthy account. The deaths from small-pox were six, three in the Mold Registration District, and three in the parish of Mold, all in persons not vaccinated. To these I will again refer, as I propose now to consider the vaccinations of the last thirteen years, as illustrative of the working of the Compulsory Vaccination Act under very favourable circumstances.

The act to extend and make compulsory the practice of vaccination came into operation on October

1st, 1853, and, although it was far in advance of any previous legislation on the subject, the reception it met with was not favourable. Had early and well-advised attempts been made to amend, instead of strenuous exertions to repeal it, I am satisfied that it would have been successful, and that deaths from small-pox would ere this have ceased to hold the prominent position they still do in the returns of the Registrar-General.

I had fortunately no difficulties to encounter in my attempts to carry out the intentions of the act, as for above twenty years I had performed most of the vaccinations in the neighbourhood. I was also then registrar of births and deaths for the Mold District, and at once acknowledged the importance of the registration scheme, which, although not perfect, was workable, and tolerably efficient. The registers of successful vaccinations have been therefore carefully kept, and are up to the present time accurate records of the working of the act, as they contain not only the vaccinations of the public vaccinator, but also those of other medical men residing in the district, who have with great regularity forwarded their certificates of successful vaccination to the registrars. I have taken care that all deaths previous to vaccination, and changes of residence, should be noted, so that every entry is accounted for.

The registers of successful vaccination from which the following table was compiled on the first of last January, contain entries of the births of all children born on and after the 1st day of August, 1853, who were alive at the date of registration, also the vaccinations of children born in the district, but not registered, and of others, who, not born in the district, were resident therein when vaccinated.

Return of Successful Vaccinations in the Mold District from October 1st, 1853, to September 30th, 1866, under the Compulsory Vaccination Act.

Number of entries in Register-Books of successful vaccination.	Number in the district.	Number not born in the district.	Left previous to vaccination.	Died previous to vaccination.	Had small-pox previous to vaccination.	Have not been vaccinated.
For eleven years ending Sept. 30th, 1864	5517	1665	240	119	442	1
For the year ending Sept. 30th, 1865 ..	677	520	49	19	75	3
For the year ending Sept. 30th, 1866 ..	731	505	35	24	67	100
Totals	6925	6034	192	584	4	111

Thirty-two children registered during the last quarter were above three months old on the 1st of January; so that the total arrears are 143—the greatest number there have been since the Act came into force. It will be seen that no death from small-pox has occurred among children born in the district during the last thirteen years, who were alive when their births were registered. One infant died of it when only a few days old, whose mother was ill of varioloid disease at its birth.

We have, however, had numerous importations of small-pox during this period; and five other persons have fallen victims to it, all of whom were unvaccinated. In the year 1857, it was very prevalent and fatal in several parishes of the Ruthin Union, immediately adjoining the Mold District; and I had several cases under treatment, all of which recovered. One infant, a few weeks old, had it previous to vaccination; but it did not extend much beyond the neighbourhood in which it first broke out. In 1858

or 1859, small-pox was so severe in Chester (which is only twelve miles distant from Mold, with frequent daily communications by road and rail), that the question was seriously entertained whether a special ward should be set apart for cases of it in the General Infirmary, or a temporary hospital erected. We had several importations (servants sent home sickening); but in no instance did it spread.

In 1860, the child of a tramp, with the eruption well out, was brought into my surgery, and sent into lodgings, where it died in a few days; but every necessary precaution was taken, and no second case occurred. In 1862, small-pox appeared in the neighbouring town of Denbigh, and surrounding country; but I am not aware that we had a single case of it. In 1863, a young man came into Mold from Liverpool, and sickened next day of a mild attack of varioloid disease. His sister, who lived in the same house, took it, and was confined when ill. Her infant died of small-pox when only a few days old; but again the disease did not spread.

In July 1864, the next outbreak occurred, and was brought into the most distant part of my district from Liverpool. Three cases proved fatal before I heard of it; but, fortunately, the neighbourhood was well vaccinated, and it did not further spread. A man aged 24, a girl aged 16, and an infant three months old, whose parents had lately come to reside in the district, were the victims—all unvaccinated. About the same time, small-pox was brought into Mold; and during the next twelve months I could reckon at least thirty distinct importations of it into different parts of the district. The first cases all came from large towns (Liverpool, Birkenhead, Chester, Wolverhampton, and Birmingham); the later ones, chiefly from the neighbouring union of Wrexham, an infected part of which adjoined one of the most populous portions of my district. In no single instance did it spread beyond the locality where it broke out. One man, thirty-six years of age, died of it. He had not long lived in the neighbourhood, and had never been vaccinated. Last year, a convalescent again brought it into the district; but only two persons took it in a mild form, and soon recovered.

Such, then, are the results of vaccination steadily performed for many years in a district beyond the average size, both as regards acreage and population, which latter has greatly increased during the last five years, and must now exceed 16,000. Although I have had no difficulties to overcome, and have only to acknowledge kind and cordial co-operation, I am not the less aware of the nature of those which beset the path of the public vaccinator, and which require greater attention than they have yet received. It is in the large and densely populated towns that these difficulties appear to be the greatest; and it is in them that they must be overcome, before the success of any Vaccination Act can be perfect. Small-pox appears to be ever present in some of them, and it is generally from them that it is communicated to every accessible part of the country. In my own district, every solitary importation that I can remember came from a large town; and during the epidemic of 1864-65, as I have before mentioned, all the earlier cases were from large towns, the later ones being from neighbouring districts, which probably had in like manner received their first contagion. Railways have made it much easier to send sick servants home to the country, to our great disadvantage, as I have seen several reach their homes with small-pox eruption well developed.

As the present system has proved ineffectual, why not, for a time at least, appoint subinspectors to confer with and assist the public vaccinators in large

towns in overcoming resistance to their duty, as also to divide with them the responsibility incurred by its non-performance? What an opportunity London affords at the present time of testing the value of such a proceeding! I have long felt that the present number of inspectors is too few, and their districts too large, to allow them, able and zealous as they are, to do much practical good to the cause of public vaccination; but if to each were assigned a certain number of subinspectors, on whom should devolve the responsibility of seeing the duties of the public vaccinators in their respective districts well and ably performed, and at stated periods of reporting thereon, I will venture to say that a great step towards providing for the vaccination of the people would be achieved, and that we should soon cease to hear reports of badly performed vaccination and impure lymph, the evils of which have been somewhat magnified.

Many other matters of importance relating to public vaccination have been repeatedly urged upon the Government by the President and members of the Epidemiological Society, Mr. Griffin, and others; and it is to be hoped that the amended Vaccination Act soon to be introduced into the House of Commons will show that they have not been passed by unheeded.

I cannot agree with those who think the registration of successful vaccinations unnecessary. The present system requires some amendment, as, to make it at all perfect, it is requisite that every entry in the register-books should in some manner be accounted for; and then they will form valuable records.

To carry out, however, any improved scheme of public vaccination, a larger grant of money will be required: and can it with any good reason, be withheld? Our country spends annually vast sums of money in devising and perfecting the most deadly weapons of destruction to protect her shores from foreign foes: why, then, should she give grudgingly for the more laudable purpose of saving her children from a loathsome enemy, whose victims we count annually by thousands, and against whom we have a certain means of defence, had we the wisdom and courage properly to use it?

CASE OF

ACUTE PLEURISY, TERMINATING IN EMPHYEMA: OPERATION: RECOVERY.

By EDWIN BISHOP, M.D., Culworth.

JAMES T., a well-to-do farm-labourer, aged forty-two years, residing at Sulgrave, Northamptonshire, sent for me on March 3rd, 1865. The patient was a delicate looking man, with large brilliant eye, a hectic flush, and altogether bearing a phthisical aspect. On inquiring into his history, however, it appeared that none of his family had died of consumption, and that he had never been seriously ill but once before, which illness was an affection of the kidneys and bladder.

When I visited him, he had been ill several days from what he considered a severe cold and a "stitch." His symptoms were acute pain in the side, aggravated by painful inspiration, great difficulty in breathing, short harsh cough, quick pulse, the cheeks flushed, white tongue, and the urine scanty and high-coloured. I gave him a mixture containing two grains of tartarised antimony, a brisk purge of black draught; he was ordered to have repeated mustard and linseed-meal poultices, and a strict antiphlogistic diet.

In about four days his acute symptoms were much diminished; he continued to keep his bed, but never entirely lost his cough. He went on alternating from time to time, when the cough became much aggravated. He complained daily of shortness of breath, with flying pains about his chest and side, and his appearance altogether betokened serious mischief. He soon began to lose flesh, and his countenance became pale and anxious. There was great dyspnoea on the slightest exertion, pulse varying from 125 to 140 in a minute; tongue clean and red. Every evening, towards seven o'clock, fever came on, and was generally preceded by rigors terminating in profuse perspiration. He was unable to lie down in bed, and was supported by several pillows.

The expectoration at first appeared viscid and opaque, but soon assumed the character of purulent matter; from four to six ounces were discharged daily for a few weeks prior to the operation. His nights were passed almost entirely without sleep, unless he took half a grain of morphia. He was supported by good milk, mutton-broth, and eggs. In consequence of his not being able to lie on the right side, I was induced to examine the chest repeatedly; the right was evidently larger than the left, more particularly in the posterior surface. There was universal dullness on percussion in every posture, and on placing my hand on the swelling and desiring my patient to cough, I was fully convinced of the existence of a large quantity of fluid. As the poor fellow dreaded suffocation, I decided upon operating, though I had never seen the operation performed in such a case on the living subject. He was most anxious to be relieved from his sufferings, either by an operation, or death.

Ten weeks after my first visit to him, the evacuation of the matter was performed in the following manner: the patient was raised in bed as high as I could place him, his arms slightly elevated, the integuments were drawn up with the left hand, a deep incision was made with an ordinary bleeding lancet between the second and third rib on the posterior part of the chest, through which a full sized trocar, such as is used in tapping for hydrocele, was passed, and upwards of four pints of matter were slowly evacuated, to the great relief of the patient. After the trocar was withdrawn, a compress of lint was applied, and pressure made by means of a bandage. On the following morning I found he had requested his wife during the night to place him lower in bed, as his cough was easy, and the expectoration nearly ceased. In doing so, the compress was shifted, and at least three more pints of pus had escaped, for he was lying in a pool of it. He expressed himself greatly relieved by the evacuation of the matter, had passed a good night; he never coughed much after the operation; he changed from time to time, some days better, and again not so well; he was supported well with milk, broth, mutton, eggs, quinine, and cod-liver oil. For several months he was unequal to bodily exertion, and was urged not to attempt it. Six months after the operation, all appearance of pectoral disease was gone, and I often met him on the road with his team of horses. Yesterday (Good Friday) I entered into a long conversation respecting his present condition, which he described "as well as he ever felt in his life."

EXPERIMENTAL CINCHONA GARDEN IN NORTH CANARA. An experimental cinchona garden, established by the Bombay Government, at Gairsopah Falls, in North Canara, has met with such success that it has been proposed that another should be commenced in a still more favourable site on the Davimanny Ghaut.—*Bombay Times*.