Contributions

TO

PRACTICAL MIDWIFERY.

BY
ARTHUR B. STEELE, Esq.,

Lecturer on Midwifery, Liverpool Royal Infirmary School of
Medicine; Surgeon-Accoucheur to the Ladies'
Charity; etc.

I.—THE USE OF THE FORCEPS.*

LET us hear what eminent authorities, some even now living, have to say on the question as to when the forceps may or may not be used. Denman, who actually expressed a doubt "whether it would not have been happy for the world if no instrument had ever been contrived for, or recommended in, the practice of midwifery," says that before the forceps are applied "the head shall have rested as low as the perinæum for six hours." Dr. Robert Lee declares "that he never met with a case in which the forceps was satisfactorily applied before the os uteri was fully dilated, and an ear could be felt;" and in his published lectures he uses the following very decided language, "if you cannot pass your fingers completely round the whole head, you will do nothing but mischief with the forceps." (P. 308.) Dr. Meigs, the great authority in America, forbids their application until the os uteri is dilated and gone up over the head; he considers it unjustifiable to insert the forceps within the os uteri, or to use them in such a way as to produce compression of the head, with a view to lessen its dimensions and facilitate its descent through the pelvis. Dr. Ramsbotham, after laying down the general rule that, "if labour has lasted more than twenty-four hours from the rupture of the membranes, there is a great probability that instruments will be required," gives the following summary of the symptoms which warrant the use of the forceps, even though labour has not lasted the period of twenty-four hours:—"Gradual subsidence or disappearance of the pains, if the strength is failing, the spirits sinking, the countenance becoming anxious, the pulse 120, 130, or 140 in the minute, the tongue coated with a white slime or dry brown and raspy, rigors, green discharge, preternatural soreness of vulva, with heat and tumefaction of vagina, if the head have been locked for four hours and made no progress for six or eight hours." (*Obstetric Medicine and Surgery*. 1841: pp. 313-14.)

And, as if to make it more clear that the foregoing are the symptoms for which he would wait before interfering, he adds the following as the conditions which would indicate that interference had been injudiciously delayed; viz., "if the patient be vomiting a dark coffee-ground-like matter, if there be hurried breathing, delirium, or coldness of the extremities." I will quote one more of our highest obstetric authorities, Dr. Murphy, because the dogma which he adopts from Denman expresses forcibly and briefly the main errors to which I wish particularly to address myself. It is as follows:—"The lower the head of the child has descended, and the longer the use of the forceps is deferred, the easier in general will their application be, the success of the operation

more certain, the hazard of so doing mischief less." (*Lectures on Midwifery*, p. 250.)

Now, except perhaps the first clause, as to the facility of application, this teaching is precisely and directly contrary to that which I maintain is the sound principle upon which forceps delivery should be conducted.

The use of the forceps under the restrictions laid down by Drs. Murphy, Ramsbotham, and others, instead of lessening the perils of childbirth, would inevitably give rise to many of those mischievous and fatal complications, which their timely and appropriate application would as surely avert. I need not detain you by referring to many other authorities whose writings contain prohibitory conditions to the early and frequent use of the forceps, nor is it necessary to reproduce the graphic pictures they have given of the frightful injuries which the instrument is said to be capable of inflicting. I may remark, however, that the result of all this has been to create and perpetuate in the minds of a large body of accoucheurs in this country an unfounded prejudice against the forceps, and a repugnance to resort to them except under the most pressing urgency; and so it has come to pass that unfavourable results in practice have been attributed to the forceps, which in reality are owing to the mischief caused by deferring their application, until the local or general condition of the patient is such as to expose her to all the dangers and complications of protracted labour. It is but fair, however, to the reputation of many of those obstetricians who in their teaching place undue restrictions on the use of the forceps, to notice that in their practice they seem to follow a less prejudiced view; from which we may infer that their prohibitions are directed not so much against the skilful use of the forceps, as to deter those from having recourse to the instrument, who do not possess dexterity and experience sufficient to render it a perfectly safe instrument in their hands. It is obvious, however, that objections founded upon the abuse of any instrument should have no weight whatever in deciding upon its merits in competent hands.

If we turn from theoretical teaching to the records of actual practice, we obtain a more favourable view of the results of instrumental delivery. The maternal mortality after the use of forceps in this country, according to Dr. T. Smith, is one in twenty, according to Dr. Churchill, one in twenty-nine; the life of the child being preserved in nearly 75 per cent. In France and Germany, where the forceps are used nearly twice as often as in England, the mortality is one in thirty-four. The statistics of Mr. Harper, embracing upwards of a thousand cases in which the forceps were freely used, show that in unassisted tedious labour the mortality is one in twenty-two as to the mother, and one in five as to the child, while in forceps cases it is one in fifty-six of the mothers, and one in eighty-four of the children. The maternal mortality in craniotomy is estimated by Drs. Churchill and Tyler Smith, as one in five; in turning, as one in fifteen, and one in two and a half, as to the child; so that the maternal mortality after the use of the forceps is less than that of either craniotomy, turning, or even unassisted tedious labour.

With regard to the supposed dangers incurred in the use of the forceps; first, as to the mother, it has been alleged against them, that they are liable to injure the soft parts and to produce lacerations of the perinæum, recto-vesical and recto-vaginal fistulae, etc. Now, although these accidents may occur, especially if there be a want of skill on the part of the operator, or if the instruments be defective in construction, or if their use be deferred until the parts are in a congested or inflammatory condition, yet, with due

* Concluded from page 481 of JOURNAL for April 27th.

care, they may be avoided; and as a matter of fact, it is known, as Mr. Baker Brown and others have observed, that these accidents very rarely depend upon the use of instruments, but are the result of pressure in protracted and unassisted labour. Then, as to the child, it has been urged that the compression of the head frequently causes its death; and, as we have heard, some object to the use of the forceps as a compressor. It is, however, very doubtful whether under ordinary circumstances the compression of the head by the forceps is greater, and in many cases it is much less, than that which it frequently undergoes in unassisted labour, either when it has to pass through a strait pelvis, or when it presents in the occipito-posterior position. Every accoucheur has over and over again seen a child born alive, with a large head so moulded by strong uterine action, as to be converted from a sphere into a cone. Sir James Simpson has shown that the child's head will bear pressure in some directions better than in others, viz., in the biparietal than in the occipito-frontal; and it is in the former direction that the forceps act as compressors when properly applied. Mr. Harper, in his admirable paper in the first volume of the *Obstetric Transactions*, points out that the forceps do not surround the head and act as general compressors, but, owing to the yielding nature of the bones of the skull, when pressure is made upon two portions of the circumference, it is compensated for by the bulging out of the other portions, so that, unless the compressing power have been very great, the brain escapes injury.

Seeing, therefore, that on the one hand protracted labour is in itself dangerous both to mother and child, and that, on the other hand, the application of the forceps does not necessarily involve risk or injury to either one or the other, we conclude that it is in accordance with sound principles of conservative midwifery to encourage an earlier and more frequent recourse to this mode of delivery than is usually recommended by obstetric authors, and to discountenance the opinion that it is only to be regarded as a *dernier ressort* in extreme cases where the patient has already passed the confines of safe labour.

Adopting the proposition laid down by Mr. Harper, that, "the earlier and the more frequently the forceps are applied in proper cases, the more maternal and foetal lives will be saved, and the greater will be the freedom from the after consequences of protracted labour" (*Transactions Obstetrical Society*, Lond., vol. i, p. 180), we proceed to consider what are the circumstances which justify interference; what, in short, fulfils the conditions implied in the expression proper cases.

The decision of this point must of course in a great measure be left to the judgment and experience of the practitioner, and to the circumstances of each individual case. Dr. Churchill lays down the rule "that in no case is the forceps to be applied, until we are perfectly satisfied that the obstacle cannot be overcome by the natural powers, with safety to the mother and child." This is apparently a very simple and definite rule; but a little consideration will show that it is in reality extremely vague, and will find widely different interpretations, according to the views of different accoucheurs. Those who fully appreciate the evil results of protracted labour, will hold that its fulfilment implies interference at a very much earlier period than would be deemed proper by others, who maintain that the natural course of labour should not be interfered with, until certain definite symptoms of approaching danger have appeared. What I desire chiefly to enforce is, that the use of the forceps should not be deferred until bad symptoms have appeared, but that they should as a

general rule be applied at once, whenever the second stage of labour ceases to be actively progressive; excepting of course when contraction or distortion of the pelvis absolutely necessitates turning or craniotomy, or when the os uteri is only partially dilated, and at the same time undilatable.

I hold that it is not necessary to wait for six, four, or even one hour, so long as we are satisfied that the labour, if left to nature, will be tedious and protracted; a point which, I admit, may require much judgment and experience to determine. I altogether discard the rule that before the forceps are applied the ear should be felt, neither is it always necessary to wait until the os uteri is fully dilated; for with ordinary care the blades may be passed beyond the os uteri without inflicting the least injury, and with properly constructed forceps the os uteri is not stretched until the head itself is brought down to assist in the dilatation in the natural way.

When the fetus is known to be dead, although under certain circumstances craniotomy, as usually recommended, may be more suitable, yet, as a general rule, I prefer to deliver a dead child by the forceps, for two reasons; first, because, as we have seen, perforation is fatal in one case in five, while forceps mortality amounts to one only in thirty-three; and, secondly, assuming even that craniotomy is not more dangerous than the forceps, it is objectionable on other grounds as being, in appearance at least, a repulsive and somewhat barbarous proceeding, even though the child be dead. It must never be forgotten that, although we often have clear and positive evidence that the child is living, we cannot so surely determine that it is dead, although the usual signs of its vitality are absent.

I reject, as opposed both to theory and to experience, the notion that the forceps ought never to be used as compressors. I believe, on the contrary, that when properly applied and skilfully manipulated they conduce materially to the moulding of the head in its passage through a tight pelvis; and I have more than once proved in practice that a child may be delivered alive and unharmed, when a considerable amount of compressing force has been required. I advocate the early application of the forceps, because the earlier we apply them the better will be the position in which we grasp the head, and because we thus anticipate the disadvantages which ensue when the head is left to press for a long time on the soft parts of the mother, causing congestion and effusion, and so leaving less room for operating at a more advanced stage, besides rendering the vagina more liable to injury.

I recommend, as the best instrument for all cases, the long curved forceps of Simpson or Robertson; and on this point I speak with much confidence, from considerable practical experience, having found them more easy of application, and much more satisfactory in their effects, than the short forceps. I have for several years abandoned the use of any other except Robertson's forceps, and I feel quite sure that any practitioner, who will give them a trial, will arrive at the same conclusion. The chief points to be borne in mind in operating, besides the general directions of authors, are as follows: The operator should have constantly in his mind's eye, the curved line described by the head in its passage through the parturient canal, and direct his extractive force in accordance therewith; it will almost invariably be found that the rotation of the head as it passes through the pelvis, will be effected by nature, without assistance from the instruments, thus rendering any rotatory motion with the forceps unnecessary. The extraction should never be unduly hurried; nay, often, when the head is at the outlet it is necessary

even to delay it, and thus save the perinæum. This is a use of the forceps which I think has not been sufficiently dwelt upon, and which can only be satisfactorily accomplished by the long curved forceps.

I repudiate the dogma, that force is never to be used in artificial delivery; in employing it judiciously we imitate nature herself, and it is only when abused that it becomes dangerous. I have over and over again verified in practice that, with proper precaution, and with appropriate manipulation, delivery may be accomplished with safety to both mother and child, even when considerable force has been necessary to effect the extraction of the head.

I regard the use of the forceps as more appropriate in those cases in which many obstetricians recommend the ergot of rye, or other oxytoxics, for the following reasons: it gives the accoucheur a more certain, regular, and complete power over the course of labour, especially during the exit of the head, enabling him either to hasten or retard its progress as circumstances may require.

Their use does not involve certain risks which are admitted to attach to the action of ergot, even by those who advocate its employment in tedious labour. Dr. Johnson, for instance, says, "That when ergot is given it brings matters to a crisis, for if the child be not delivered within a certain time it will in all probability be destroyed." Drs. Hardy and McClintock state they "have on many occasions observed the ergot of rye to exercise a very depressing influence upon the maternal circulation." More recently Dr. McClintock has advocated the view, taught by Dr. Robert Lee when I was a student, that danger to the child arises from the incessant uterine action, and the consequent pressure, which the ergot is so liable to occasion. Dr. Churchill attributes to its use "retention of the placenta, from irregular uterine action, after the birth of the child." (*Theory and Practice of Midwifery*, 1866, p. 264.)

In my own practice I rarely use it in actual labour, preferring it as a means of completing abortion, when all hope of saving the ovum is abandoned, or of producing contraction in *post partum* hæmorrhage. It is my firm conviction that the more the forceps come into favour, the more rarely will ergot of rye be used, and that ultimately it will be altogether set aside as a substitute for what I cannot but regard as a safer and more efficient means of affording assistance in tedious labour.

In conclusion, I must beg your indulgence for many imperfections in this mere sketch of a very large and comprehensive subject. Those who desire to follow it out more completely than my limits on this occasion will permit, I would refer to the detailed observations of the authors I have quoted, and especially to two admirable papers in the first volume of the *Transactions of the Obstetrical Society of London*, by Dr. Tyler Smith and Mr. Harper.

I have purposely abstained from wearying you by a detail of cases; I may, however, observe that the views which I have brought forward are founded not upon theory, or the teachings of others alone, but are an exposition of the opinions I have formed during a somewhat lengthened and extensive course of practical experience as one of the accoucheurs to the Ladies' Charity of this town; an institution which comprises the results of about two thousand cases of labour annually, and those chiefly in a class of women especially liable to various forms of dystocia and other complications of parturition.

[ERRATA. At page 480, line 9 of second column, for "posterior", read "anterior"; and in the succeeding line, after the word "temple", insert "the other the posterior over the side of the occiput."]

REPORT OF

TWENTY-FOUR YEARS' VACCINATION IN THE MOLD DISTRICT OF THE HOLYWELL UNION.

By EDWARD T. HUGHES, M.D., etc.,
Public Vaccinator.

DURING the last twenty-four years I have reported to the Holywell Board of Guardians 10,344 successful vaccinations, and noted their influence in checking the spread of small-pox.

For the first five years, the Mold district consisted of three parishes; population 10,851; acreage 21,505. It was also the Mold Registration District. The registered births were 1535, and vaccinations 1340. Allowing for deaths previous to vaccination, this would have been a satisfactory number, had not a great majority of the children been above a year old, when vaccinated. Small-pox prevailed in some part of the district every year, and eighteen children born therein died of it, nine of whom were under one year, and only one above six years of age, all unvaccinated.

At the end of 1847, the parish of Northop was added to the Mold district, its population according to the census of 1851 being 14,577, and acreage 29,505. From January 1st, 1848, to December 31st, 1853, (when the Compulsory Vaccination Act had just come into force,) the number of registered births was about 2530, and my vaccinations 2609, of these 1303 were in children above a year old; so that, although the district had been on three occasions during the six years vaccinated almost from house to house, the arrears were considerable, as six hundred and forty-six of the children vaccinated were born before January 1st, 1848. These were well accounted for in subsequent vaccinations.

Twenty deaths from small-pox were registered during the same period; twelve of these occurred in the Mold Registration District, in which there had been but one fatal case, and that an imported one, from August 24th, 1847, to September 16th, 1852, when it was brought into Mold by the family of a knife-grinder. Five children died of it, four the children of tramps lately come into the town, and the fifth a sickly child sixteen months old, who had been unfit for vaccination from birth, and whose parents lived next to the house in which it first broke out. The other seven cases were in persons from seventeen to thirty years of age, all unvaccinated but one, who was said to have been vaccinated by her father, a village shopkeeper in Carnarvonshire, of which there was but faint proof. The other eight deaths in the parish of Northop, which forms part of the Flint Registration District, were all in young unvaccinated children.

My vaccinations during the remaining thirteen years have been 6395, and of these 403 were in children above a year old. The registered births must have been 7000 or thereabouts, but of these I have no trustworthy account. The deaths from small-pox were six, three in the Mold Registration District, and three in the parish of Mold, all in persons not vaccinated. To these I will again refer, as I propose now to consider the vaccinations of the last thirteen years, as illustrative of the working of the Compulsory Vaccination Act under very favourable circumstances.

The act to extend and make compulsory the practice of vaccination came into operation on October