towns in overcoming resistance to their duty, as also to divide with them the responsibility incurred by its non-performance? What an opportunity London affords at the present time of testing the value of such a proceeding! I have long felt that the present number of inspectors is too few, and their districts too large, to allow them, able and zealous as they are, to do much practical good to the cause of public vaccination; but if to each were assigned a certain number of subinspectors, on whom should devolve the responsibility of seeing the duties of the public vaccinators in their respective districts well and ably performed, and at stated periods of reporting thereon, I will venture to say that a great step towards providing for the vaccination of the people would be achieved, and that we should soon cease to hear reports of badly performed vaccination and impure lymph, the evils of which have been somewhat magnified.

Many other matters of importance relating to public vaccination have been repeatedly urged upon the Government by the President and members of the Epidemiological Society, Mr. Griffin, and others; and it is to be hoped that the amended Vaccination Act soon to be introduced into the House of Commons will show that they have not been passed by unheeded.

I cannot agree with those who think the registration of successful vaccinations unnecessary. The present system requires some amendment, as, to make it at all perfect, it is requisite that every entry in the register-books should in some manner be accounted for; and then they will form valuable records.

To carry out, however, any improved scheme of public vaccination, a larger grant of money will be required: and can it with any good reason, be withheld? Our country spends annually vast sums of money in devising and perfecting the most deadly weapons of destruction to protect her shores from foreign foes: why, then, should she give grudgingly for the more laudable purpose of saving her children from a loathsome enemy, whose victims we count annually by thousands, and against whom we have a certain means of defence, had we the wisdom and courage properly to use it?

CASE OF

ACUTE PLEURISY, TERMINATING IN EMPHYEMA: OPERATION: RECOVERY.

By EDWIN BISHOP, M.D., Culworth.

JAMES T., a well-to-do farm-labourer, aged forty-two years, residing at Sulgrave, Northamptonshire, sent for me on March 3rd, 1865. The patient was a delicate looking man, with large brilliant eye, a hectic flush, and altogether bearing a phthisical aspect. On inquiring into his history, however, it appeared that none of his family had died of consumption, and that he had never been seriously ill but once before, which illness was an affection of the kidneys and bladder.

When I visited him, he had been ill several days from what he considered a severe cold and a "stitch." His symptoms were acute pain in the side, aggravated by painful inspiration, great difficulty in breathing, short harsh cough, quick pulse, the cheeks flushed, white tongue, and the urine scanty and high-coloured. I gave him a mixture containing two grains of tartarised antimony, a brisk purge of black draught; he was ordered to have repeated mustard and linseed-meal poultices, and a strict antiphlogistic diet.

In about four days his acute symptoms were much diminished; he continued to keep his bed, but never entirely lost his cough. He went on alternating from time to time, when the cough became much aggravated. He complained daily of shortness of breath, with flying pains about his chest and side, and his appearance altogether betokened serious mischief. He soon began to lose flesh, and his countenance became pale and anxious. There was great dyspnoea on the slightest exertion, pulse varying from 125 to 140 in a minute; tongue clean and red. Every evening, towards seven o'clock, fever came on, and was generally preceded by rigors terminating in profuse perspiration. He was unable to lie down in bed, and was supported by several pillows.

The expectoration at first appeared viscid and opaque, but soon assumed the character of purulent matter; from four to six ounces were discharged daily for a few weeks prior to the operation. His nights were passed almost entirely without sleep, unless he took half a grain of morphia. He was supported by good milk, mutton-broth, and eggs. In consequence of his not being able to lie on the right side, I was induced to examine the chest repeatedly; the right was evidently larger than the left, more particularly in the posterior surface. There was universal dullness on percussion in every posture, and on placing my hand on the swelling and desiring my patient to cough, I was fully convinced of the existence of a large quantity of fluid. As the poor fellow dreaded suffocation, I decided upon operating, though I had never seen the operation performed in such a case on the living subject. He was most anxious to be relieved from his sufferings, either by an operation, or death.

Ten weeks after my first visit to him, the evacuation of the matter was performed in the following manner: the patient was raised in bed as high as I could place him, his arms slightly elevated, the integuments were drawn up with the left hand, a deep incision was made with an ordinary bleeding lancet between the second and third rib on the posterior part of the chest, through which a full sized trocar, such as is used in tapping for hydrocele, was passed, and upwards of four pints of matter were slowly evacuated, to the great relief of the patient. After the trocar was withdrawn, a compress of lint was applied, and pressure made by means of a bandage. On the following morning I found he had requested his wife during the night to place him lower in bed, as his cough was easy, and the expectoration nearly ceased. In doing so, the compress was shifted, and at least three more pints of pus had escaped, for he was lying in a pool of it. He expressed himself greatly relieved by the evacuation of the matter, had passed a good night; he never coughed much after the operation; he changed from time to time, some days better, and again not so well; he was supported well with milk, broth, mutton, eggs, quinine, and cod-liver oil. For several months he was unequal to bodily exertion, and was urged not to attempt it. Six months after the operation, all appearance of pectoral disease was gone, and I often met him on the road with his team of horses. Yesterday (Good Friday) I entered into a long conversation respecting his present condition, which he described "as well as he ever felt in his life."

EXPERIMENTAL CINCHONA GARDEN IN NORTH CANARA. An experimental cinchona garden, established by the Bombay Government, at Gairsopah Falls, in North Canara, has met with such success that it has been proposed that another should be commenced in a still more favourable site on the Davimanny Ghaut.—*Bombay Times*.