CLINICAL STUDIES OF CANCER.

BY CHARLES H. MOORE, F.R.C.S.,

Surgeon to the Middlesex and to St. Luke's Hospitals.

TREATMENT OF PRIMARY TUMOURS: DELAY.

GREAT as I deem the error of treating primary Cancers by the injection of acetic acid, it is small in comparison with the delusive trifling with them which occurs in the practice of certain persons, and especially in homoeopathy. When acetic acid fails, the disappointed surgeon resorts to the operation which he should have performed at the first. But, in other practice to which I now refer, inadequate or negative treatment is sometimes persisted in until tumours have become well nigh or wholly unsuited for operation. It has been usual, I believe, for the homoeopathic practitioners to confine their treatment to internal diseases, which combine the advantage of definite names with indefinite outward signs. But lately—for how long I do not know—some of these gentlemen have ventured with their globules to assail external tumours. Some patients who have been so treated have complained to me loudly of their wasted confidence; while others, gathering hope from the doubtful changes which occurred in their tumour from week to week, rather than recognising its manifold increase in the whole period of treatment, have appeared to cherish a lingering persuasion that homoeopathy might even yet cure them. It is hard to tell by what foregone professional mishaps, or under what anti-professional influence, the opinions of these persons have been so perverted; but there is among them a surprisingly inflexible confidence in irregular ways. While actually abandoning one form of quackery which has beguiled them nearly to death, they will take our opinion of the nature and the requisite treatment of their cancerous tumour, that they may with the surer reliance betake themselves to quackery again.

It might be useless, not to say improper, to write thus, if the subject were less serious. In the medical practice of the drawing-room, the physicking of those who are sound, there may be practically little harm in homoeopathy. But the case is different when a patient confides to a practitioner the dispersal of her cancerous tumour. There is no power in any treatment to cure a disease of that nature; and the attempt to do so beyond its early and doubtful stage—certainly a prolonged attempt—demands a hardy conscience.

To what extent may success be looked for in the treatment of primary Cancer? It is a fact, both that small variations take place in the apparent size of a tumour, and also that a certain diminution of its bulk may sometimes be obtained by improving the general health, or by suitable local applications. And this statement is not to be confined to innocent tumours, but is true of those also which are cancerous. Indeed, the greater perfection attained in the growth of an innocent tumour confers on it a more stable vitality than is possessed by the elements of Cancer, which are at once more multiplied and less mature. Approaching the natural textures in the quality of elaboration, an innocent tumour imitates them in the resistance to artificial destruction, and is therefore much less than Cancer is under the control of remedies. But the diminution of a cancerous tumour which follows upon suitable treatment, apart from the direct effect of local applications upon the

skin, is not mainly due to the removal of essential constituents of the tumour. It may be doubted if any part of it which retains the germinating power can be thus destroyed. Contained liquid may be absorbed, surrounding oedema may disperse, and to that extent the growing of the tumour may be arrested, as well as its size reduced. It may be said that, in some cases, a texture once stiff with Cancer becomes supple and all but sound again. Even there, however, a residue of the living cancerous element still exists, which may at any time break forth in a renewed growth. Such temporary and partial reduction of a cancerous tumour affords no justification for pursuing inadequate treatment, when the nature of the disease has been recognised and the tumour may be otherwise removed. But what shall be said of a reckless dallying with growing tumours, such as is illustrated in the following two cases? Surely fairness towards our patients requires that we should neither persuade them to adopt useless treatment in the stead of that which is efficient, nor practically consent to their pursuing it when it is hopeless. The connivance of the practitioner in such proceeding results in a fatal, though unseen, dissemination of Cancer into adjoining healthy structures, or in a serious increase of the extent and risk of the inevitable operation.

A widow, aged 58, whose life had been always healthy, found a tumour of the size of a pea in the right breast. She committed herself to homoeopathic treatment for nine months, after which I saw her. Deep in the upper part of the right breast was an oval tumour, having its long direction horizontal, in size between that of a hen's egg and that of a goose's egg. It was slightly granular on its surface and hard. Though feeling moveable, it was in fact closely connected with the breast-tissue, the looseness of which permitted the tumour to roll about. The skin was healthy and still unattached to it; the nipple could be drawn out; there was no adhesion to the pectoral muscle, and no trace of glandular disease. The tumour was sometimes the seat of shooting pain, and it ached after handling.

Satisfied of the cancerous nature of this disease, I advised that it should be removed. The patient adopted my advice; but, notwithstanding her disappointment with homoeopathy, she continued true to the feeling of distrust which she had conceived towards the profession, and she resorted to some unrecognised practitioner for what operation he might practise upon her.

The left breast of a delicate lady, under 40 years of age, contained a tumour of the size of a small cocoon. During two years, in which this tumour had grown from the size of a walnut to its present dimensions, the patient had taken homoeopathic medicines, and had not only encouraged herself to hope that they might reduce the tumour, but had apparently failed to notice the fact of its steady increase. It was lobed, convex, inelastic, firm, and tender. The skin adhered closely to it in two parts, and was there red, thin, shining and tense. Large veins passed from this portion of the tumour over the healthy part of the breast. The mass moved freely upon the pectoral muscle, and there was no trace of axillary disease.

It was still possible to remove this breast, and I gave advice that it should be done.

A disastrous delay arose in the following case also, from misplaced confidence in the efficacy of treatment. Whether the error were on the part of the patient or of his medical adviser, a man was allowed to get worse, and so much worse that it was doubtful if his case must not be abandoned, when a double operation rescued him from early death, and restored

The procedure was so well tolerated, a case of extreme pain, a sharp shooting of the optic nerve, or a light in the pupil, did not ensue in the first experiment, and continued his observations every day. "By the time the action of the atropin has subsided, he will, if endowed with good vision and some manual tact, find himself master of a step in ophthalmic investigation which had hitherto been considered by him as insurmountable.