narcotic properties are beneficial in certain conditions arising in acute disease.

10. No remedy which is potent for good can be impotent for evil.

CASES OF DUODENAL PERFORATION.*

By ANDREW CLARK, M.D., F.R.C.P.,

Physician to and Lecturer on Medicine at the London Hospital.

CASE V. Of this case I possess only imperfect notes; but the main facts were these. A girl, about 10 years old, was brought to the London Hospital for an extensive scald. She did well till about the fourteenth day after admission, when she began to complain of pain in the right side of the belly, shooting through to the back. This continued for nearly three days, sometimes better, sometimes worse. At the close of the third day, peritonitis set in, and ended fatally within twenty-four hours. After death, it was discovered that the peritonitis had arisen in consequence of the occurrence of perforation in one of Curling's ulcers of the duodenum near the pyloric extremity of the stomach.

CASE VI. W. T., 43 years old, a short, stout, but pale and flabby looking man, was admitted into the London Hospital as an extra case, about the middle of July, complaining of intense pain, increased by pressure, on the right side of abdomen, occasionally shooting through to the back, of sickness, and of a sense of extreme prostration.

The patient had been for some months liable to pain in the right hypochondrium, coming on about two hours after food. The pain had never before been severe, and never before had either arisen or subsided suddenly. There was no history of jaundice, hæmaturia, or gravel. The patient's present attack came on quite suddenly, shortly after a hearty meal taken four hours before admission.

When first seen by me, the man lay upon his back, with his knees drawn up and slightly apart. The face was dusky-pallid and anxious; the breathing hurried and costal; the skin bedewed at parts with a cold sweat; the manner restless and dejected. The belly was flat; and no pressure could be borne in the region of pain, which extended from below the right hypochondrium to near the crest of the ilium, and was felt at intervals under the lower angle of the right scapula. The tongue, yellow-furred at base, was red and eroded at tip and edges. There was occasional vomiting of greenish frothy mucus streaked with blood. The introduction of water into the stomach produced no increase of pain. The bowels had been freely and naturally relieved the day before. The urine was normal. No disease could be detected in the heart, lungs, or brain. The pulse was about 100, small and somewhat sharp.

Plainly enough, this was no case of intestinal colic. Was it, then, one of hepatic or of renal neuralgia due to the presence of a calculus, or arising independently of one? Here was an assemblage and order of symptoms not uncommonly attendant upon the passage of a gall-stone into the bowels. Why, in the present instance, then, were not these symptoms ascribed to this cause? In the first place, because the chief seat of pain was not over the liver, but someway below it, and because there was neither swelling nor tenderness over the gall-bladder. You could percuss the liver and press its free margin,

without adding to the patient's sufferings; but, when the hand was laid upon a spot about an inch below that and a little to the right of the umbilicus, the pain, throughout severe, became at once insufferable; moreover, it had arisen and remained in that spot. In the second place, though the absence of jaundice was no disproof of the presence or passage of a gall-stone; the absence of bile from the urine, and its presence in the faeces, showed that the biliary passages were free. In the third place, a number of minor circumstances concurred in disavowing the notion of a gall-stone as the cause of the patient's symptoms. Among these were the history of painful digestion with an eroded tongue, the well defined position and fixity of the abdominal pain, the quick small pulse, and the partial collapse.

In the light of these considerations, it seemed impossible to ascribe the patient's symptoms to any disease of the liver. Did the fault, then, lie in the kidney, and was the cause a renal calculus? I concluded not. There was no pain, and there had been none, in the region of the kidney; no retraction of the testicle; and no numbness or pain in the right thigh. The urine was healthy and abundant; and there was no history of hæmaturia or gravel. Having put the kidney out of the question, I came to the conclusion, that I was dealing with a case of ulceration of the duodenum in which a slight leakage had occurred. Acting upon this conviction, the man was ordered to be kept scrupulously quiet on his back in bed; and to take no food or drink by the stomach for forty-eight hours. A small pebble was kept in the mouth to allay thirst; the skin was sponged with tepid water; and nutritive injections were administered by the bowels. He was put freely under the influence of opium; and hot flannels sprinkled with turpentine were kept constantly applied to the abdomen. For the first fourteen hours, there was no improvement in the patient's condition; but, after that time, he began and continued to amend. At the end of thirty-six hours, I was obliged to forego the rigid abstinence which had been practised up to that time; ounce doses of milk were allowed to be taken every hour. On the third day, the patient was much better; the pain had subsided; the countenance had become more natural, the skin warm, the breathing deeper, the pulse fuller; vomiting had ceased; and the weight of the hand could be borne on the abdomen. Milk and beef-tea were now allowed more freely in small quantities at short intervals. In a week, the patient had become so much better as to be able to sit up. In a fortnight, he expressed himself as well, though confined to a sloppy and almost exclusively milk diet. There was still, however, uneasiness after food; and pressure could not be borne along the former seat of pain. The patient now took to nursing; and, after a few weeks' careful dieting upon pultaceous food, abstinence from stimuli, counterirritation to the epigastrium, and the use of bismuth, left the hospital, as I understand, quite well.*

Whilst the last-mentioned case was under treatment, a woman was admitted into Redman Ward suffering from somewhat analogous symptoms, but arising from quite a different cause, and, for the sake of comparison therefore, worthy of brief narration in this place. She was 42 years old, had a fat flabby body, a sallow anxious countenance, and a slight icteric tinge in both conjunctivæ. Tossing restlessly about in bed, and uttering loud and frequent groans, she complained of agonising pain in the pit of the stomach, shooting through to the back and upwards

* Concluded from page 689 of last number.

* Another case, similar in all essential points, was admitted into my wards at the hospital in the beginning of May, and has just been discharged cured.

into the chest and shoulders. The pain ebbed and flowed in severity, and was accompanied by a sense of constriction round the body, by chilliness, and by occasional sensations of nausea and faintness. The abdomen was retracted and covered with a cold sweat. The epigastric and right hypochondriac regions were tender to pressure, which, when considerable, induced severe pains in the chest and shoulders. The feces were reported to be pale. The urine obviously contained a good deal of bile. The pulse was about 62, small and feeble. The hands and feet were cold. There was no history of indigestion, or of any former attack like the present, which had come on the day before when the stomach was empty. Patient also noticed, of her own accord, that the pain was lower, and stretched further to the right side, than when it first arose. Moreover, though suffering atrocious pain, the woman was keenly alive to all that was passing around her; there was none of that introversion of thought which, when present, excites one's suspicions of grave disaster within.

Taking all these facts into consideration, especially the remitting character of the pain and its change of place, the tenderness of the hepatic region, the contraction of the abdomen, the absence of bile from the feces and its presence in the urine, the jaundiced aspect of the conjunctive, and the slowness of pulse, I concluded that I had to deal with a case of biliary calculus, and treated it by means of warm drinks, full doses of opium (with which a little antimony was conjoined), and hot fomentations to the belly. Next day, the woman became slightly jaundiced; the pain shifted lower down; and then, towards evening, suddenly ceased. In another day, she was quite well.

A review of the cases of duodenal perforation here recorded, however imperfectly, brings into striking relief some important points in the diagnosis and treatment of this terrible accident; and with a short summary of them I shall close this communication.

First with respect to diagnosis: here are the points which, when present together, appear conclusive as to the occurrence of perforation of the duodenum:—

- a. A history of more or less painful digestion.
 - b. The sudden occurrence of pain after food a little to the right of the mesial line, and somewhere between the liver above and the crest of the ileum below.
 - c. The continuity of character and fixity of place of this pain, which may increase, but does not remit, and which may extend in almost any direction, but never quite leaves its place of origin.
 - d. Swelling and tenderness of belly; costal breathing; a quick, small, feeble pulse; clammy sweats; pinching of features; and dusky skin.
- The next event, if death does not occur from collapse, is the setting in of general peritonitis, which does away with any further doubt concerning the nature of the case.

The feeling of something having given way within, followed by a sensation of liquid heat diffusing itself for some distance in the belly, often experienced in cases of perforation, was not observed in any of the patients whose histories are recorded above. It is not improbable, therefore, that the presence or absence of this feeling may be made a means of distinguishing between a large and small perforation—between one for which we entertain some hope of recovery, and one for which there can be none at all.

Secondly, with respect to treatment, it is plain, if we aim at recovery (which, if the perforation be very small or imperfect, we may fairly do), that we must keep the parts quiet; that we must prevent the ingestion of food (liquid or solid) which may escape into the peritoneum; and that we must not only

allay pain, but soothe that organic unrest which springs out of such accidents, and puts the patient in present peril of their worst effects. It seems to me that these indications will be most happily fulfilled by keeping the patient scrupulously quiet on his back or left side in bed; by allowing nothing to be taken by the mouth for the first forty-eight hours after the accident, or for so long as may be consistent with safety; by giving opium in the way of hypodermic injection; by the application of hot moist cloths to the skin for the supply of fluid to the body; and by the use of nutritive and stimulating enemata, if there be need of such support.

My clinical and pathological experience quite warrants the opinion that there are cases of minute perforation of the duodenum which might be saved by the timely practice of such rules of treatment as I have touched upon above.

Reports OF HOSPITAL PRACTICE: METROPOLITAN AND PROVINCIAL.

BIRMINGHAM GENERAL HOSPITAL.

THREE CASES OF HEMIPLEGIC EPILEPSY, EACH PRESENTING A DIFFERENT GROUP OF SYMPTOMS.

(Under the care of Dr. RUSSELL.)

I APPLY the term epilepsy to the cases which follow in compliance with what I believe to be the better use of this term, suggested by Dr. Jackson; viz., to indicate a condition of the nervous tissue, rather than a special group of symptoms. The different attacks in the three patients presented all the suddenness of the epileptic paroxysm, and its brevity; in each instance, fits of the ordinary epileptic type also occurred; finally, the three cases, read together, will I think afford powerful support to the hypothesis that we may refer the condition of the nerve-tissue in all these attacks, to a similar occurrence, the running down of nerve power.

In this latter point of view, the three cases bear a highly interesting relation to each other, and strengthen the opinion that spasm and neuralgic pain are both attributable to the same condition of the centres with that which originates paralysis and anaesthesia. Taking the cases together, we have the three conditions, spasm, paralysis, and pain, each one prominent in a particular case, but more or less associated with the other two symptoms. Thus in the first case, spasm takes the lead, with paralysis waiting upon it, and anaesthesia; and what is very much to our present purpose, on one occasion hyperaesthesia actually took the place of the anaesthesia. In the second case, paralysis is the main symptom, with anaesthesia and pain, the pain occupying a very subordinate position. In the third, pain is the chief phenomenon; anaesthesia and paralysis are also present, but in a less prominent place. In both of the last two cases, there was also a tendency to spasmodic action.

But not only were these different symptoms present in the same case, thereby sufficiently indicating their connexion with each other; but one symptom actually passed into the other. Such was especially the case with the first patient, in whom the spasms