perience in consultation felt assured that their professional interests and reputation were perfectly safe in his hands. I have to add, that Mr. Smith was formerly on the Council of this Society, and at the time of his death was honorary local secretary for Leeds.

My next desire, gentlemen, is to return you my best thanks for your earnest and well directed endeavours to promote the welfare and usefulness of this Society; and to the honorary secretaries our cordial acknowledgments are also most undoubtedly due for their indefatigable and valuable services during the year. And now, in conclusion, let me venture to hope, as we have indeed reasonable grounds to expect, that our Society will be the means of knitting together, and firmly, the bonds of good fellowship amongst us, and, when our work is finished, amongst our successors for all time. That, in its abundant and valuable fruits, our Society will ever, in pursuance of its high object, continue to prosper for the good of those tender beings who have the strongest claim upon our sympathy, whose safety it should ever be our anxious care, if not our sacred duty, to watch over, help, and comfort, in their peculiar bodily afflictions and sufferings.

ARE THE TYPES OF FEVER LIMITED?*

By HENRY KENNEDY, A.B., M.B., Dublin.

THOUGH the subject of fever be one of great extent, I promise to make the following remarks very brief, and to avoid, as far as possible, ground which has been already trodden. My object now, in venturing to come before the British Medical Association, is to advance a few general considerations on the several types of fever. I have the strongest convictions that these types are much more mixed up, intermingled, as it were, than is commonly supposed, or than books teach. That they are more numerous, too, than the works on the subject describe, I cannot doubt; and I may give, as an illustration, the type of fever known as "gastric"—a type very well marked in itself, described by many English writers of the last and of the present century, and yet a type which seems to have died out, if I may so express it, in London and other cities; for I can find no notice of it in any of the recent works which treat of fever. How this is to be explained, it is not easy to say. With us in Dublin, it may be said to be always present, though it must be allowed to prevail much more at some seasons than others.

To this type of fever, then, known by the name of "gastric," I would ask special attention. Though often met as quite a distinct type in itself, and as distinct as typhus is from typhoid, it yet seems to me to form a connecting link, as it were, between the two. It would be quite out of place here to enter on a description of this type of fever. I shall only, then, state that the leading symptoms are, a red tongue, frequently sore throat, a desire for cold drinks, nausea, vomiting, pain on pressure over the stomach, and all these attended by a very prolonged fever, at least as long as typhoid. In very severe cases, too, the tongue will become quite dry, and there will be marked tympany; whilst the brain will be much excited. Now, within the last three years I have twice, at intervals, had occasion to note a large number of cases of typhus, as shown by the presence of a copious rash, where this gastric type of fever was also present. The duration of these cases was something remarkable, being far beyond the ordinary typhus. Here, then, was a very striking fact, proving, as it seems to me, that these two types of fever may be conjoined, though usually quite distinct and separate.

But this gastric type of fever sometimes presented other features, showing how difficult it is to draw a boundary between one type of fever and another. Within the last year, measles prevailed much in Dublin; and in many cases, and before the patients had left their beds, was followed by an attack of gastric fever, which proved most obstinate. One of these I saw with Mr. McCarthy of Caple Street. The patient was a boy of nine years old. But it was not because gastric fever existed, that I notice it here, but because, during this fever, a well-marked eruption of typhoid spots made their appearance, and yet there was no other sign whatever of the ileum being engaged. The spots, I should add, were bright red, well defined, and limited to the abdomen and sides of the chest. Nor was this at all a solitary instance of gastric fever with typhoid spots. In a very prolonged case of this kind which I saw with my friend Dr. Johnstone, not only one, but two distinct crops of eruption appeared in the progress of the case. The rash might here have been taken as a model of what appears in enteric fever, and yet no

other symptoms of this type of fever existed. The bowels were much confined all through the attack. Another case, too, I saw with my friend Dr. Denham. It occurred in a boy of five years old, and was a very aggravated case of gastric fever. In its progress, when the boy was now ten days ill, a well-marked crop of spots, having all the characters of the typhoid rash, appeared. Yet, though looked for, not another symptom of enteric fever could be discovered. Lastly, I may allude to the case of a lady aged 27, whom I saw in the spring of the present year, in consultation with my friend Dr. J. B. Montgomery of Penzance. She was attacked with gastric fever in a very marked form. The nausea continued for the greater part of a week; and during the first two days there was diarrhoea; but the discharges were not characteristic of enteric fever, nor did it recur, though the fever ran out to the twenty-fifth day. When the patient was nine days ill, a very well marked typhoid rash appeared; but the spots did not exceed seven or eight altogether. Still they were well pronounced, and could not be mistaken. As it was thought advisable, a sketch of this case, but much fuller than that now given, was sent to an eminent London physician, who pronounced it to be one of "modified typhoid"; which, for my part, I must say, appeared to me an excellently chosen name, and bore out well, though coming from a London physician, what it is the object of these remarks to prove—viz., that the types of fever do not always exist as isolated types, in the way that books describe, but that it is by no means uncommon to find them conjoined, or the one gliding insensibly into the other. That this is so at times, as regards typhus and gastric fevers, I cannot doubt; neither can I doubt that gastric and typhoid may be met together; whilst the peculiar spots, on the presence of which so many place great weight as being characteristic of enteric fever, are, beyond question, not limited to that type of fever, as the cases alluded to this day sufficiently prove; and I could have given several others, had it been necessary.

In close connexion with the line of argument I have been pursuing, I may advert to the epidemic fever which prevailed so generally, and in so many countries, in the years 1847-8. With us in Dublin, it was essentially what is known as the relapsing fever; that is, it was made up of two parts. A sharp attack of ordinary fever, not presenting any special symptoms, occurred, and ran on for five, seven, or nine days; a complete lull would then occur, very often apparently brought about by perspiration. The patient would ask for food, and say he was quite well. But the improvement was only in appearance—not real; for, at an interval varying from a day to seven or even more, fever again set in, invariably of a much more severe character than the first, and running a longer course. It did not appear to me possible to look upon the two attacks in any other light than the one being but a part of the other. No other view, indeed, of the matter would have constituted the fever a relapsing one. And yet the two attacks frequently presented the most marked contrasts, not only in severity and duration, to which I have already alluded, but in the much more distinctive character of the absence of spots in the first, and their presence in the second part of the fever. In fact, a fever, beginning as an ordinary continued one, very frequently, after the lull, merged into genuine typhus, attended by a copious petechial rash. I know not that I could advance anything which, to my mind, would more conclusively prove that the types of fever are very much more united than is commonly thought; and that even the existence of spots, usually considered so characteristic of certain types of fever, does not prove the contrary. I shall be glad to hear the opinion of any gentleman on this special point.

In another and a very different aspect I would speak of this subject: I mean in relation to the morbid anatomy of enteric fever. It seems to me that writers have taken too narrow a view of it, when they speak of the affection of the lower part of the ileum and enlargement of the contiguous glands as being the essence of the disease. It is quite notorious that these morbid states frequently do not bear a relation with the symptoms present during life. I have seen cases where the morbid appearances were very slight, though not to be mistaken; and yet the fever was so severe as to cause the death of the patient. In fact, it was the fever that killed the patient, not the local lesion. Hence I am led to the conclusion—and it appears to me a fair one—that enteric fever is much more of an essential fever than is commonly thought; and that a patient may even die of this fever, and yet not a trace of morbid lesion be found after death. On the other hand, when the lower portion of the ileum is profoundly engaged, it seems to have been overlooked that it frequently does not constitute the whole of the disease. On the contrary, the colon has been ulcerated, and, in my own experience, more frequently far than was met with by Louis. In a specimen which I showed last winter, the ulcers in the colon predominated, though there were also ulcers in the ileum, and enlarged glands. The symptoms during life were those of well-marked typhoid fever. In addition to all this, Louis has shown that other parts of the intestinal canal, but more particularly

* Read in the Medical Section at the Annual Meeting of the British Medical Association in Dublin, August 1867.

the stomach, are often engaged in this type of fever; to which, amongst other names, he gives that of gastro-enteric. Is it any wonder, I ask, with such a variety of morbid lesions, whether of the slightest on the one hand, or of the most profound and extensive on the other—is it any wonder that the symptoms during life should vary in a very marked way, or exhibit symptoms having the closest resemblance to gastric fever, or even to typhus, as I myself have often seen? And have not the symptoms described as characteristic of enteric fever been too circumscribed? As examples, I may mention bleeding from the nose or bowels, both of which, beyond all question, occur in other types of fever besides the enteric, as I have proved in another place. Were it necessary, it would be easy to pursue this part of my subject much farther; but I must hasten on.

From the tenor of the preceding remarks, it will be understood what my object has been. I wished to show that, while the types of fever may all usually be distinguished the one from the other, yet they are frequently conjoined, or glide the one into the other. I cannot doubt that this is the case as regards gastric and enteric fever; neither can I doubt that the relapsing fever and typhoid may coexist. I believe, too—and here I would wish to speak with much more reserve, on account of the many able physicians who differ from me—that I have often had to treat cases of typhus and typhoid conjoined. When speaking of this subject in former papers, I detailed a number of cases of this kind; but I have purposely avoided doing so now, not because I wanted cases of the kind, for during the past year several of great interest have come under my notice, but because I have given many similar cases in former papers. There are, however, two points which appear to me worthy of a special notice. In two cases within the last year, the treatment adopted to check the diarrhoea of enteric fever proved somewhat too potent; but what was the effect on the patients? It literally changed, as I believe, the very type of fever before my eyes; and what had begun with all the signs of enteric fever ended literally in typhus. In each case, the diarrhoea (and I may state, in passing, the discharges were most characteristic) suddenly ceased, and the brain became seriously engaged. One case was a man of thirty-two years of age. Though he presented no petechia, he had otherwise well-marked typhus; but his life was at no time in danger. The second was a woman, aged twenty-six. She had all the signs of enteric fever, and in a very severe form. The diarrhoea was most profuse; and, what I believe to be most characteristic of the disease, the quantity of fluid passed was out of all proportion greater than what she took. This patient had a few well-marked spots on the chest and abdomen. They were very well defined, but were a shade darker than those usually met in enteric fever. On the cessation of the diarrhoea, the patient's brain became much engaged; the eyes were deeply suffused; she raved constantly, and was scarcely to be kept in bed. In this state, her life was in the balance for many days; but she finally made a good, though an unusually slow recovery. This case was witnessed by Dr. Grimshaw. These two cases open up a new phase of the subject, as showing how one type of fever may be converted into another. Nor are such cases without parallel; and more particularly in acute disease, where the attack, beginning as inflammatory fever, often, and very suddenly, puts on all the characters of typhus. But the subject is too extensive to follow out further in this place.

The second point which appears to me worthy of notice is the fact that, in those cases of fever where, as I believe, typhus and typhoid co-existed, the duration of the attack was much longer than if the disease had been typhus alone, by a period of at least ten days. In other words, the attack, as far as duration went, occupied the number of days which enteric fever usually runs—that is, five- or six-and-twenty days. This fact appears to me one of moment, as showing, as far as the nature of the subject admits, that a something was present in these cases which prevented the attack ending in fourteen or fifteen days, which is the common course of typhus; and that these cases partook of this character, was proved by the presence of a copious petechial rash. I must leave to those who hold different views from my own the explanation of this fact.

In the foregoing remarks, I have purposely avoided going over again old ground. I might, however, have given a number of additional cases where enteric and typhus fevers seemed to me to be conjoined. For the present, however, I will hold these over for another occasion. I shall only observe in reference to them, that they appear to me much more common at one period than at another; and this is quite in keeping with what we might expect, for the phases of fever are ever varying. Each month differs from its fellow, and thus I have met many more cases of conjoined types at one time than another. The same has occurred to others; and Huss states, in the most explicit terms, that he met typhus and typhoid conjoined, and occurring under identical circumstances. On the contrary, it seems to me more than probable that, in some localities, the types are not met conjoined, but present themselves

separately, just as has been described by the London physicians. But that the time will come when the diseases will most probably present themselves under the aspects seen by Huss and myself, I can scarcely doubt. For a long period, it was asserted that enteric fever did not exist in India. Now, it is known to prevail there, and, if I recollect right, has also been described as being met in the conjoined form. This state of things was also met during the Crimean war, and has been particularly described.

The conclusion, then, at which I arrive from these facts, is, that the types of fever are often conjoined; and that, if this be so, if the gastric type and typhus, and the gastric and enteric, be found so mixed up, there is no straining of analogy in supposing that typhus and typhoid may coexist, as I firmly believe I have often seen. With this remark, I leave the matter for the present.

NOTES OF AN AUTUMN HOLIDAY IN AMERICA.

By T. SPENCER WELLS, F.R.C.S.,

Surgeon to Her Majesty's Household, and to the Samaritan Hospital.

I.

A HOLIDAY in August or September is now happily becoming very general amongst us. The town doctor takes his family to the country or to the seaside. He gets some fishing or shooting with country friends in England; perhaps in Wales, Scotland, or Ireland. He tries some French or German watering-place; visits the Alps or Pyrenees; or perhaps does a little yachting. Somehow or another, he gets rest, sunshine, and what a learned friend of mine says is not less valuable than change of air—namely, *change of mind*. Having run through all the more common varieties of a doctor's holiday, I determined last autumn to gratify a long standing desire to see the falls of Niagara, and some of the great lakes and rivers and forests and mountains across the Atlantic, and visit some of the chief cities which have so rapidly grown up, and at the same time to make the personal acquaintance of some of our fellow-workmen known before only by their names and writings. Consulting with various American friends who have visited London—with Sir Henry Holland, who has repeatedly crossed the Atlantic—and by the aid of the American Bradshaw, *Appleton's Railway Guide*—I found that a great deal could be done in six or seven weeks; and such is the perfection of the ocean mail-steamers, and the regularity of the steamboats and railways in Canada and the United States, that the plan marked out before leaving England was followed with scarcely a day's alteration. For the information of any one who may wish to enjoy a similar holiday, I have noted exactly what we did comfortably in seven weeks. I say we, because my wife and one of my daughters accompanied me.

August 15th. Left Liverpool.

August 27th. Arrived at Quebec.

August 29th. On St. Lawrence to Montreal.

August 30th to September 1st. At Montreal.

September 2nd. Up river to Ottawa.

September 3rd. At Ottawa.

Sept. 4th and 5th. By the Thousand Islands and Lake Ontario to

September 6th. Niagara. Stayed there till

September 9th. Travelled by night train to

September 10th. The Catskill Mountains.

September 12th. By Hudson River to New York.

September 13th to 17th. Philadelphia.

September 17th to 20th. New York.

September 21st. New Haven.

September 22nd to 25th. Lowell and Boston.

September 26th. Halifax. Sailed and arrived

October 5th. At Liverpool.

By this arrangement, no part of the country is travelled over twice, except between New York and Philadelphia, and that is varied by choosing different routes in going and returning. By going first to Canada, the cooler country is visited during the hottest part of the holiday, leaving New York and Philadelphia till the latter part of September, when the great heats are over. By making the holiday a fortnight or a month later, the autumnal tints in the American foliage would be seen to greater advantage; but this would not accord so well with London seasons, and the return Atlantic voyage would probably be more