why that lesion is unlikely to happen frequently. Be it noted, that the displacement observed was primary, taking place at once in the original injury, and of such a kind as could scarcely be produced after lodgment of the head of the humerus in the upper part of the axilla, without injury to the thoracic walls or the shoulder-girdle. While the patient was walking on crutches, his right crutch slipped; he lost his footing; and, the whole weight of his body being thus thrown on the left crutch, the head of that crutch glided outwards from the axilla, and raised his elbow as he fell to the ground. This is the patient's story. The head of the crutch was the fulcrum on which the dislocation took place; and, as it glided outwards, it gave more and more leverage to the forces by which the head of the humerus was depressed—viz., the action of the latissimus dorsi and pectoralis major muscles in the commencement of the process, and the weight of the patient's body throughout. But, in the ordinary dislocation forwards and downwards, the elbow or hand comes into contact with the ground while the head of the humerus is at a higher level, with the weight of the body over it; and the latissimus dorsi and pectoralis major muscles, which are important agents in producing the dislocation, likewise draw the arm and the trunk together, and pull the head of the humerus to the deep part of the armpit. Now, in the present instance, although these muscles doubtless aided in withdrawing the head of the humerus from the glenoid fossa, they were immediately thereafter with great rapidity opposed by the descending weight of the body, which, pressing the acromion against the displaced neck of the humerus, caused the head of the bone to pass between the insertions of the muscles in question, undoing the fold in the tendon of each, and reversing, or more probably destroying, the action of both.

HYPODERMIC INJECTION OF MORPHIA IN A CASE OF FIBROID TUMOUR OF THE UTERUS.*

By SAMUEL MILLS, Esq.,
House-Surgeon to the County Hospital, Lincoln.

It is perhaps worthy of mention that great relief was afforded during life by means of the hypodermic method of injection of morphia to a poor woman (S. G., aged 40) who was lately an in-patient in this hospital, suffering from a very large and irregular fibroid tumour of the uterus. One of the main features of the case, and in fact the one which I am anxious to bring before this meeting, is the pain which she suffered during the last fortnight of her existence; pain of the most dreadful and agonising character during its continuance, and apparently caused by efforts on the part of the uterus to expel the fibroid mass from its interior. The pain was almost of daily occurrence, and, unless speedy relief was afforded her, would continue unabated for six or eight hours at once. In these attacks, she presented more the appearance of a woman in very severe labour than any other state with which I could compare her. She complained of the pain being always either in "her body" over the seat of the uterus, or in her back—most frequently in her back, which latter part, in her agony she often cried out, "must break in two."

Various opiates, amongst others from half a drachm to forty minims of chlorodyne, were given by the mouth; but they all appeared either to disagree with the stomach, or give little or no relief to her suffering, so that I was obliged to resort to the hypodermic administration of morphia, which was first commenced by injecting one-third of a grain of morphia under the skin of the abdomen, over the seat of the uterine tumour. In exactly three minutes (as timed by a watch) from the time of injecting, she was quite free from pain, and certainly presented a sudden contrast to her former state of agony. She was left in a perfectly quiescent state, and breathing tranquilly. She stated that the injection first made her feel very "lightheaded," and then acted like a charm in sending her to sleep. The first injection kept her free from pain for twenty-four hours, after which she had a recurrence in every way as severe as before, but this time the seat of pain was entirely in her back, in the lower dorsal and lumbar regions. The same quantity (one-third of a grain) was injected into the thick cellular tissue of the back, under the skin of the lower dorsal region. The relief was equally significant as in the former instance, and was afforded in four minutes from the time of injection. The pain, principally in the back, recurred almost daily for the space of a fortnight, when she became completely exhausted and worn out by it, and sank, as it were from sheer exhaustion. The quantity of morphia injected was obliged to be increased towards the latter part of her existence, as the system gradually became used to it, but never more than one grain was injected at a time, and this dose had to be repeated occasionally as often as twice in the twenty-four hours. No injurious local effects were ever produced by the injecting process.

* Read before the Lincoln Medical Society, January 1868.

REMOVAL OF FOREIGN BODY FROM THE LEFT BRONCHUS.

By JOHN WALTERS, M.B., Reigate.

T. W., aged 40, tailor, twelve years ago had tracheotomy performed for disease of the larynx and glottis, following fever. He had worn the same tube ever since, not being able to do without it. The voice had been distinct, but husky.

On March 22nd, 1867, at 8.30 A.M., while he was attempting to remove the tube for cleansing, the rim came off, and he could not get hold of the cylinder. When I saw him shortly afterwards, I could see the tube at the bottom of the wound. I went down stairs to get instruments from the carriage, and to ask Dr. Holman to assist me. When I returned, the tube had slipped down the trachea, and could only be felt by a bent probe at a distance of five inches from the external opening, to the right. Respiration was imperfect over the right side of the chest; perfect on the left. Attacks of dyspnoea were at times moderately severe, and cough very troublesome. Inversion was tried, with smart blows on the back and front, without altering the position of the tube. At 3 P.M., having obtained a pair of long curved forceps, we resolved to make further attempts to remove the tube. A stilette, bent at the end, was first introduced, and caught something which seemed like the tube; but, on making traction, smart bright hæmorrhage came rather freely from the tracheal wound, so the hook was disengaged and withdrawn. Chloroform was then administered over the mouth and trachea; it excited him much at first, but he soon went under its influence. The long forceps was then introduced; and the tube was felt, but it could not be caught. Bloody mucus came freely from the mouth and wound, and he retched freely. The respiration became so much impaired, that the chloroform had to be given up, and time given him to eject the accumulation in his air-passages, this condition being somewhat perilous. When he had sufficiently recovered, I made a fresh examination, and found the tube rather more than five inches down and to the left. After one or two failures, the tube was seized with the long forceps, and drawn to the wound, where it stuck; and, while Dr. Holman was enlarging the opening to admit its removal, it slipped back. I managed, however, to catch it again immediately—fortunately, this time in the long axis; and it was readily removed, and found to be filled with mucous membrane. After the removal, he complained merely of being wearied, and of some soreness of the chest.

He did not suffer pain during the attempts at removal—merely distress from obstruction to the breathing. His recovery was very rapid, and in a few days he was able to resume his work.

ON OPERATING FOR STRANGULATED HERNIA.

By G. OLIVER, M.B.LOND., M.R.C.S., etc.

I OPERATED on two cases of strangulated hernia during the past year. The fibrous structures which caused strangulation were dilated by a simple manoeuvre of the tip of the forefinger, after which proceeding the bowel was easily reduced. The hernia-knife was not used. The operation was so simple, and yet so successful, that I am induced to record it.

My patients were both females; and in each case a single reducible femoral hernia had existed for several months before it became strangulated. In one case, symptoms of strangulation existed twenty-four hours, in the other case four days, before the operation was performed. In the latter case, there was persistent and profusely stercoraceous vomiting.

The structures covering the seat of strangulation were incised in the usual way. A wound of about two inches in length was produced, which passed in a direction outwards and slightly upwards across Poupart's ligament. The forefinger exploring the wound felt some firm fibres encircling the neck of the tumour. The knife was dispensed with, and the tip of the forefinger was insinuated under the constricting fibres; and these, as well as firm tendinous bands at the crural ring, were forcibly stretched by firm pressure of the finger in a direction away from the bowel; this, now free from constriction, was easily returned to the abdomen by gentle pressure. The wound, closed by metallic sutures, was healed in a few days. Both patients recovered uninterruptedly.

Since the above cases occurred to me, I have learnt that Baron Seutin successfully dilated the stricture of strangulated hernia by insinuating the tip of the forefinger into it, and by stretching it forcibly.