many of the nuclear fibres and spindle-shaped cells supposed to be illustrations of the cellular development of fibrous tissue.

5. When the disease is not of constitutional origin, there will be found evidence either of the pursuit of some occupation exposing the patient to the inhalation of irritating particles (Peacock; Greenhow), or of the existence of tubercular phthisis, long-continued bronchitis, or of some former attack of acute disease, such as fibrinous pleurisy, pericarditis, or what the author has called the corpuscular form of pneumonia.

6. The disease commonly affects one lung, and more commonly the left than the right.

7. If the disease affect both lungs, it is as a rule, either the effect of mechanical irritation, or it is accompanied and has probably been caused by the presence of tubercles.

In rare cases, the double affection may be caused by rheumatism or syphilis (Wilks).

8. In Fibroid Phthisis there is always contraction of the chest-wall. The percussion dullness is harder, higher-pitched, and more uniformly continuous than in any other lung-disease. Resistance of the thoracic parietes is greatly increased. The intercostal spaces are depressed. Sometimes the dullness is tubular. Vocal fremitus is at one time increased, at another greatly diminished. Over the fibroid lung one hears blowing breath-sounds often without audible prolongation of expiration; occasionally coarse dry and moist râles, superficial creaking, and diffuse bronchophony. Over cheesy deposits of any extent the breath-sounds are sharply tubular, the expiration prolonged, and the vocal resonance bronchophonic, sniffling, and circumscribed.

9. If the apex of the diseased lung be involved in the solidification, there is reason to suspect the existence of tubercles; if there be moist crackling in the supraspinous fossa, the suspicion amounts to probability; and if with the slightest dullness over the summit of the opposite lung there be any moist crepitation, doubt is practically no longer possible.

10. When the lung below the solidification is healthy, expiration is peculiarly prolonged and puffy.

11. When cavities exist, they commonly occupy the mammary region: occasionally they are found in the base, and with extreme rarity in the summit of the lung. (Sutton's cases appear to have been complicated with tubercles.)

12. When about the middle of the lung several cavities lie near together in a horizontal or diagonal line, and are bounded by solidified lung, they are in all probability due to dilated bronchi. Cavities resulting from the breaking up of cheesy deposits are commonly isolated, irregularly placed, larger, and yield on auscultation unmistakable cavernous or amphoric breathing and pectoriloquy. The only certain evidence, however, of the existence of cheesy cavities is the presence of areolæ of elastic tissue in the sputum.

13. In the neighbourhood of the solidified lung it is not uncommon to find patches and tracts of lung in a condition of extreme vesicular emphysema; and the atrophic changes constituting it appear in many instances to be due to plugging, or some other obliteration of branches of the pulmonary artery. It is therefore a statical as opposed to a dynamical emphysema.

14. When the left lung is affected, the heart is usually displaced upwards, and a little outwards. When the right lung is affected, the heart is drawn chiefly outwards, and a little upwards. In both cases, a low-pitched systolic bruit is commonly heard over the pulmonary artery.

15. The cough is paroxysmal, and ordinarily induces vomiting. The expectoration varies; it is usually yellowish, greenish, or ashen grey, studded with pigment streaked with blood; sometimes foetid, and ejected with difficulty after several ineffectual fits of exhaustive coughing.

16. The general symptoms, viewed collectively, are strikingly different from those of tubercular phthisis. The skin is, local sweats excepted, dry and inactive; there is no continuous elevation of temperature, no evening fever, seldom any hectic flushing. Till the disease is far advanced, the breathing is quiet, and the pulse below 84.

17. Slight œdema of the lower extremities is very common in the course of this disease, and almost always present at its close.

18. Patients who have long suffered from Fibroid Phthisis, whether of local or constitutional origin, become at last pallid, waxy-looking, and cachectic.

19. The disease is commonly slow in its progress, and when it complicates tubercles, it retards disintegration and greatly prolongs life. This may explain some of the cases of unusually prolonged phthisis in spirit-drinkers.

20. The frequent occurrence of even copious hæmoptysis, and the setting in of diarrhœa in the course of this disease, may justify a suspicion of the development of tubercles, but are not conclusive evidences of their presence. Hæmoptysis, diarrhœa, and ulceration of the bowels often occur during life without a solitary tubercle being found after death.

FATAL CASES OF POISONING BY THE ABSORPTION OF CARBOLIC ACID.

By E. S. MACHIN, Esq., Erdington, Birmingham.

ON Wednesday evening, February 5th, shortly before eight o'clock, I was summoned to the sick ward of the Aston Union Workhouse by a messenger, who stated that three women had been dressed for itch with carbolic acid, in mistake for sulphur-lotion. I at once proceeded to the workhouse; and, on arriving in the itch-ward, I perceived a very powerful odour of carbolic acid, and the three women lying in a state of great prostration, with spasmodic breathing, and being totally insensible to all around. The patients were, Mary Pritchett, aged 60; Mary Ann Pritchett, her daughter, aged 23; and Mary Baker, aged 68.

On inquiry, I elicited that the acid, after being warmed, had been uniformly applied to the entire surface of the bodies by means of a sponge; that, within a few moments of its application, they had each complained of smarting pain and headache; and that they then became giddy and rapidly insensible. As near as I can ascertain, I saw them within twenty-five minutes of the commencement of the "dressing". I at once gave instructions for removing the acid from the skin by washing with tepid water and soft soap; the water being continually changed as it became impregnated with the acid. I endeavoured to administer some brandy with ammonia and sulphuric ether, and some of it was swallowed. The surface of the body felt harsh and dry; but there did not appear more redness than that produced by the washing. I then had the patients carried down to another ward, as the room was full of the fumes of the acid. The headache and giddiness complained of by the patients before they became insensible were felt by all present, especially by my friend Mr. Elkington, whose attendance I had requested, and who rendered me valuable assistance for many hours.

The elder Pritchett did not rally, but continued unconscious till 11.30, the breathing becoming more laboured and slow. There appeared to be no convulsive action, except an occasional spasm of the diaphragm. The pupils were natural, and answered to the light.

The younger Pritchett, after lying insensible for nearly five hours, gradually became conscious, and an attempt at vomiting was set up. An emetic of mustard was administered; but very little of the contents of the stomach was evacuated, although the retching was considerable. A very strong odour of the carbolic acid was perceived with each expiration. Some strong coffee was given, and she appeared much better. The respiration, although losing its spasmodic character, continued quick and irregular; and pain in the head and throat was complained of. Cold milk with a little brandy was given. In the course of the morning, an epileptiform convulsion came on. Leeches were applied to the temples, and she soon again became conscious; she said the pain in the head was better, that she felt better, but that the pain in the throat was unrelieved. An alkaline mixture with chlorate of potash was prescribed, and afterwards chlorine in solution; but symptoms of congestion of the lungs came on, and she gradually sank, being conscious to the last. Death occurred on the 7th ult., about 10 A.M.—i. e., about forty hours after the application of the acid. This girl was reported to be an epileptic; but she had not had any fit since she had been in the workhouse, a period of about seven months. The attendant who applied the acid said that she had a convulsion directly the "dressing" took place.

The third patient (Baker), who is recovering, states that she was "dressed" first; and that she stood away from the fire during the application, whereas the two Pritchetts stood close to it. She was carried down stairs while unconscious, and rallied in about four hours. She informs me that, immediately the dressing was used, she felt giddy and as if she were drunk; and her head felt as if tightly fastened round with a band. She remembered no more till she found herself in another ward, and the master of the workhouse giving her restoratives. She complained of great smarting over the surface. There was no vomiting in this case, and the bowels have acted naturally; the pulse is regular, but rather feeble (80). The skin appeared at the time harsh, dry, and puckered; there is not (nor was there in any of the cases) any vesication. The skin has been thrown off in small scaly patches; and to-day (February 26th) the woman appears well, being able to get up and take exercise.

A *post mortem* examination of the bodies was not obtained. Externally, there appeared to be small patches of sloughy skin dotted about the bodies.

The carbolic acid used was Calvert's carbolic acid for disinfecting purposes, and was in appearance dark and oily—I presume, an impure acid. As far as I can ascertain, I believe that about six ounces were used for "dressing" the three cases.