whether from disease or after accident; nature of the operation; removal of limb, or in joint-diseases only of the heads of the bones, when the limb is removed; whether a circular incision is made through the integuments, and the edges of the wound brought together with strapping; or whether the flap-operation and the skin brought together by sutures, whether the operation heals by the 'first intention' or by granulation; number of arteries or veins taken up; when removed; stump or wound closed." This direction respecting "arteries or veins" is given twice over. In case of hæmorrhage or phlebitis, such a note might give rise to unfair and injurious comment. The whole is enough to turn the head of any nurse.

If pupil-nurses are to obey these instructions, they must have the foremost places at operations. They actually have, as I am told, at another hospital. Is the nurse qualifying herself to operate? Whenever public opinion demands that lithotomy and other great surgical operations shall be performed by women, instead of surgeons, it will be proper, and therefore decent; but not until then will I endure a group of girls peeping over my shoulders.

As educated women are clearly contemplated, how will the scheme work? Are they to be lady doctors, under another name? or will they criticise, and control, and counteract without responsibility? Will the medical man become a mere ceremony; a convenience, hired reluctantly, to take the blame, if any; just as the regular practitioner is sent for, when a case grows too perilous for the reputation of the quack? Rely upon it, nothing of the kind is thought of; but to that it will come.

Gentlemen, do not infer that I am repressive of improvement, but the painfully menial services, the direct personal assistance, which it is a nurse's bounden duty to render, without flinching, to the helpless and infirm, would be most unseemly if undertaken by a young gentlewoman, fresh from the tutelage of the schoolroom and the nursery, and reared in the most scrupulous abstraction from whatever is equivocal or indelicate.

My own idea of society is, a great collection of all sorts of people, each one contributing his own peculiar quality for the common weal. If the intelligent or the refined, the laborious or the humble, be absent, society is incomplete; and if any section affect the actual business of another, it will furnish but sorry substitutes. This does not forbid an earnest striving to excel in one's own proper duties—without which indeed all would stagnate; but whoever aims at a position other than his own, should be conscious and sure of powers which are irrepressible.

Now there is an old book, and in it a quaint passage which we all learnt when children, though some seem to think it becoming to have forgotten it; so I would suggest to the vain and restless of both sexes and all ranks, that they should be content "to do their duty in that state of life to which it has pleased God to call them".

CASE OF LOW FEVER: DELIRIUM: INCOMPLETE DEMENTIA: CONVULSIONS: DEATH: TUBERCULOSIS.

By C. HANDFIELD JONES, M.B., F.R.S.,
Physician to St. Mary's Hospital.

M. G., female, aged 25, Irish, was admitted into St. Mary's Hospital on December 27th, 1865. She had been ill three weeks; was taken first with pain in the stomach and vomiting. Cough came on soon afterwards, and had become worse; and now it made her vomit whenever it came on. She had no thirst. The skin was hot. The next day it was markedly hot; temperature 104.5°. There were no distinct typhoid spots, or any other eruption. Jan. 3rd, 1866. Sickness had quite ceased. Tongue nearly clean. Pulse 108, weak. There was much restlessness. The countenance was expressive of stupor and uneasiness; the abdomen rather tender; bowels relaxed three times; eyes clear. She had much noisy delirium at night. Jan. 17th. During the last fourteen days, her condition had been quite that of a patient in low fever; she had had more or less stupor, alternating with delirium, mostly nocturnal; she was also very deaf, and did not hear when she was spoken to even with a loud voice. A large bed-sore was forming on the right trochanter. The temperature had kept at 101-102° about the middle of the day. The pulse had been 100 to 110. Jan. 20th. Temperature 99.8°. Pulse 104. She was quite conscious and rational at times, at others very noisy. Jan. 25th. A little blood appeared in the stools to-day. Jan. 30th. She continued extremely deaf; and seemed but half conscious. By March 1st, the bed-sore was healed, but she looked very thin and pallid and semidemented. Her deafness was extreme some times; at others she was able to hear better. March 7th. She looked very sallow and earthy complexioned, and quite as if demented; the intellect was evidently much impaired. There was harsh

but pretty full breathing in both fronts. March 16th. She was much better, getting hungry, much more rational, and less deaf. March 19th. She was doing well yesterday; but passed a very bad night, and was very noisy and delirious; she refused food to-day, unless it was forced upon her. Pulse very weak. Skin cool. March 23rd. She was improving. She took her food well, and enjoyed oatmeal porridge. The sister observed that she always heard better in mornings; after midday she became more deaf. March 26th. She had a fit to-day at noon, and was convulsed much for ten minutes. A few days after this, she became highly delirious at night and screamed violently; her cough was also very bad. Asthenia increased, and she sank in the night of April 3rd, having continued to be noisy almost to the last.

The treatment consisted of nitric acid during the fever period; subsequently she had cod-liver oil and sedatives and peppine wine. Brandy was administered in moderate quantity, and as much nourishment as she could be made to take. About a fortnight before death, her weight was only 66 lbs.

At the autopsy, there were several purpuric patches on the surface. The brain weighed forty-three ounces. It appeared healthy everywhere; was very pale, and rather wet. The heart weighed only six ounces. Both lungs were full of tubercle, which was undergoing softening in numerous foci. There was some effused blood on the surface of the left lung and in the pleural cavity. The liver and kidneys were normal. The small intestine was carefully examined; but no trace was found of ulceration, except a small nearly healed cavity or depression close to the ilio-cæcal valve. Peyer's patches were very little developed, less almost than normal; there was no trace of any morbid alteration in them. In the large intestine, several small ulcers were found not quite cicatrised; towards the lower end, numerous solitary follicles were seen with the central blackish spot, which gives to them, when aggregated in patches, the shaven-beard appearance.

CLINICAL REMARKS. The duration of disease in this case seems to have been about four months; the earlier half or two-thirds of this period presenting the phenomena of low fever, and the remainder those of an imperfect and unsuccessful attempt at convalescence. During the progress of the case, I entertained no doubt that it was one of typhoid fever; nor was it until I came to think over the results of the autopsy, that any doubt arose. The question is, Was the malady primarily and essentially acute tuberculosis of the lungs, masking itself, to a certain extent, under the guise of low fever; or was truly an attack of typhoid which precipitated a tuberculous diathesis into rapid evolution? The first view is favoured by the existence of pretty severe cough throughout, by the existence of bronchial rûle in both backs on admission, by the frequent vomiting, the absence of eruption, the advanced stage of the disease in the lungs, and the absence of any marked lesion of Peyer's patches. On the other hand, the general aspect of the patient, the course of the disease, the high temperature (104.5°) on the twenty-second day, and its fall to 99.8° by the forty-fifth day, the formation and healing of a bed-sore, the deafness, and the mental disorder, are phenomena much more accordant with the second. I have little doubt that the latter is the correct interpretation. But, if it be, I cannot avoid remarking that the case tends to increase the belief I have for some length of time entertained, that other forms of fever exist beside the familiar ones we are agreed to recognise. The specific lesion of typhoid did not exist in this instance, at least not in the lower part of the bowel; and, though I know it is said sometimes to occur only in the upper part, I cannot but think it highly improbable that such was the case here. The ulcers in the mucous membrane of the large intestine were much more like the results of fever than of tuberculosis. Their existence affords a pretty good proof that the freedom of Peyer's patches from morbid change, was not the result of their having had time to recover their normal state. With regard to the delirium and deafness, which were such prominent symptoms, the results of the *post mortem* examination go to show that they were more dependent on defective supply of blood than on atrophy of nerve-tissue. I had fully expected to find the brain very much atrophied; and I was not a little surprised to find it not more than one or two ounces below the normal figure. On the contrary, there can be little question that the heart was unduly small—in fact, atrophied; just as it was plain that the total amount of blood was. Such atrophy of the heart is well known to occur in various wasting disorders, as cancer and diabetes. While thus recognising defective supply of blood as one very important factor of the above disorders, I think it is also certain that it was not the only, and perhaps not the most, influential. I have certainly witnessed cases of complete, or almost complete, pulselessness, when the intravascular pressure must have been reduced to a minimum, where the functions of the nervous centres were less defective than they were in this fever-wasted sufferer. I cannot doubt that the action, the functional energy, of the nerve-cells was greatly impaired by the deleterious influence of the fever miasm. The connection

of the delirium, the incomplete dementia, and the epileptic convulsion, with impaired nutrition of the brain, is most evident. No doubt can exist that, could the digestion and assimilation have been improved, and the blood brought into a proper state as regards its quantity and quality, all the nerve-disorders would have ceased. This may be set down as a typical case of cerebral malady from enfeeblement, and may be instructively contrasted with one recently recorded, where venesection was beneficial. The varying quality of the nerve-disorders may be remarked, the condition of stupor and fatuity alternated with delirious raving, yet both essentially depended on the same conditions. Just so, in the nerves, we find hyperaesthesia and neuralgia, spasm and paralysis, often similarly associated. The improved condition of hearing in the morning, and its subsequent deterioration later in the day, is quite in accordance with what is observed in many instances of functional paresis. Purpuric extravasation in the skin could evidently have depended in this case only on an altered state of vessels, and not on any increase in intravascular pressure. In my experience, it is far most frequently connected with asthenia and defective nervous power. The last case in which I witnessed it was a weakly man who had recently had a good deal of delirium during rheumatic fever.

REPORTS OF HOSPITAL PRACTICE:

METROPOLITAN AND PROVINCIAL.

ST. BARTHOLOMEW'S HOSPITAL.

CASE OF PELVIC CELLULITIS.

(Under the care of Dr. GREENHALGH.)

THE chief point of interest about this case is the pointing of the abscess below, instead of above, Poupart's ligament—a most unusual circumstance, which has not, we believe, been recorded before. We are indebted to Mr. H. Jarvis for the following notes.

Jane S., aged 23, was admitted on Feb. 4th, complaining of loss of power in the left leg, and of a swelling at the upper part of the thigh; also of being very low and weak. She had had two children, the first two years and a half ago. She had a good labour and quick recovery. The second confinement was on the 4th of November last. She had a good labour; no bad results from it. She was confined at the Lying-in Hospital, St. Luke's; and on the tenth day afterwards left the hospital for her own house. The third day after being home, she was suddenly seized with sharp cutting pains all over the abdomen, quickly followed by cold shivers, which occurred about once a day for four or five days afterwards; and night-sweats, sufficient to make the night-dress quite damp; these continued a fortnight. The pains were relieved by hot linseed and bran-poultices; but they continued more or less severe up to the time of her admission here. Her appetite had been very bad since the commencement of the attack; she ate scarcely anything. On Saturday, February 2nd, she discovered a lump at the upper part of the thigh; and now she found herself suddenly relieved from the abdominal pain, from which she has not since suffered.

On her admission, a swelling was discovered just bulging above Poupart's ligament, and more distinct below it. The swelling bulged also into the upper part and left wall of the vagina. Soon after her admission, the skin was discovered to be red; and fluctuation could be felt about an inch and a half below Poupart's ligament. On the seventh day after becoming an in-patient, the swelling was punctured half an inch below Poupart's ligament, and half an inch to the outer side of the anterior crural nerve, giving vent to about ten ounces of pus. The thigh had been flexed to about an angle of 60° one month before the abscess was opened. Two days after this, she was again enabled to extend it. From this time, she rapidly improved in health; her pulse becoming stronger and firmer; and she gained flesh. Ten days after the opening of the abscess, the upper part of her thigh was observed to be inflamed and painful; pus formed; and the opening was enlarged. The inflammation subsided a few days only after it was first noticed.

SUPPURATION OF KNEE-JOINT: ARTICULATION LAID FREELY OPEN: RECOVERY.

(Under the care of Mr. HOLMES COOTE.)

Jane Lysaght, aged 14, was admitted April 1st, 1867, with extensive suppuration around, but not involving, the knee-joint, following the prick of a pin eleven days previously. She had a somewhat anxious expression; rather rapid pulse; and furred tongue. She suffered a great deal of pain in the knee. At the time she pricked her knee, she did not take much notice of it, and continued her household duties as usual. From her description, the pin seems to have entered just below the joint. She was ordered nourishing diet, stimulants, bark, and am-

monia. A free incision was made, and a large quantity of pus evacuated, with great relief to her symptoms. In spite of this, the suppuration extended, and was followed up by other incisions.

Five days after admission, there was noticed some slight effusion into the joint, with an aggravation of pain. This effusion gradually increased until the fourteenth day after admission, when the knee was very considerably swollen. There was also great swelling and tenderness extending half-way up the thigh. She was at this time in a very low hectic condition, and amputation was contemplated, when Mr. Coote, as a last resource, made a large incision four inches in length, freely opening up the joint on its outer side. Five ounces of pus were evacuated. She did not get much better for a few days; but, after that, the suppuration began to decrease, her general condition became better, and the appearance of the knee much quieter. From this time, she gradually improved, and the incisions healed up.

She now (June 26th) has the limb incased in stout leather splints, and walks about with the aid of a stick. There is a little movement in the joint.

The great feature in this case was the laying freely open of the joint; for, if this had not been done, amputation through the thigh must have been the almost inevitable consequence, which would have exposed the girl's life to great danger, to say nothing about the loss of the leg. As it was, the danger to life was not only avoided, but she has recovered with a limb which will, in all probability, be eventually little inferior in utility to the other.

UNIVERSITY COLLEGE HOSPITAL.

OLD STRICTURE OF THE URETHRA; DISEASE OF THE BLADDER; NEPHRITIC AND PERINEPHRITIC ABSCESES; DEATH; AUTOPSY.

(Under the care of Mr. ERICHSEN.)

THE following case, for the notes of which we are indebted to Mr. Poore, House-Surgeon, is a capital illustration of the danger of neglecting an old stricture, and of the complications which are sure to follow in the course of time, in the shape of disease of the bladder, dilatation of the ureters, and inflammation of the substance of the kidneys themselves.

Joseph C., a farrier, aged 36, was first admitted into the hospital in the beginning of February 1867, on account of an old and very tight stricture of the urethra, the consequence of gonorrhoea contracted eighteen years ago. The stricture was cured by means of Holt's dilator, and the patient discharged in the course of the same month. He continued to present himself at the hospital from time to time to have bougies passed, and one day complained of pain in one of his testicles which had swelled; ultimately an abscess formed in it and burst.

On June 11th, he was again admitted into the hospital, but this time under Dr. Jenner's care, because of pain in his loins, and of his passing urine "as thick as milk." He first felt the pain about three weeks before admission; and stated that it was greatly aggravated by stooping or by pressure. Since its first appearance, he had been often seized with fits of vomiting which occasionally left him for three or four days, but returned again.

After his admission, he passed a great quantity of urine, which was loaded with pus. There were no casts of tubes to be seen, and the urine was of the same quality throughout, not thicker at the beginning or towards the end of micturition. There was a distinct fullness on the left side of the back, extending from the tenth dorsal vertebra to the crest of the ilium, and from the middle line to a hand's breadth outside the spiral column. The patient complained of pain when the part was compressed. The fullness over the right kidney having become more prominent, Mr. Erichsen saw the patient, at Dr. Jenner's request; and, having diagnosed the fullness to be due to a perinephritic abscess, he made a deep incision at the most prominent part and let out a considerable quantity of pus. The patient was then transferred to the surgical wards. After his removal, the abscess continued to discharge freely. Large poultices were kept constantly applied over the whole right side. A drainage-tube was also introduced into the abscess, to the depth of about three inches. The quantity of urine was greatly diminished, but it still contained a considerable amount of pus. The vomiting continued every day, chiefly in the morning, and no improvement was obtained in this regard in spite of all remedies. The patient gradually became weaker and weaker, and sank at last on July 3rd.

A post mortem examination of the body was made on the following day. The bladder was found enormously distended, thickened, and fasciculated. Both ureters were enlarged and thickened. The left kidney was small and nodulated, and, on section, was found to contain a number of small abscesses, all of which communicated with the pelvis of the organ. The right kidney was rather large, of a pale colour, and also full of abscesses; and it had contracted adhesions to the right lobe