LECTURE

ON THE

CHARACTERS OF THE ARTERIAL PULSE,

IN THEIR RELATION TO THE MODE AND DURATION
OF THE CONTRACTION OF THE HEART IN
HEALTH AND DISEASE.*Delivered at the Royal College of Physicians, London.*

BY

J. BURDON SANDERSON, M.D., F.R.S.,

Physician to the Hospital for Consumption, and Assistant-Physician to
Middlesex Hospital.

PART III.

THE striking peculiarities of form which the sphygmographic tracing exhibits in cases of aortic regurgitation, have afforded to some writers a ground for regarding the sphygmograph as an instrument of diagnosis in heart-disease. The truth is, that all such cases can be just as satisfactorily made out by the ordinary methods of examination. In these two patients,* for example, there was no manner of doubt as to the nature of the lesion. Indeed, the character of the pulse was so obvious that, with the finger on the wrist, one might have made a diagnosis with one's eyes shut.

In the more numerous instances in which the same peculiarities present themselves in a less marked degree, the presumption they afford in favour of aortic regurgitation falls far short of evidence of its existence.

Fig. 13.—J. S., aged 28. Mitral valvular disease.

Fig. 14.—J. B., aged 25. Aortic valvular disease. The two forms are essentially the same.

To illustrate this, I refer to the two tracings, figs. 13 and 14. One (fig. 13) is the pulse of a man, aged 28, who was at the time recovering from an attack of paralysis. This illness had come on somewhat gradually. When first seen, he had motor paralysis of the right arm and right side of the face, which had come on a few days before somewhat gradually. With this exception, and that of slight metæsthesia of the affected arm, there was no disorder referable to the nervous system. The paralysis rapidly diminished, and the patient regained health and appetite. The tracing was taken three weeks after the seizure, when, on examination of the chest, I found that there was a bellows murmur lasting during the whole systolic period at the apex, but no indication of aortic regurgitation. Let us compare with this the pulse of another patient (fig. 14), in whom the physical signs were so exactly similar to those already detailed with reference to fig. 12, that I need not occupy space in describing them. It is, in fact, so plain a case of aortic regurgitation, that the merest tyro in auscultation could not mistake it. The comparison of figs. 13 and 14 illustrates how completely the organic is simulated by the functional disorder of the heart.

To the question, What is the nature of this functional disorder? I venture to hope that the principles laid down may, in some measure, afford an answer. The condition of the heart which simulates aortic regurgitation, is that in which its contraction is smart, or even violent, but soon over. The bounding and collapsing pulse may, in fact, arise from three causes, all of which must be borne in mind before a conclusion can be come to as to its significance. It clearly cannot exist independently of a relaxed state of the arteries; but this may be produced either by the reflux of blood from the aorta into the heart during the relaxation of the ventricle; by the preternatural velocity with which the blood finds its way through the relaxed capillaries to the veins; or, lastly, it may happen because the heart itself, although contracting sharply, does not contract effectually. To illustrate this, I pass to the next examples, those of functional irregularity of the heart's action. (Figs. 15 and 16.)

It is well known, that persons of gouty constitution are liable to be affected with irregularity of pulse quite independently of valvular lesion. In this affection, the irregularity is one of rhythm. During the period of inspiration, when, as I have elsewhere shewn, the frequency of the contractions of the heart is increased, the pulse becomes dicrotic, its form being entirely different from that which it assumes during the

respiratory pause, when it is relatively retarded. During the respiratory pause, on the other hand, the contractions of the heart are less frequent; the diastolic intervals are longer, so that the heart

Fig. 15.—T. R., aged 57. Irregular pulse of gout. In this patient, during the attack, the pulse becomes regular, as seen in the lower line. At other times, the largest sweeps of the lever occur during the respiratory pause; the smallest, during and after inspiration.

has time to fill completely before it contracts. Hence, the quantity of blood delivered into the aorta is much larger in proportion to the quantity which can be transmitted by the capillaries. The duration of the ventricular systole is greater, and the arteries remain a much longer time distended. The pulse is no longer dicrotic.

Thus, in one and the same individual, you have dicrotism during inspiration—absence of dicrotism during the respiratory pause; the only

Fig. 16.—E. H., aged 58. Irregular pulse of mitral valvular disease.

difference in the state of the circulation being that, in the one case, the diastolic pause is shortened, and, consequently, the ventricle contracts upon an insufficient supply of blood; whereas, in the other, its expansion is complete and its systole effectual.

Let us refer to another case. Fig. 16 shews a pulse often seen in rheumatic valvular disease, with large mitral regurgitation. The heart acts very irregularly. At times, the ventricle contracts effectually; but, at other times, the systolic expansion of the arteries is imperfect and abbreviated, in consequence of which the mean arterial tension declines. Whenever this is the case, the pulse assumes a form which is as distinctly dicrotic as that of typhus; so much so, indeed, that its double character can be readily recognised by the finger.

I have now said all that I have time to say of those forms of pulse which exhibit arterial atony. We have still to consider those which represent the opposite condition, that of excessive resistance.

Increased arterial resistance may be connected with a variety of causes. Thus, it may arise either from a contracted state of the capillaries, from diminished elasticity of the arteries, or from narrowing of the aortic valve. Figs. 17, 18, and 19, serve to illustrate these points. An example of the most simple case of resistance to the contraction of the left ventricle—viz., that which occurs in aortic stenosis—is given in fig. 17. This is the pulse of a patient who died a few days ago in the

Fig. 17.—G. M., aged 40. Pulse of aortic obstruction, with hypertrophy of the left ventricle.

hospital. The heart, with the adherent pericardium, weighs thirty ounces. The left ventricle is both thickened and dilated, and the aortic valve so deformed and beset with vegetations, that the orifice would scarcely admit the tip of the index finger. The other valves are healthy. All the other morbid appearances found were distinctly referable to the cardiac lesion as their cause. Thus, the kidneys, although enlarged and congested, are healthy. The tracing scarcely needs explanation; its characters are seen to be just the opposite of those we have been hitherto considering. The second event, which, in the pulse of aortic regurgita-

* Vide *supra*, p. 40.

tion, and indeed in all atonic pulses, is either entirely suppressed or difficult to distinguish, is here extremely well marked; and the form of that part of the pulse-curve which represents it shows that the systolic expansion of the artery is of nearly equal duration with the diastolic interval. (Compare with figs. 3 and 4.)

The patient whose pulse is represented in fig. 18 was complaining, at the time when the observation was made, of dyspnoea on the slightest exertion and precordial pain. He was subject to nocturnal paroxysms of cardiac distress, which, he stated, always came on with pain at the scrobiculus, followed by palpitation and nausea, with violent respiratory efforts. On examining his chest, I found that the cardiac dullness extended from the sternum to the mammary line, and that the precordial impulse was difficult and expansive. The systolic sound was prolonged, but no abnormal murmur could be made out. He died several months afterwards, when it was found that the heart weighed twenty ounces, and that the left ventricle was both hypertrophied and dilated, without valvular disease.

Fig. 18.—W. S., aged 59. Hypertrophy of the left ventricle without valvular disease.

Fig. 19.—R. B., aged 59. The same.

Fig. 19 is the pulse of a patient, aged 58, whose physical signs were very similar. The cardiac dullness was not, however, so extensive, nor were the symptoms referable to the heart-affection so distressing as in the last case. Both of these patients had bounding, visible pulses, the arteries knocking against the finger like a hammer; and in both, after death, it was found that the kidneys were much contracted, that their capsules were thickened and adherent, that their surfaces were coarsely granular and studded with numerous cysts, and their cortices atrophied. The tracings, therefore, represent that condition of the heart which accompanies advanced renal degeneration, and appears to be its inevitable result. The differences between the two are not important. In the one case (fig. 18), the evidence of diminished elasticity is less marked than in the other.

I regret that time will not permit me to give more numerous examples. Those already placed before you will, I trust, serve to illustrate the principle on which the discrimination of increased arterial resistances rests; viz., the exaggeration of the systolic expansion, or second event. The state of the circulatory apparatus which these conditions imply, although it cannot be said to constitute a disease, is something more than a mere symptom. So strongly am I convinced of its value as an element of diseased action, that I regard its detection and estimation as the most important purpose to which the sphygmograph can be applied. If this condition could be detected only in well pronounced cases of hypertrophy of the heart when the precordial dullness is enlarged, and the patient perhaps is already suffering from well marked signs of disordered circulation, it would be of little value. But, if we could establish as a fact what appears more and more probable, that increased arterial resistance may not only exist independently of any other sign of organic disease, but may be detected with the aid of the sphygmograph, the practical value of the result would be unquestionable. With reference to it, I would, in conclusion, submit to you a consideration which appears to me to be interesting. Anatomical researches lead us to believe it probable that the earliest beginnings of what may be called degenerative disease consist in such structural alterations of the minutest arteries, as, by rendering them less pervious to the circulating blood, must inevitably lead to increased arterial resistance estimable by the sphygmograph. If I am right in believing that this may be measured by the sphygmograph—if, in other words, we can ascertain, by the examination of the pulse, that the heart is overtaxed long before any change can be detected by auscultation or percussion, it is obvious that we have made a step forward of practical utility.

In conclusion, let me state that it is with these considerations in view that I have elsewhere expressed the opinion, that the sphygmograph is likely to be of greater use as an aid in forming an opinion as to the probable duration of life, than in any other department of medical practice. There are many persons in whom, in the absence of any other trace of ailment, the pulse-curve indicates that the arterial resistance is excessive. The question is, Are such persons sound? I think not; but, as yet, we know too little to speak dogmatically.

ON THE INFLUENCE OF THE ALPINE CLIMATES ON PULMONARY CONSUMPTION.*

By HERMANN WEBER, M.D., F.R.C.P.,

Physician to the German Hospital.

IN Europe itself there were up to quite lately no arrangements for the wintering of delicate persons in elevated situations; but it possesses several localities well adapted to the purpose; and in some of the valleys of the Grisons there are not only villages and inns to live in, but also trustworthy medical men to superintend the invalids. One of these valleys is the Upper Engadin, which is already well known to the profession as a summer residence, but is almost equally well suited for the reception of invalids during winter. Another but much less known valley is the Davos, at an elevation varying from 4,500 to 5,100 feet above the sea-level. It runs about eleven miles in length, parallel to the Engadin, but in the opposite direction, i.e., from north-east to south-west, and is traversed by a stream called the Landwasser. Few of the mountains which border the valley on both sides exceed nine thousand feet in height, and most of them take their rise in a comparatively gentle slope, thus admitting the rays of the sun during the greater part of the day; and nowhere in the upper portion of the valley is that sense of oppression felt which dwellers of open countries often perceive in mountainous countries. The geological substratum of the valley is principally gneiss with hornblende and mica-slate; the crests on the right side of the valley consist of various formations of limestone and dolomite, which descend on some places into the valley itself, as in the neighbourhood of Davos am Platz. There are no glaciers and large snowfields in the neighbourhood, a circumstance which, though it takes away from the grandness of the scenery, increases the value of the locality as a health resort, as being less subject to those icy winds descending from glaciers and extensive snowfields. The principal village is Davos am Platz, about 5,100 feet above the sea, with two good inns situated still on the slope of a hill facing the south. Two excellent medical men reside there, Drs. Spengler and Unger, who devote themselves particularly to pulmonary affections, and keep accurate notes of the progress of the cases under their charge. Dr. Unger, to whom I am indebted for much valuable information, himself had a cavity in his right lung when he first resorted to mountainous climates, and now, after a lapse of four years, is not only free from cough, but is also able to undergo considerable bodily exertion. The population of the valley, especially the male, has the appearance of robust health, with the broad shoulders and wide chests peculiar to mountaineers; and the locality has for many years past enjoyed a good reputation for its influence on scrofulous diseases.† I am not yet provided with accurate meteorological observations, but they are being made by Dr. Spengler. The winter is, of course, cold; but this cold is not excessive, and appears to be rather less than in some lower localities of Switzerland, owing principally to the fact that the valley is protected on the north and north-east. The number of clear days is comparatively great in autumn and winter, in fact greater than in summer; and, owing to the configuration of the locality, the rays of the sun have access during the greater part of the day, and their force in the rarefied air is so great as to allow the invalid on fine winter days to sit in the open air or at the open window.

With regard to the diseases prevalent in the Davos, there are, according to Dr. Spengler's experience extending over more than twelve years, no endemic diseases. Pneumonia is the most frequent of the more serious affections, but it terminates in general favourably on the seventh day without active treatment; the lower lobes are those most usually affected, but occasionally the disease occupies the upper lobes; the termination in consumption seems to be unknown. Chronic bronchitis and emphysema are not rare, but are probably less frequent than is usually believed. Croup is a comparatively frequent disease, especially in spring. Rheumatic affections and diseases of the heart are frequent, and not less so various forms of dyspepsia, and chronic affections of the stomach, especially the round ulcer. The mortality of children is small. Scrofulous diseases originate very rarely in the valley, and those introduced from abroad are favourably influenced by a prolonged residence. Dr. Unger especially mentions the entire absence of tubercular diseases of the lungs among the natives who have never left their valley; and Dr. Spengler perfectly corroborates this statement, and has also mentioned to us cases of men who had become consumptive while working abroad, and have regained their health by a timely return to their native home. Dr. Brügger, it may be remembered, has made the

* Continued from page 42.

† Dr. Meyer Ahrens, "Die Heilquellen und Kuroerte der Schweiz." Zurich, 1869, p. 702.