tion, and indeed in all atonic pulses, is either entirely suppressed or difficult to distinguish, is here extremely well marked; and the form of that part of the pulse-curve which represents it shows that the systolic expansion of the artery is of nearly equal duration with the diastolic interval. (Compare with figs. 3 and 4.)

The patient whose pulse is represented in fig. 18 was complaining, at the time when the observation was made, of dyspnoea on the slightest exertion and precordial pain. He was subject to nocturnal paroxysms of cardiac distress, which, he stated, always came on with pain at the scrobiculus, followed by palpitation and nausea, with violent respiratory efforts. On examining his chest, I found that the cardiac dullness extended from the sternum to the mammary line, and that the precordial impulse was difficult and expansive. The systolic sound was prolonged, but no abnormal murmur could be made out. He died several months afterwards, when it was found that the heart weighed twenty ounces, and that the left ventricle was both hypertrophied and dilated, without valvular disease.

Fig. 18.—W. S., aged 59. Hypertrophy of the left ventricle without valvular disease.

Fig. 19.—R. B., aged 59. The same.

Fig. 19 is the pulse of a patient, aged 58, whose physical signs were very similar. The cardiac dullness was not, however, so extensive, nor were the symptoms referable to the heart-affection so distressing as in the last case. Both of these patients had bounding, visible pulses, the arteries knocking against the finger like a hammer; and in both, after death, it was found that the kidneys were much contracted, that their capsules were thickened and adherent, that their surfaces were coarsely granular and studded with numerous cysts, and their cortices atrophied. The tracings, therefore, represent that condition of the heart which accompanies advanced renal degeneration, and appears to be its inevitable result. The differences between the two are not important. In the one case (fig. 18), the evidence of diminished elasticity is less marked than in the other.

I regret that time will not permit me to give more numerous examples. Those already placed before you will, I trust, serve to illustrate the principle on which the discrimination of increased arterial resistances rests; viz., the exaggeration of the systolic expansion, or second event. The state of the circulatory apparatus which these conditions imply, although it cannot be said to constitute a disease, is something more than a mere symptom. So strongly am I convinced of its value as an element of diseased action, that I regard its detection and estimation as the most important purpose to which the sphygmograph can be applied. If this condition could be detected only in well pronounced cases of hypertrophy of the heart when the precordial dullness is enlarged, and the patient perhaps is already suffering from well marked signs of disordered circulation, it would be of little value. But, if we could establish as a fact what appears more and more probable, that increased arterial resistance may not only exist independently of any other sign of organic disease, but may be detected with the aid of the sphygmograph, the practical value of the result would be unquestionable. With reference to it, I would, in conclusion, submit to you a consideration which appears to me to be interesting. Anatomical researches lead us to believe it probable that the earliest beginnings of what may be called degenerative disease consist in such structural alterations of the minutest arteries, as, by rendering them less pervious to the circulating blood, must inevitably lead to increased arterial resistance estimable by the sphygmograph. If I am right in believing that this may be measured by the sphygmograph—if, in other words, we can ascertain, by the examination of the pulse, that the heart is overtaxed long before any change can be detected by auscultation or percussion, it is obvious that we have made a step forward of practical utility.

In conclusion, let me state that it is with these considerations in view that I have elsewhere expressed the opinion, that the sphygmograph is likely to be of greater use as an aid in forming an opinion as to the probable duration of life, than in any other department of medical practice. There are many persons in whom, in the absence of any other trace of ailment, the pulse-curve indicates that the arterial resistance is excessive. The question is, Are such persons sound? I think not; but, as yet, we know too little to speak dogmatically.

ON THE INFLUENCE OF THE ALPINE CLIMATES ON PULMONARY CONSUMPTION.*

By HERMANN WEBER, M.D., F.R.C.P.,

Physician to the German Hospital.

IN Europe itself there were up to quite lately no arrangements for the wintering of delicate persons in elevated situations; but it possesses several localities well adapted to the purpose; and in some of the valleys of the Grisons there are not only villages and inns to live in, but also trustworthy medical men to superintend the invalids. One of these valleys is the Upper Engadin, which is already well known to the profession as a summer residence, but is almost equally well suited for the reception of invalids during winter. Another but much less known valley is the Davos, at an elevation varying from 4,500 to 5,100 feet above the sea-level. It runs about eleven miles in length, parallel to the Engadin, but in the opposite direction, i.e., from north-east to south-west, and is traversed by a stream called the Landwasser. Few of the mountains which border the valley on both sides exceed nine thousand feet in height, and most of them take their rise in a comparatively gentle slope, thus admitting the rays of the sun during the greater part of the day; and nowhere in the upper portion of the valley is that sense of oppression felt which dwellers of open countries often perceive in mountainous countries. The geological substratum of the valley is principally gneiss with hornblende and mica-slate; the crests on the right side of the valley consist of various formations of limestone and dolomite, which descend on some places into the valley itself, as in the neighbourhood of Davos am Platz. There are no glaciers and large snowfields in the neighbourhood, a circumstance which, though it takes away from the grandness of the scenery, increases the value of the locality as a health resort, as being less subject to those icy winds descending from glaciers and extensive snowfields. The principal village is Davos am Platz, about 5,100 feet above the sea, with two good inns situated still on the slope of a hill facing the south. Two excellent medical men reside there, Drs. Spengler and Unger, who devote themselves particularly to pulmonary affections, and keep accurate notes of the progress of the cases under their charge. Dr. Unger, to whom I am indebted for much valuable information, himself had a cavity in his right lung when he first resorted to mountainous climates, and now, after a lapse of four years, is not only free from cough, but is also able to undergo considerable bodily exertion. The population of the valley, especially the male, has the appearance of robust health, with the broad shoulders and wide chests peculiar to mountaineers; and the locality has for many years past enjoyed a good reputation for its influence on scrofulous diseases.† I am not yet provided with accurate meteorological observations, but they are being made by Dr. Spengler. The winter is, of course, cold; but this cold is not excessive, and appears to be rather less than in some lower localities of Switzerland, owing principally to the fact that the valley is protected on the north and north-east. The number of clear days is comparatively great in autumn and winter, in fact greater than in summer; and, owing to the configuration of the locality, the rays of the sun have access during the greater part of the day, and their force in the rarefied air is so great as to allow the invalid on fine winter days to sit in the open air or at the open window.

With regard to the diseases prevalent in the Davos, there are, according to Dr. Spengler's experience extending over more than twelve years, no endemic diseases. Pneumonia is the most frequent of the more serious affections, but it terminates in general favourably on the seventh day without active treatment; the lower lobes are those most usually affected, but occasionally the disease occupies the upper lobes; the termination in consumption seems to be unknown. Chronic bronchitis and emphysema are not rare, but are probably less frequent than is usually believed. Croup is a comparatively frequent disease, especially in spring. Rheumatic affections and diseases of the heart are frequent, and not less so various forms of dyspepsia, and chronic affections of the stomach, especially the round ulcer. The mortality of children is small. Scrofulous diseases originate very rarely in the valley, and those introduced from abroad are favourably influenced by a prolonged residence. Dr. Unger especially mentions the entire absence of tubercular diseases of the lungs among the natives who have never left their valley; and Dr. Spengler perfectly corroborates this statement, and has also mentioned to us cases of men who had become consumptive while working abroad, and have regained their health by a timely return to their native home. Dr. Brügger, it may be remembered, has made the

* Continued from page 42.

† Dr. Meyer Ahrens, "Die Heilquellen und Kuroerte der Schweiz." Zurich, 1869, p. 702.

same statement with regard to the neighbouring valley of the Upper Engadin; and Dr. Bener of Klosters likewise states that, in the upper part of the Prättigau, immediately adjoining the Davos, although below the level of 4,000 feet, he has not met with consumption; while it has repeatedly occurred to him in the lower part of the same valley.*

The absence of consumption and scrofulous diseases among the inhabitants of the valley who have never left their home, the influence exercised on these diseases, when the natives have acquired them during their stay in other localities, by a timely return to the valley, and the usually favourable termination of pneumonia, are important facts in recommendation of the curative influence of such mountain valleys in scrofulous and consumptive diseases—facts which cannot be urged in the same manner for other localities greatly in favour with the profession and the public. I can, however, not deny that there are also disadvantages connected with a prolonged stay in these mountain regions, especially the want of the habitual society and healthy excitement, and of the accustomed comfort. I do not undervalue these disadvantages, which to some invalids are most serious drawbacks; but, if it could be proved that, in a certain class of cases, the curative influence is much greater than in the usual and more agreeable health-resorts, the profession would sooner or later declare in favour of the elevated regions, and the present disadvantages would thereby gradually be much lessened. I shall, for this purpose, give an outline of those cases which have particularly influenced me in forming a favourable view, and then add a short abstract of the observations made by Drs. Spengler and Unger at the Davos.

CASE I. J. H., aged 24, a clockmaker from the Black Forest, who formerly had enjoyed good health, came to England in the summer of 1848. He had repeated attacks of sore-throat and bronchitis in the same and in the following winters, and became an out-patient of the German Hospital in the summer of 1851, with cough and dyspeptic symptoms. He had slight dulness and impaired respiration in the right clavicular and infraclavicular region; and, with slight variations, gradually lost weight up to September 1852, when he was seized with bronchopneumonia of the lower part of the left lung, slowly creeping upwards, until, in the beginning of November, the dulness reached on the back up to the upper third of the scapula; the crepitant rhonchus having ceased in the lower portion of the dull space, but being persistent in the upper. On the right, the originally diseased side, the dulness extended from the apex to the fourth rib, and was accompanied by a slight degree of bronchophony and large-sized rhonchi. There was great emaciation; pyrexia every evening, followed by perspiration towards the morning. Pulse 110 to 120; and chest-expansion 33 to 34½ inches. In this condition he returned to his home in the Black Forest, about 2,800 feet above the sea-level; where, after some weeks, on milk diet, he gained strength, lost the night-perspiration, and gradually also the cough. In July 1853, when he returned to London, nothing abnormal could be discovered on the left side. The upper part of the right was considerably flattened; but the dulness extended only from the apex to the second intercostal space; the bronchophony was replaced by absence of vesicular murmur and prolonged expiration; and there were no rhonchi. Pulse 82; chest-expansion 34½ to 36½ inches; increase in weight, 21 lbs. He remained well up to August 1854, when he again began to cough; not long after which, while at work, he was seized with violent hæmoptysis, and died on the fifth day.

Post Mortem Examination. The upper lobe of the right lung was adherent; the apex hard, contracted, consisting entirely of slate-coloured, dense, fibrous tissue, almost grating under the knife. The lower part of the same lobe was expanded, as in emphysema. The two other lobes were free from old disease; but several bronchi were plugged up with coagulated blood. The left lung was everywhere adherent. In the centre of the upper lobe was a small fresh cavity filled with blood, with grey soft infiltration in its neighbourhood. In various parts of both lobes were apoplectic nodules; several bronchi were filled with decomposing coagulated blood. The heart was fatty. The liver was likewise slightly fatty. The kidneys were congested.

This case, which evidently was one of Addison's pneumonic phthisis, is of great interest, not only on account of the rapid recovery after the patient's return to his mountain home, but also on account of the relapse and fatal termination after the too early return into unfavourable conditions; and, further, on account of the light thrown upon the nature of the affection by the result of the *post mortem* inspection.

CASE II. J. K., aged 22, a native of Switzerland, formerly healthy, came to London in 1849, and was employed as a waiter in a City dining-room. He began to cough in 1850, and had hæmoptysis in February 1852. Dulness and rhonchi extended from the right apex to

the third rib. There was increase of cough and pyrexia in July. Pneumonic affection of the lower part of the left side was discovered in August. The dulness and crepitant rhonchi gradually ascended to the supraspinal region on the left side (in November); and the older affection of the right side had much extended. The patient was never entirely free from fever. Loss of weight, 30 lbs. in eight months; pulse 90 to 95; respirations 30 to 36; expansion of chest, 34½ to 35½ inches. In this condition he returned home in November to an elevation of about 5,000 feet; and there, on an almost pure milk-diet, he gradually improved, and came back to London in May 1853, having gained 24 lbs. The left side appeared normal; the upper part of the right was sunk in; but there were no rhonchi, and the dulness and defective breathing extended only to the second rib. Pulse 75; respirations 24; expansion of chest, 35 to 37½ inches.

He remained well until March 1855, when he contracted bronchitis, followed by pleuropneumonia of the right side, beginning at the apex, and extending rapidly downwards. In June, the upper part of the left side became likewise affected. Death ensued at the end of this month, from hæmoptysis.

Post Mortem Examination. There were extensive adhesions on both sides. The upper lobe of the right lung was much contracted and puckered; the apex was occupied by several chalky concretions, surrounded by dense, fibrous, slate-coloured, airless tissue. The remainder of the right lung was in a state of soft grey infiltration, with several fresh irregular cavities. The apex of the left lung was in a similar condition; but the greater part of this lung was permeable to air; it contained many hæmorrhagic spots. There were no milary tubercles. The heart was fatty. The kidneys were in the first stage of Bright's disease.

This was again a case of Addison's pneumonic phthisis, occurring under the influence of confined air in a youth accustomed to pure mountain air. Great improvement or cure was produced by a prolonged stay in his native mountain climate—an improvement which continued for some years after his return into unfavourable hygienic conditions. At last, however, a fresh attack of catarrhal pneumonia led to rapid breaking down of tissue, the formation of cavities, and death accelerated by hæmoptysis.

[To be continued.]

CONVULSIONS IN UTERO.

By JAMES WHITEHEAD, M.D., Manchester.

ON March 25th, 1867, Mrs. D., in the ninth month of her second pregnancy, was seized, at 6 P.M., with severe abdominal pain of paroxysmal character, recurring at intervals of three to five minutes. Each attack was attended with an inclination to down-forcing effort, and hence appeared to indicate the near approach of labour.

I saw her two hours after the first seizure, and learnt that the paroxysms had steadily increased in severity and duration from the onset. For two or three minutes after my arrival, the patient was tranquil and half-dozing, the uterine tumefaction presenting its natural roundness and uniformity; but, while in the act of carrying the flat hand over the abdominal surface, a sudden shock of the system took place, followed immediately by the elevation of a tumour of the size of a fist on the left side of the umbilicus, very sensitive under slight pressure, and alternately increasing and diminishing slightly in bulk and tension. The throes were of an irregularly pulsating character; and the tumour, under the gentle grasp of the hand, felt as if caused by a succession of violent spasmodic plunges of a foot of the foetus against this part of the uterine wall. Sometimes I thought I could define the outlines of the foot as if embedded in the oval-shaped tumour. After the commencement of a paroxysm, the plunges, each one being succeeded by a very perceptible quivering motion, were repeated in rapid succession, once every four or five seconds during the first half-minute, then somewhat less frequently during the next half-minute, after which they diminished in force and frequency, and ceased about the end of the second minute, the local tumour at the same time subsiding, and merging itself in the general uniform rotundity of the abdomen, as it had appeared during the preceding interval of quiescence.

The patient, on the subsidence of the paroxysm, was greatly exhausted, drowsy, and disposed to slumber. Her skin was bathed in cold perspiration; the breathing was slow, with occasionally deep sobbing; the circulation was languid; pulse 64, small and feeble.

The lower section of the uterus, now examined for the first time, could be felt above the brim of the pelvis, but not protruding within it. The uterine orifice was not dilated nor dilatable; the cervix, though short, was still tubular and unoccupied. The foetal head could be felt through the uterine walls on each side, but its exact position was not

* I owe this statement of Dr. Bener's experience to a communication from Dr. Unger of Davos am Platz.