Hemorrhage, and to the admirable manner in which the local anæsthetic was employed by Dr. Richardson. In conclusion, a word or two as to after treatment, which, upon broad principles, should depend on the general condition of the patient, and the nature and severity of the symptoms. It is worthy of remark, that although five out of the nine cases lived for periods varying from two to eighteen days, not one died from the effects of abdominal or uterine inflammation, but from causes more or less interfering with, or taxing, the vital powers. This I conceive to be a most important fact, and one worthy of serious consideration. In the majority of the cases a supporting and cautiously stimulating system was carried out from the first with good results; so much so, that I have strong reasons to believe, and those who watched some of the cases with me have expressed similar convictions, that had this plan been carefully and systematically followed out in all the cases recorded, I should have been enabled to place before you not one, but three, successful cases.

CASES ILLUSTRATIVE OF PROGNOSIS IN HEART-DISEASE.

By W. H. BROADBENT, M.D.,

Senior Assistant-Physician to St. Mary's Hospital.

IN the BRITISH MEDICAL JOURNAL for 1866, there was published a paper of mine on Prognosis in Valvular Disease of the Heart; and it is my wish to illustrate some of the points therein advanced by a few cases in which the principles laid down have been applied. The Elements of Prognosis I gave as based upon the following considerations.

1. The organic condition of the heart as ascertained by physical signs.
2. The degree of impairment in its functional efficiency, as manifested by symptoms.
3. The general state of health and soundness of tissues and organs.
4. The presence or absence of conditions tending to aggravate the valvular or structural lesions in the heart, or to precipitate the occurrence of complications.

Each of these general considerations was subdivided into special indications.

1. The organic condition of the heart was shown to depend upon:
 - A. The particular valve affected, and the effect produced by the lesion, constriction, or insufficiency, causing obstruction on the one hand, regurgitation on the other.
 - B. The extent of the damage sustained by the valve, and the degree of mechanical difficulty in the transmission of the blood to which it had given rise.

The murmurs enable us easily to locate and characterise the valvular alteration; and the relative gravity of the different affections I gave, according to Dr. Walshe, as follows: Mitral regurgitation and constriction; aortic regurgitation; pulmonary constriction; aortic constriction; pointing out the special tendencies and dangers of each.

A considerable degree of change in any valve will cause derangement of the circulation, and out of this the danger arises. But the delay may be of two kinds: from an obstacle or impediment to the transit of the blood through the heart, causing it to accumulate in the veins—this I have called "obstruction"; or from deficient *vis a tergo* in the expulsion of the blood from the heart, which primarily causes sluggishness and feebleness of the onward current in the arteries—this I termed "stagnation". The two are always more or less combined in different degrees; but, speaking generally, the tendency of mitral disease is to obstruction of the circulation, and the danger congestion; the tendency of aortic disease is to stagnation, and the danger syncope.

The degree of change which has taken place in the valve is of far greater import than the particular valve affected, and is far more difficult to ascertain. The character of the murmur is no guide; the great indication is the amount of structural change in the walls and cavities of the heart—hypertrophy and dilatation—to which the mechanical difficulty has given rise, and, when cardiac symptoms are present, their nature and severity.

2. The degree of impairment in the functional efficiency of the heart, the result of its organic condition, and aiding in the estimation of this, is shown:
 - A. By the evidence of habitual derangement in the systemic and pulmonary circulation, such as shortness of breath on exertion, a congested condition of the capillaries, dusky face, blue lips, cold extremities, etc.
 - B. By the liability to paroxysms of dyspnoea or attacks of palpitation.
 - C. By the existence of secondary consequences—pulmonary complications or dropsy.

3. The general state of health and soundness includes:
 - A. The condition of the blood and tissues—the existence or not of anæmia and in-nutrition.
 - B. The soundness and functional activity of the important viscera, or the reverse.

4. The conditions tending to affect the state of the valves, or to precipitate or retard evil consequences, are:
 - A. The progressive or stationary character of the disease in the valves.
 - B. The structural health or degeneracy of the walls of the heart and coats of the great vessels, when this can be ascertained.
 - C. The mode of life of the patient.

There can be no case of heart-disease into the prognosis of which all the four great general considerations will not enter, and few in which most of the special indications will not find a place. It would, however, be impossible to illustrate each point by individual cases. Some are at once obvious. One of the points which I wish especially to enforce by the examples I have to bring forward, is the importance, as an element in prognosis, of the character of the pathological process by which the alteration in the valves has been brought about.

The importance of inquiring into the nature of the change in the valve consists in the fact that, according as it is due to one or other of the common affections to which the valves are liable, it will probably be stationary or progressive. If it be traceable to an attack of acute rheumatism, in the absence of further attacks, it may be expected to remain in its existing condition. If it be recognised as due to atheromatous change or other disease of progressive character, this will continue, and will increase the degree of impairment already induced. This consideration has, of course, most weight before serious symptoms have developed themselves. In the one case, they may be postponed indefinitely, unless precipitated by some powerful exciting cause; in the others, they will be evolved sooner or later from the increasing imperfection in the valvular apparatus. It has its value, also, when complications have set in. In the one case, we may have reason to believe that compensatory changes in the heart-walls have for some time balanced the changes in the valves, and that rest and favourable circumstances may restore equilibrium. In the other, we can never have this assurance.

Another point which I wish to illustrate is, that the changes in the walls of the heart, the hypertrophy and dilatation, are indicative of the degree of mechanical difficulty caused by the morbid changes in the valves, and that their prognostic indications are best estimated in this light. Recent authors of great eminence have enunciated the following views: 1. That the symptoms which appear in heart-disease, and the dangers attending it, arise not so much from the state of the valves as from co-existing hypertrophy or dilatation, which are said to increase, and not to counteract, the derangement in the circulation caused by the valvular disease. 2. That this hypertrophy and dilatation are not due to mechanical difficulty occasioned by the imperfect function of the valves, but to inflammation, congestion, innutrition, or other affection of the walls of the heart arising independently of the valvular change.

I hold, on the contrary, that the hypertrophy and dilatation are, as a rule, caused by the mechanical difficulty arising out of the change in the valves, and are, at the same time, the result and the measure of the difficulty. I hold, again, that hypertrophy and dilatation are, for the most part, alike conservative, and tend to counteract the effects of the valvular disease; and that the symptoms, instead of being due to the hypertrophy and dilatation, are due to the valvular obstruction or insufficiency, the effects of which they fail perfectly to neutralise.

According to the view first stated, in a case where a valvular murmur was found to co-exist with alterations in the walls and cavities of the heart, there would be valvular change of uncertain amount, complicated with hypertrophy and dilatation of uncertain origin, and the symptoms which might exist would be due in uncertain proportion to the one or the other. As I should look upon it, the degree of hypertrophy and dilatation would indicate the extent of the valvular disease—being caused by the mechanical difficulty to which this gives rise, and tending to overcome it. In the absence of symptoms, the compensation is effected, and the structural change would furnish an exact measure of the valvular change. Any symptoms which may arise are due primarily to the imperfect function of the valves; and their comparative severity marks the degree in which the hypertrophy and dilatation have failed to overcome the obstruction.

Hypertrophy and dilatation are looked upon as furnishing unfavourable prognostic indications under both views, but for different reasons—on the one hand, as in themselves an evil superimposed upon the lesion of the valve; on the other hand, as marking the great extent of this lesion.

We may be called upon to give an opinion as to the future of a case in which a valvular murmur exists under various circumstances. 1. When the subject of the murmur is apparently in perfect health; as, for instance, when examining for life assurance. 2. When suffering from cardiac symptoms of various kinds and in different degrees, not immediately threatening life. 3. When in some stage of dropsy, or in the crisis of some pulmonary complication.

My illustrations are drawn chiefly from the out-patient practice of St. Mary's Hospital; and the cases, in consequence, belong chiefly to the

second of the three categories. In the press of an out-patient room, a murmur is not likely to be sought for unless attention is called to the heart by symptoms; and, on the other hand, when these have reached a certain degree of severity, the patient is too ill to attend, and must be admitted or he disappears.

CASE I.—*Mitral Disease of Rheumatic Origin, Overwork, and Insufficient Food*.—C. H., a woman, aged 32, looking at least ten years older, came as out-patient to St. Mary's Hospital on May 31st, 1866. Six or seven years before she had had acute rheumatism, and had never been well since. During all the previous winter, she had had a cough and had spat blood. She now complained of cough and shortness of breath on exertion, especially on going upstairs, pain in the loins, giddiness, and great weakness. The legs were œdematous and the abdomen large. Her lips had a peculiar livid colour, as if the blood were highly venous; but the countenance generally did not give a corresponding indication. She had to stand a great deal, and to work hard, having to maintain herself and an invalid husband; she was on her legs from 6 A.M. till midnight. Her diet also was insufficient. The heart was rather large; the apex beat being felt in the fifth space, about an inch to the left of the nipple line. The impulse was not violent; the heart's action regular. A systolic mitral murmur was audible at and near the apex. She took at first compound squill mixture, with twelve minims of tincture of sesquichloride of iron. The cough was improving, but the œdema was increasing. On June 14th, she had larger doses of the tincture of sesquichloride of iron with ether; and on the 25th, I began to order her daily dinners at the St. Mary's Kitchen. She improved greatly, and the œdema quite disappeared when she had been supplied with the nourishing food for a month (July 23rd). I could not continue this indefinitely, and she began to fall back, having still to work hard and being unable to take the rest she needed. August 23rd. The œdema was returning in the legs; she was very sick. Sept. 13th. The legs swelled especially at night after a day's standing. She had much pain in the head, which was worse at night; also pain across the loins. She was extremely weak. The heart fluttered on exertion, and it seemed to her as if her breathing would be stopped. Her aspect was not so good. The pulse was irregular as to force and rhythm, nearly intermitting. Murmur as before. October 1st. The legs were now swollen above the knees. She had cough, with much expectoration and pain in the chest. She was very anxious and nervous. The breathing was short, and she had to stop many times in coming to the hospital. Pulse irregular as to force, and small. The systolic murmur at the apex was louder and heard over a more extensive area upwards and towards the sternum.

This poor woman might live for many years, so far as the heart-disease is concerned. It is stationary, and does not give rise to any very great embarrassment in the circulation. But everything is against her in her mode of life. She is overworked and ill-fed. The blood is poor and the tissues imperfectly nourished; and the anæmia and enfeebled state of the heart-walls are certain to hasten the occurrence of complications. The improvement effected by the good food I was enabled to procure for her was striking, confirming the opinion that, under more favourable conditions, she might suffer little from her heart-disease.

CASE II.—*Mitral and Aortic Disease of Rheumatic Origin: Great Hypertrophy and Dilatation*.—E. P., aged 43, a labourer, came as an out-patient to St. Mary's Hospital on October 2nd, 1865. He had had acute rheumatism eight times; first when very young, last eleven years since, when he was six months in recovering. Ten years ago, he fell from a cart, injuring his back, and was ill for six months in consequence. After this, he worked at his occupation up to the time of attending at the hospital. Two winters ago, he had cough and shortness of breath. He complained of pain in the precordial region, short breath, and extreme weakness, and the feet were swollen. His expression was anxious, and the face was of a deep general red colour, almost amounting to purple. The pulse was very irregular in force and rhythm, and intermittent. On examination, the heart was found to be very large. The impulse was felt over a large space; the apex beat was most evident in the sixth space below and to the left of the normal situation; but it could be felt also much beyond this point for an inch and a half lower down, and as far as the line of the axilla laterally. To the left of the apex a systolic murmur was audible; to the right of it a diastolic murmur which became louder nearer the sternum. At the third right space a systolic and a diastolic murmur were audible; the latter louder and longer. At the right sterno-clavicular articulation the systolic murmur was loudest. There was no second sound in the neck. The pulse did not correspond with the heart, and many systoles failed to reach the wrist. There was here every indication of serious embarrassment in the circulation. Great irregularity in the heart's action, short breath, congested systemic capillaries, commencing dropsy. He presented a remarkable example of dyspnoea become so habitual, that, while it was at

once evident to the observer, he was scarcely conscious of it. Both the aortic and mitral valves were diseased, and the heart was greatly enlarged, showing that very considerable mechanical difficulty existed. Notwithstanding all this, my prognosis as to the existing attack was favourable; and I recommended him to enter the hospital, assuring him that in a few weeks he would be out and at work. The grounds of this opinion were, that the change in the valves was traceable to a rheumatic attack; that consequently they had been for eleven years at least in the same state, and that all this time the disease had not given rise to symptoms of sufficient severity to prevent him from working as a labourer. I concluded that the changes in the walls and cavities of the heart, the hypertrophy and dilatation found to exist, though very great, had been so far compensatory to the changes in the valves as to re-establish a kind of equilibrium. Apprehending no new change in the valves to disturb this, I felt sure that, with rest and good diet, the symptoms would soon disappear. He declined to become an in-patient; and I gave him tincture of sesquichloride of iron, tincture of digitalis, and spirit of chloric ether, of each ten minims, in quinine mixture. He improved, and on October 26th felt so much better that he went to work without asking permission, and continued at it for a week, but was then compelled to desist. He afterwards again improved, and finally ceased to attend on Nov. 23rd in order to resume work.

[To be continued.]

IRREDUCIBLE INGUINAL HERNIA:

STRANGULATION: OPERATION: SLOUGING OF GUT: COMPLETE RECOVERY.*

BY HENRY J. ALFORD, M.B.Lond.,
Surgeon to the Taunton and Somerset Hospital.

ON Tuesday, May 14th, 1867, I was asked by Dr. Bridge of Wellington, to see, with him, Mrs. G., who had been suffering from strangulated hernia since the preceding Friday. Her past history was as follows. She was a married woman, aged about 47, strong and very active. She had been ruptured for many years. Six years ago, Dr. Bridge, with great difficulty, reduced the hernia for her, and ordered her to wear a truss; this she, however, soon left off, and for four years the hernia had been irreducible. On the previous Saturday she was seized with vomiting and great pain in the bowels. On Sunday Dr. Bridge saw her, and applied the taxis, tobacco enema, and other remedies, but to no purpose. About seven o'clock on the Tuesday evening I saw her with Dr. Bridge. I found her in bed; with great anxiety of countenance, constant vomiting, said to be stercoraceous, and severe paroxysmal attacks of abdominal pains. On the right side was a large inguinal hernia, of the size of a child's head. Chloroform was administered, and the taxis applied; but no impression was made on the tumour. I therefore at once operated, and divided a tight stricture at the external ring. On opening the sac, some very offensive serum, having a gangrenous odour, escaped, and the gut was purple and compressed. I found such firm adhesions that it was impossible to return the hernia. A few stitches brought the edges easily together; warm flannels were applied, and thirty drops of Battley's solution given. This was repeated at bedtime. She took the chloroform remarkably well, and recovered soon after the operation, expressing her surprise at its completion. She continued doing very well until Friday, May 17th. She had no sickness, pain, nor tenderness on pressure; no discharge from the wound; and the ruptured parts looked healthy. The warm fomentations and opium were continued, and seemed to afford her great relief. However, on the Friday night she was seized with vomiting and rigors; and when Dr. Bridge saw her he found her in a cold clammy sweat, and almost pulseless.

I was to have seen her on the Sunday, but hearing she was moribund I did not go over. On Monday morning, however, I heard that at ten on Saturday night she had rallied. She had taken brandy, and ammonia and bark; and her pulse recovered, and she became warmer. The ruptured part in patches was black and offensive.

Sunday, May 19th. All pain had ceased; she was much better; vomited a little in the afternoon. Some flatus and fæces also passed *per anum*.

Tuesday, May 21st. I saw her, and found that just before our arrival the tumour had given way at the lower and under part, and a quantity of exceedingly fetid matter had made its escape. Her pulse was good, and she took her food well. Poultices, with permanganate of potash, were ordered to the part. After this she progressed most favourably. For about ten days the fæces escaped through this artificial anus, but after that the passage was thoroughly established *per anum* only.

* Read at the meeting of the West Somerset Branch.