LECTURES

ON THE

PRINCIPLES OF SURGICAL DIAGNOSIS:

ESPECIALLY IN RELATION TO SHOCK AND
VISCERAL LESIONS.*Delivered at the Royal College of Surgeons of England.*

By F. LE GROS CLARK, F.R.C.S.,

Surgeon to St. Thomas's Hospital, and Examiner in Surgery in the University of London.

LECTURE II.—*Concluded.*

Symptoms and Signs.—Antecedent Condition of Patient: his Habits, Temperament, and Occupation.—Exciting Causes of Injury or Disease.—Existing Subjective and Objective Indications.—Signs, Certain, Equivocal, Negative, and Accidental.—Sources of Obscurity and Fallacy.—Functions of Animal and Organic Life.—Intellect, Expression, Colour, Volition, Sensation, Excito-motor Phenomena—Rigor, Specific Sensation.

THE many functions of animal and organic life are variously influenced in maladies which it is the lot of the surgeon to treat; those which are under the influence of the cerebro-spinal nerve-centre more directly, and those over which the cyclo-ganglionic system presides more indirectly. Disturbance of the assimilative functions, in some of its varied forms, is commonly present in any local lesion which is of sufficient importance to produce a constitutional impression; and for the purpose of correctly interpreting these deviations from a healthy standard, it behoves the surgeon to have acquired familiarity with many details in diagnosis, which it was formerly too much the custom to regard as the special province, if not the exclusive privilege, of the physician to expound. The tongue, the pulse, the state of the excretions, all afford invaluable assistance to the surgeon in diagnosis, and, therefore, in treatment; and where it is needed no trouble should be spared, no means of investigation should be considered beneath his careful attention, in order to elicit the requisite information to guide him in forming an opinion. Chemical analysis, especially of the urine, will throw invaluable light on many symptoms which are otherwise inexplicable, and where there are no other more direct means of learning that the kidney or bladder is the seat of functional disturbance or even organic change. And where chemistry fails, the microscope, with its simple lenses, or its polarising arrangement, will supply the required information. The varying development of heat, as tested by the self-registering thermometer, has a significance which requires only the accumulation of facts, statistically arranged, to constitute a far more valuable guide in diagnosis, than is the case in the present limited state of our knowledge on this subject. It is not because the testimony we obtain from this source is, in some instances, apparently contradictory, that we are justified in disparaging its value; such, as I remarked in my last lecture, is often the case in the early steps of scientific investigation; and it is not unfrequently the most unpromising plant, that which requires most jealous watching and most careful tending, which ultimately bears the fairest and the richest fruit.

The state of the *Circulation*, as tested by the pulse and by listening to the beating of the heart, is too important to need a comment. Yet, many precautions are needed, many influencing conditions are to be allowed for, and many accidental circumstances taken into consideration, in estimating the value of the pulse in diagnosis. In some individuals the normal standard is low in frequency, in others high; or the power may be feeble, or the natural pulse strong and full; and these conditions are by no means determined by the general physical development, nor even by the ordinary health of the individual, whether weakly or robust. Again, the nervous excitability of one patient, or the stolid indifference of another, necessarily introduces a discordant element into a source of information which we would willingly have precise. Further, the surgeon will not forget the anatomical lesson he has learned in the dissecting-room, that deviations in arterial distribution may, if overlooked, lead to serious error in testing the pulse at the wrist. For these and other reasons, I think the functional derangements of the circulation must be accepted, in many instances, with the allowance of a considerable discount for possible sources of fallacy, and consequent misdirection. The same remark applies with similar and even greater force to the act of *Respiration*, i.e., apart from the stethoscopic examination of the lungs. It is from the persistence of an abnormal standard in these functions that we derive the most valuable information; especially when the conclu-

sions which we draw from such condition are fortified by collateral evidence of another kind.

Of the *Tongue*, as the index of functional changes which are otherwise hidden from our ken, I entertain a very different estimate. This little member rarely deceives the intelligent inquirer; I mean by its *silent* response to his queries. It tells its simple tale truly; and on its face we may read with certainty not only the day to day variations of our patient, but full often his approaching end. I would trust the tongue, if it gave me encouragement, when almost all beside might seem to forbid hope; and I should know that its glazed and brown surface tell more plainly than the parched member could articulate, that food is useless, the chemistry of organic life is at fault, nourishment has ceased, and that dissolution is near at hand.

Turning from these organic functions, which are under the guidance of the cyclo-ganglionic system, we will now contemplate briefly other functions which are controlled by the cerebro-spinal centre of nerve-power. Under this head we have to consider the intellectual faculties, voluntary movement, common and specific sensation, and likewise the excito-motor movements which are associated with the spinal cord as their source or centre.

In the deviations of these organs from a healthy condition the surgeon is deeply interested; and especially so in the direct appeals which are made to them in accidental lesions from violence. In a future lecture I shall have occasion to illustrate this subject somewhat fully; at present I shall confine myself to noticing the indirect influence exercised on the nervous centres, as exemplified in the aberration of these functions from a normal standard.

Wandering of the *Intellect* occurs in a variety of circumstances; and, as a casual occurrence or taken *per se*, it is not to be trusted with any confidence as a diagnostic symptom. Not unfrequently I have known temporary delirium to occur after an operation, without being accompanied or followed by any circumstance to create anxiety. This may be the result of mental reaction after much tension or subdued feeling; it may be the consequence of wakeful watching, or the effect of the anodyne administered to relieve it. It is rarely produced by pain, unless long continued and severe; but it often supervenes in extreme feebleness and exhaustion, whether from hæmorrhage, profuse suppuration, or any other source of depression of the vital powers.

Yet, though often deceptive by itself, when viewed in connection with other circumstances, it is frequently a very alarming symptom. When accompanied by a calm and even pulse and a clean tongue, we know that the wildest delirium is the offspring of nervous irritation, and not of inflammation. But when associated with other indications of febrile excitement, anxiety is naturally aroused; and if, further, such disturbance is consequent on some injury to the head or spine, we have little room left to doubt that the nervous centres or their membranes are the seat of inflammation.

The delirium of nervous excitement—nervous delirium—and that of drunkards, delirium tremens, or, as a critical scholar and physician under whom I studied medicine used to term it, *delirium cum tremore*,—have much in common. Indeed, it has been denied by many surgeons, amongst whom is Dupuytren, that they are distinguishable. Probably this is altogether so in many cases; the quiet pulse, the clean tongue, the excited talk, the restlessness, the profuse sweating, are characteristic of both; but the tremor is more uniformly present in one than in the other; there is not sufficient difference, perhaps, to constitute this a diagnostic distinction between them; but still tremor is rarely absent in the drunkard's delirium, and not unfrequently so in the subject of nervous delirium. Perhaps I should express myself more accurately by terming the latter *agitation*, the former *tremor*. Each is attended by a remarkable insensibility to pain.

There is a close analogy, if not identity, between the delirium and other symptoms which attend the later stages of an exhaustive disease, such as gangrene, and that which is the sequel of typhus or typhoid fever. Is this wandering of the intellect the consequence of a specific poison circulating in the brain, or simply indicative of the declining powers of life, and the presence in the blood of the uneliminated products of organic conversion? Probably the latter, when unattended by excitement, and of that low muttering form which babbles incoherently of things long forgotten, and unintelligible to, or unheeded by, the bedside watcher. Misguided by the failing senses, the hands wander forth in quest of imaginary objects; and the poor wrecked intellect is busied with its own suggestions, as life ebbs away and leaves it stranded on the confines of eternity.

Here, as the fittest opportunity, I will introduce a few remarks on *Expression*, in attitude and feature, as an adjunct in surgical diagnosis. I say in attitude as well as feature; because in the former there is often much which is expressive of pain or ease, and also of the character of the suffering. The mode in which the arm is carried in axillary dis-

location; the position of the outstretched leg in luxation into the thyroid foramen; the drawn-up knees in peritonitis; the uplifted arms in dyspnoea; the curving spasm of tetanus; the retracted limb in knee and hip disease, are all more or less characteristic of the maladies with which they are allied. Again, how much expression is there in the restless tossing to and fro of the nervously-excited patient, and in the languid, extended and flaccid limbs, and outspread hands of the worn-out sufferer. The hypocrite is sensible of the influence of posture in imposing upon the credulous and uninitiated; and the imitative faculty of hysteria in this respect is often such as to deceive even those who are alive to its Protean characteristics. Freedom from pain is expressed by an easy and unconstrained attitude, which may be detected even in the recumbent posture. I remember, among the many anecdotes which he used to gratify me by telling of his uncle, that Mr. Bransby Cooper said Sir Astley always augured favourably of a patient, who received his visit to his bedroom with his hands clasped behind his head. I have often verified the correctness of this diagnostic sign of ease of body, as well, generally, as of mental quietude. It is scarcely necessary to add that the surgeon's knowledge and sagacity should supply him with a ready tact in convicting an impostor of assuming an attitude which is inconsistent with the malady he wishes to affect; the detection of fraud may often be accomplished in this way.

In the *Face*, as the index of the mind and feelings, we have a guide on which, when allowed to tell its own tale, we may place much reliance. We are all, naturally, physiognomists; and our ability in reading expression is sharpened by contact with the world; but more truly so by familiarity with disease and suffering. In the intercourse of society, good breeding requires that the facial expression should be under control; though, I fear that, from selfish motives, the features are too often made to subserve the purposes attributed by Talleyrand to words, viz., that of concealing ideas. But, in the sick room, human nature is stripped of this mask, and physiognomy may be studied usually without such fear of deception. Each feature has its own peculiar language, but the meaning is generally more intelligible in their combined expression than when they are studied separately.

The tint or colour of the skin is, especially in our climate, often though not invariably a measure of the state of health; more so, however, in youth than in the middle period of life or in old age. But the normal diversity of tint is so manifold, that it behoves the practitioner to be cautious as to the inferences he may draw therefrom. To the surgeon the information drawn from this source is of far less value than to the physician, in special diagnosis. To the latter, the yellow tint of jaundice, the blue colour of cyanosis, the copper dye in suprarenal disease, and the ashy hue in ague, are characteristic signs, severally, of these various maladies. It is said, indeed, that there is a peculiar tint of skin allied with malignant disease: but its frequent absence deprives it of its worth. Not unfrequently disease of unequivocally malignant type is associated with a healthy complexion and well-nourished frame.

On the aspect and the expression of the features in disease much might be written; it is a prolific subject, and yet one which, in its bearing on diagnosis, it is difficult to reduce to system and rule. The expression which betrays various degrees of pain is familiar to all observers; indeed, it is intuitively known and understood. But it may be, and is, easily simulated, for the purpose of eliciting sympathy, or of extorting admiration for fortitude shown. Yet, there is a silent utterance in the compressed mouth, dilated nostril, and upturned eye of genuine suffering, which makes a more forcible appeal than the violent distortion of feature which is indulged in or assumed.

The expression of the face may aid the surgeon in his diagnosis, when taken in conjunction with other signs or symptoms. Such are the crimson spot of hectic, the pinched features of peritonitis, and the peculiar but indescribable combination of anxiety and vacancy in shock. How anxiously does he look for any evidence of loss of symmetry in the face of a sufferer from head-injury; and often the first look of greeting cheers or depresses him in regard to the improvement or retrogression of his patient, whatever may be the disease of which he is a victim. I shall have occasion again to refer to this subject; and will, therefore, leave it with this cursory notice, and proceed to comment upon some other functions allied with the cerebro-spinal nerve-centres.

The exercise of *Volition*, as evinced in muscular movements, is liable to be influenced by any cause which is operative upon the nerve-centre from which it derives its power. Such influence may be transient or persistent, local or general, the consequence of functional sympathy or of organic lesion; or, lastly, it may be due to some cause operating on the brain, or on some intermediate part of the nerve-communication between that centre and the paralysed muscles. The state showing this disturbance varies in intensity from partial to entire loss of power; or it may include the signs of irritation—local or general spasm. Further,

there are accidental circumstances relating to lapse of time or to special seat of injury and treatment, which materially modify these conditions both in degree and character; but of them I shall have to speak more especially in my next lecture, and, therefore, I will defer any further observations on this subject until then.

The phenomena witnessed in association with the true spinal centre and the excito-motor system of nerves are deeply interesting and of great significance in the diagnosis of surgical as well as medical disease. They are essentially distinct from, though often complicated and sometimes confused with, those referable to the encephalic centre; and they are elicited by motive causes acting upon the cord itself, or upon the peripheral distribution of the nerves. The fact that the phenomena in question are, in some instances, more pronounced or more apparent when the cerebral functions are interfered with or in part suspended, enhances their interest and value in the diagnosis of brain or spinal injury; and the recognition of the special functions of this system, and their alliance with the cord as their centre or source, in some part (not yet identified) of its complex structure, has led to an accuracy in diagnosis, which has already borne ample fruit in the treatment of disease, and still offers a field for further investigation, which has been and continues to be successfully cultivated by the physiologist, the anatomist, and the practical physician and surgeon. It would be invidious to attempt to enumerate the many labourers in this deeply interesting field of science: but I may be permitted to mention Sir Charles Bell, as the pioneer of this century; Marshall Hall, whose name is imperishably associated with these discoveries; Brown-Séquard; and Lockhart Clarke, who is still prosecuting his admirable and fruitful investigations in this delicate branch of histology.

Local or partial morbid manifestations of excito-motor phenomena are of frequent occurrence, and may be occasioned by trivial causes, acting directly upon distant organs, and mediately or by sympathy upon the true spinal centre. The varying and doubtful character of the exciting cause in determining an uncertain result, deprives these pathological indications of much of their diagnostic value; but where the characteristic spasm extends more generally throughout the frame, the source is, or becomes by transfer, more directly central; as in tetanus, chorea, epilepsy, and hysteria. In some instances there seems to be reason for believing that persistent general symptoms are due to organic change in the cord itself, by transfer of irritation from the wounded periphery of a nerve. I have been favoured with a view of some preparations of Mr. Lockhart Clarke, which appear to demonstrate this fact, and to which I may again refer in a future lecture. In spinal lesions from violence there can be no doubt that such is the case. Partial and limited damage of the cord may produce involuntary spasm; and organic change following injury may develop tonic spasm or rigidity of the limbs after volition and sensation are suspended.

Rigor is an excito-motor expression, often of great value, in diagnosis, to the surgeon. But its full significance is elicited only by association with other symptoms; by itself, the most severe rigor may denote nothing more than the response to a transient appeal which has been made to the sentient extremities of the excito-motor nerves; as, for instance, by the introduction of a catheter into the bladder, or by the passage of a renal calculus along the ureter. Rigor may be due to cold applied to the surface of the body, to some depressing mental influence, or to miasmatic exhalation when it assumes an intermittent type. It is found to occur, indeed, in all forms of blood-poisoning, and in vital inflammations, which are allied in many of their phenomena, though unintelligibly so in their cause, with toxæmia. But, when associated with symptoms which are suggestive of suppuration, the critical rigor marks the period often with unerring accuracy; or this indication of the formation of pus may, from the obscurity of the case, be the first symptom to arouse suspicion as to its true nature. Probably under no circumstances does the operating surgeon look with more alarm on this symptom, than when allied with an accelerated pulse, a hot skin, and profuse perspiration, under circumstances which induce him to apprehend the occurrence of pyæmia. Rigor is so usual—I may say so constant—an event as ushering in this fatal malady, that, although its presence may not be accepted as conclusively diagnostic, in combination with other indications, of the disease, yet the absence of this symptom may induce the surgeon to be hopeful, even when the condition of the patient might otherwise excite great anxiety.

Many circumstances indicate an affinity between rigor and other convulsive affections, such as epilepsy and the convulsive stage of shock. It is not unnatural there should be this relationship, as the spinal centre is appealed to in all, the points of difference being in degree of intensity or duration, and in the fact that the encephalic centre is involved in the severer forms of epileptic convulsion. An interesting phenomenon connected with these conditions is the variation of temperature in the different stages. Rigor is marked by a rise in temperature, and in the

subsequent stage of sweating it falls, in some instances, below the normal standard. What may be the proximate cause of these changes, so entirely at variance with the feelings of the patient, is matter of conjecture. Dr. Johnson, in some interesting observations on this subject, remarks that probably the sympathetic plays an important part, by its influence on the muscular contractility of the capillaries; and that there seems reason for believing that in rigor there is an anæmic condition of the cord; and, in epilepsy, also an anæmic condition of the brain. Of epileptiform convulsion, in association with injury of the spine or head, I shall have to speak hereafter.

The information to be derived from the various states of *Sensibility* or common sensation is very similar to that which is acquired from the allied function, as regards its cerebral source, of motor power. As in the latter, so in the former, the cause of disturbance may be transient or local, or it may be such as to entail permanently suspended function over a considerable portion of the frame. The frequent coincidence of symptoms, indicating simultaneous derangement of the exercise of voluntary power and of sensibility, is accounted for by the anatomical relations of the centres of nervous influence, and of their prolongations in the spinal canal. I shall return to this subject in a future lecture; but there are certain conditions associated with common sensation, which it will be well to examine at the present time, as they have a more general application than in their special relation to the lesions of the nervous centres.

I have already remarked that intoxication has the effect of lowering the sensibility by its direct influence on the brain. I have observed a similar condition in severe shock, and apparently from the same cause. Loss of blood does not induce this effect, unless complicated with shock, or to such an extent as to cause faintness. Hysteria influences sensibility in various ways; in some instances suspending it for a time entirely, in others exalting it locally, and even generally. Depressed sensibility is, therefore, an equivocal diagnostic symptom, which may be interpreted in a variety of ways, and requires to be used with caution. But when the tactile insensibility of any part is complete and persistent, without evidence of nerve-lesion, the indication is much more definite, and denotes, as the case may be, pressure on, or disorganisation in, some portion of the encephalic centre, or of the spinal cord. Transient paraplegia may result from shock to the cord; but this symptom, under those circumstances, is limited in duration, and has no tendency to increase; extension upwards of the loss of sensibility is a fatal indication of organic or vascular lesion.

Localised tactile insensibility is indicative of pressure upon the supplying nerve, or of some interference, material or functional, with the nerve-centre from which it springs. But the motor nerve of the face is more obnoxious to such lesion than any of the sensitive nerves passing out of the skull; it is frequently paralysed by injury or disease of the petrous bone, or by the pressure of a tumour in the parotid gland. Under these circumstances this symptom is diagnostic; but it should be remembered that the facial nerve is also liable to functional paralysis from cold.

Hyperæsthesia, or exalted sensibility to touch, is to be distinguished from pain. It has a certain analogy to spasm or rigidity in muscles, and is often due to the same cause, viz., some irritating agent applied to the nerve-centre. This is sometimes sympathetic, as in the tenderness, as well as pain, referred to the knee in hip-disease. The presence of simple tumours is sometimes accompanied by remarkable sensitiveness,—as great as, or even in excess of, that which attends malignant growth; and, therefore, this symptom cannot be regarded as diagnostic of the nature of a tumour, but rather, in many instances, of a preternaturally susceptible nervous system. There is a rare form of skin-tumour, of which I have met with some few examples; it is usually very small, and remarkable sensitiveness is its chief characteristic. I remember removing one from a lady's thigh, which was the seat of such exquisite sensibility, that it was a source of almost constant suffering to her; yet this tumour scarcely exceeded a barleycorn in size.

The feeling of heat or chill is a modification of common sensation, which has nothing to do with the actual temperature of the body; indeed, as I just now remarked, a sense of chill is often accompanied by a rise in the thermometer, and the converse is also the case. The occurrence of these alternations in sensation is indicative of a disturbing cause, usually affecting both nervous and circulating systems apparently simultaneously. This condition often precedes rigor, and introduces fever. The sense of chill is to rigor what the epileptic *aura* is to the convulsive fit—the premonitory symptom of the true convulsion, or, it may be, its mild and transient representative. As a diagnostic symptom, this sensation is too inconstant to be of value, and it is often unimportant by itself, though deserving attention in concert with other symptoms, when we are seeking to clear up some obscurity in a perplexing case.

The nerves of *Specific Sensation* evince, in their various deviations from a normal standard and a healthy condition, different states of disease in the organs themselves, which it is not my intention, as it would be beside my purpose, now to discuss. Unquestionably these morbid changes are of infinite importance in the diagnosis of these special diseases, yet it would not subserve any good purpose that I should simply enumerate them; but they are also indicative or symptomatic of other diseases, and this circumstance imparts to them an interest and significance quite independent of the special affection of the organ implicated.

With a few remarks I have still to make on this subject, I shall close the present lecture, and resume the thread of these general observations in the next.

The insensibility or morbid sensitiveness of the *eye* to light are, when not due to any local cause, indicative of cerebral disturbance. This is the case likewise with the other senses; yet this disturbance is not necessarily an organic lesion, or, if so, it may elude our scrutiny. This remark applies rather to increased sensitiveness than to dulness or loss of perception. Indeed, the former condition is closely allied with that hyperæsthetic state of which I have already spoken, as sometimes existing in the nerves of common sensation, and it is frequently met with, and diagnostic of, similar morbid changes. Yet, since writing the above, I have met with a singular instance of temporary suspension of sensitiveness existing in the retina, or, more probably, in the perceptive organ in the brain. A youth, who had fallen on his head, was admitted, with slight concussion, into the hospital. His most marked symptom was total loss of vision; he could not even perceive light. This condition was very transient; in half an hour he could just detect the presence of my fingers when interposed between his eyes and the bright day-light, and in the course of another hour he had recovered natural vision. He made a speedy recovery.

The foregoing remarks are equally applicable to corresponding conditions of the organ of *hearing*, but with this reserve: that, in diagnosis, the latter is less to be depended on, because the morbid changes to which it is liable are more occult, and therefore cannot be referred with the same certainty to a local or cerebral origin. In the details of these functional or organic derangements, there is a close analogy between the two organs. The flitting notes, the dancing sparks, or the flashing light of varied colour, and the low, humming murmur, the continuous rush, or the intermittent tinkle, as of the sheep-bell on the hill-side, and other similar conceits of sight and sound, mark these derangements, though generally with too much obscurity to be surely identified with their origin. It is not infrequent in certain cerebral conditions, and under the deceptive prompting of the distempered sense, for the imagination to frame, out of these fictions, definite sights and sounds. There can be little doubt that such is the case in delirium, especially that of drunkards; and in the dying, these impressions assume a character in accordance with the habits, or, it may be, the hopes or fears of him who is about to sleep the sleep of death. The arms of lost and loved friends seem to be extended, in waiting attitude, to receive the spirit which is struggling for its release; or the averted eye and horror-struck countenance may be in unison with the exclamations which betray the terror occasioned by the imaginary sight of the emissaries of torture. The recurring boom, as of a distant church-bell, will conjure up the vision of the impending funeral procession, or the ideal sound of soothing music may light up the dying countenance with placid hope. As with the eye, so with the ear, a morbid sensibility to sounds denotes increased sensitiveness in the perceptive organ, and usually indicates cerebral excitement; dulness of hearing or absolute deafness is valuable in diagnosis only in connection with other circumstances. It may indicate suppuration within the petrous bone or fracture at that part, when associated with paralysis of the facial nerve. When separately affected, the mischief may be cerebral, or it may exist in the auditory passage or drum. I have known deafness to precede suppuration of the mastoid cells, and in another case necrosis of the squamous portion of the temporal bone, the duration of the affection being too long to ascribe it to obstruction in the external canal; in company with complete blindness of one eye, I have known deafness to supervene on paraplegia following injury to the lumbar spine.

The senses of *taste* and *smell* are still more closely allied than those of sight and hearing; because there is not only an analogy in their morbid derangements, but likewise a close association and interdependence in their functions. It is rarely that one organ is troubled without the other being involved; the loss of scent entails the loss of taste. But there is little to be gathered from these morbid conditions. A disagreeable sense of odours or taste implies perversion of the organs, which is often met with in fever and other maladies, involving derangement of the assimilative functions, yet without being specifically characteristic, except in confirming what is learned from other symptoms: but their successively suspended office sometimes indicates the progress of disease in the brain,

when vegetative life alone survives; or, it may be, as I have witnessed, that the intellect is left to watch, unmoved, the onward march of that advancing disorganisation which marks the degradation of the earthly tabernacle, calmly anticipating the moment when the last prop shall be withdrawn from the tottering fabric, and seeing in "the damp vault's dayless gloom" the welcome portal to unbounded liberty and unfailing light.

OBSTETRIC MEMORANDA.

HEAD AND HAND PRESENTATION.

[By R. B. RUDDOCK, Esq., Clifton.

DR. TENISON'S case in the JOURNAL of the 4th inst. reminds me of a similar presentation I met with last year.

Mr. Hore called on me late one night, asking me to assist him in a case of turning. The woman had been in labour twenty-four hours; the membranes had ruptured eight hours. On visiting the patient, the midwife, an experienced woman, who had called in Mr. Hore, told us that the case had made considerable progress in his absence, and was going on very well. Under these circumstances, we thought it prudent not to interfere, and left. Next morning, ten hours afterwards, we found the patient in precisely the same state we had left her over-night. Mr. Hore administered chloroform whilst I turned the child. It was by no means an easy matter, jammed as it was into the pelvis. She was under chloroform about two hours, and about three ounces were used. The child, a large male, was dead. The woman made a good recovery.

HEAD AND RIGHT HAND PRESENTATION, TOGETHER WITH FUNIS.

By C. H. COLLINS, Esq., Chew Magna.

MRS. H., aged 35, in labour with her fourth child, had been attended for some hours by her mother, who acted as a midwife. On my arrival, I found about two or three inches of the cord down externally, and, on examination, the head and right hand presenting. The pains were regular and persistent; but the expulsive efforts appeared unable to overcome the difficulty of forcing the head, which was, together with the hand, rather firmly impacted in the brim. All pulsation had long ceased in the cord. Not having my instruments with me, I despatched a messenger for them, with the intention of performing craniotomy, as it was quite impossible to force the hand back. During the interval, not wishing to be idle, I succeeded, by a little manipulation, in hooking my finger over the ridge of the orbit; and, by using as much traction-power as I was able to command by that means, during the expulsive efforts of the pains, I managed to deliver the patient of a dead child, of full size and development. The patient herself made a good and rapid recovery.

PUERPERAL CONVULSIONS SUCCESSFULLY TREATED BY DEPLETION.

By W. WEAVER JONES, Esq., Cleobury Mortimer.

IN the year 1863 (May 23rd) I reported in the BRITISH MEDICAL JOURNAL five cases of puerperal convulsions successfully treated by depletion. All the patients are now living. I now beg to forward the notes of another case that occurred on the 29th of May.

J. C., aged 24, married, of full habit, primipara, was at the full period of utero-gestation. Labour commenced at 5 A.M.; she was seized with convulsions immediately after a thunder-clap, 8 A.M. Several fits occurred, and at 10 A.M. she became totally insensible. I saw her at 2 P.M.; she had had up to this time seven convulsions; the pulse was 120; the countenance blue; the breathing stertorous; the pupils dilated; the tongue protruded and swollen; the os uteri was open fully, and the membranes tense. She was bled to twelve ounces; the membranes were ruptured; cold was applied to the head. She was delivered by forceps of twins; the first child was a male, still-born; the second a female, living. Up to 8 P.M. she had five convulsions, not so violent; the breathing was less stertorous, but she was quite insensible. An injection of gruel, with two ounces of brandy, was administered. On the 30th, at 8 A.M. she had had four convulsions since 8 o'clock on the previous night, the last occurring at 1 A.M. She was now conscious: another injection was administered. She has progressed favourably since.

REPORT OF THE EDINBURGH COMMITTEE

ON

THE ACTION OF MERCURY ON THE LIVER.

By J. HUGHES BENNETT, M.D.,

CONVENER AND REPORTER.

PART I.

It may be remembered that, in an Address I had the honour of delivering two years ago at Chester, I urged upon the British Medical Association the propriety of doing something, in its collective capacity, towards solving one or more of those doubtful points in practical medicine connected with the action of medicines or the treatment of diseases. The proposal, having met with the general approbation of the meeting, was committed to the Council; and, in the middle of the following October, I received a letter from the Secretary, informing me—"That, in accordance with a resolution passed at the last annual meeting, the sum of £25 be granted, out of the funds of the Association, to a Committee, to be appointed by, and under the direction of, Professor Hughes Bennett, to investigate the Action of Mercury on Animals." After hesitating for some time, in consequence of the inadequate sum voted, and after communicating with the President, who explained that although small there was no further balance available, I finally accepted the commission. In doing so, I was actuated by the feeling that this first effort of the Association to investigate an important medical question, with the aid of its funds and by a combined effort of its members, ought to be prosecuted at all risks; I also felt confident that an earnest effort in this direction would ultimately receive the support of the Association.

My next step was to secure the co-operation of gentlemen who were capable in various ways of assisting the inquiry; and ultimately I secured the cordial co-operation of—Dr. Christison, Professor of Materia Medica and Therapeutics, Edinburgh; Dr. MacLagan, Professor of Medical Jurisprudence, Edinburgh; Dr. James Rogers, formerly of St. Petersburg; Dr. William Rutherford, Assistant to the Physiological Department of the Edinburgh University; Dr. Frazer, Assistant to the Materia Medica Department of the Edinburgh University; and Dr. Gamgee, Assistant to the Medico-legal Department. The Physiological, Materia Medica, and Medico-legal Laboratories of the University were also to be placed at our disposal.

The Committee thus formed appeared to me to unite in itself all those elements required to give confidence in the results arrived at; viz., "The skill of the anatomical operator, the analytical power of the chemist, and the varied knowledge, theoretical and practical, of the histologist, physiologist, physicist, pathologist, and therapist, as well as of the physician whose knowledge of diagnosis is unimpeachable." In the Committee thus constituted, it appeared to me, every confidence might reasonably be placed, as "the energy and sanguine character of youth were sufficiently tempered with the caution and reasoning power of age, while every necessary instrument, appliance, and chemical, together with a public hospital, were at its disposal."

The Committee met on Nov. 16th, 1866, and, after discussing what branch of this extensive subject should be first investigated, it unanimously agreed to direct its attention to the Action of Mercury as a Cholagogue. On proceeding to consider by what method this was to be accurately ascertained, it was clearly pointed out by Dr. Gamgee, as the result of the opinion of modern chemists, that no kind of examination of the feces would yield trustworthy results. Supposing that the chief and characteristic constituents of the bile found their way into the alvine evacuations unchanged, imperfection in the analytical methods at our disposal render their quantitative analysis impossible. The plan of ascertaining bile-acids indirectly by means of nitrogen and sulphur determinations of the alcoholic extract, while most unsatisfactory in the case of