douche, but to cool the blood, and thus to restore the contractility of the minute arteries of the lungs. The condition of the pulmonary vessels in this disease is the exact opposite to their state in cholera collapse. In cholera collapse, the minute pulmonary arteries are in a state of extreme contraction; and, as a consequence, the capillaries are extremely anæmic. In heat-apnoea, the pulmonary arteries are extremely relaxed; and the capillaries, consequently, are excessively engorged. In cholera collapse, external warmth in some degree, but much more rapidly and decidedly a warm injection into the veins, relaxes the arterial spasm, and restores the circulation. In heat-apnoea, on the contrary, the object is to cool down the over-heated blood, so to revive the contractile power of the minute pulmonary arteries, to relieve the capillaries from their embarrassing excess of blood, and thus to remove the state of apnoea. A clear apprehension of these physiological principles cannot fail to be of great assistance in practice.

In the treatment of heat-apnoea, the following appear to be the main points which require attention. The patient should be placed in a recumbent position in the coolest possible place, with a free current of air. The clothes should be removed, and cold water applied to the whole surface; or, if the symptoms be urgent, the clothes should immediately be saturated with cold water without waiting to remove them. If the respiratory movements be failing and feeble, the cold douche is a powerful excitant; but if the breathing be rapid and laborious, it is better to envelope the body in a wet sheet, and to quicken evaporation and cooling by a fan or a pair of bellows. If the patient can swallow, let him drink iced water freely. Whether he can swallow or not, iced water may from time to time be injected. The marvellous effect of hot venous injections in cholera collapse, and the urgent need for cooling the blood in heat-apnoea, suggest the expediency, in extreme cases, of injecting into a vein the same saline solution as has so frequently been employed in cholera, only injecting it cold instead of hot.

Dr. Morehead (*Clinical Researches on Diseases in India*) recommends venesection in the early stages of the attack, while Dr. Maclean condemns the practice. A routine practice of venesection would be destructive; but, when symptoms of excessive venous engorgement are present, a cautious venesection would be quite justifiable, and probably beneficial, on the well known principle of lessening distension of the right side of the heart, and thus increasing its contractile power. When respiration has suddenly and quite recently ceased, artificial respiration by Dr. Silvester's method may possibly restore animation. While symptoms of apnoea continue, however great may be the apparent exhaustion, no alcoholic stimulants are to be given, for the reason that alcohol, as well as anæsthetic vapours and narcotics, impede oxidation of the nervous and other tissues, and therefore increase the risk of death from apnoea. Ammonia may be applied to the nostrils as a stimulant, and, if the patient can swallow, it may be given internally. Ammonia is a powerful diaphoretic, and the restoration of the cutaneous secretion is an important step towards recovery. When the skin becomes cool and moist, of course all cold applications are to be discontinued. To sum up then—as *hot air* and *hot blood* are the cause of this form of apnoea, so *cold air* and *cold water* are the chief means of cure; all other means are subsidiary to these.

NOTES ON THE SURGERY OF THE NINETEENTH CENTURY AND THE OPHTHALMOSCOPE.*

By JAMES VOSE SOLOMON, F.R.C.S.,

Surgeon to the Birmingham and Midland Eye Hospital, etc.

In the very eloquent and learned address on Surgery, delivered at the Dublin meeting of the British Medical Association by Professor Smith, it was deplored that so little attention is paid, in the present day, to the Patristic literature of our profession; and it was asserted that in those noble writings were to be found some things which had in our own day been presented as new discoveries.

Assenting then, as I do still, to much contended for by Professor Smith, a man of great modesty and of rare gifts and acquirements, I yet felt at the time of the delivery of his address, and on reflection find no reason to alter my opinion that the nineteenth century surgeon has no occasion for shame at the present state of his art. Can he not fairly boast that in his day scalpel and microscopic human anatomy attained perfection; that the true nature of the inflammatory process was understood, and a rational and successful treatment adopted for its relief? He can

claim that in his day the knife was robbed of its terrors by chloroform and local anæsthetics; while Eve's curse was mitigated by the same divine agency. The surgery of tendons, that of the iris, and that of acute glaucoma, were perfected.

I believe the future historian of the epoch will recognise as characteristic of it a marvellous improvement in the diagnostics of disease, and a large diminution of death-rate; great economy of time in the attainment of a stage of convalescence, with considerable diminution of the restraints and severities incident to our methods of cure; the substitution, for destructive operations, of procedures which reduce the parts submitted to surgical interference as near as is possible to a physiological state; the cure of certain diseases by bloodless operations, where the knife had ever before been the sole remedial resort; the healing of large wounds without occurrence of the process of suppuration. He will dilate with admiring wonder upon the rapid perfection of ovarian surgery, and the accomplishment of this by the labours of scarce more than twelve men—a death-rate of twenty per cent. having been attained. As a striking instance of economy of time, of recovery, and of physiological surgery, he will adduce excision of the elbow-joint in cases of articular disease, whereby years of suffering have been abridged to an easy and almost painless recovery extending over no more than a few weeks. In the list of bloodless operations, we doubt not, will be found the cure of stone in the urinary bladder through its eliminative canal by lithotripsy; and, greatest of marvels, the consolidation of aneurism, even of the descending abdominal aorta, by compression. On the other hand, the historian will find much to tell of the invasion, and successful subjugation, by surgery, of districts which had, from the birth of the healing art, been considered as belonging to the domain of pure medicine; as, for example, the successful treatment by operation of acute glaucoma, reflex ophthalmia, recurrent iritis, and, it may be, of myopia. He will attribute a large measure of our progress to the important aids contributed to minuteness and accuracy of diagnosis by physical science, and by mechanical ingenuity and skill. He will relate that the living bladder ceased to be a dark obscure cavity; its diseases being pictured to the surgeon's eye on the field of the endoscope, and encysted and loose calculi being accurately differentiated. He will tell how the pathology of the larynx, painted on the most ancient of mirrors made by the hand of man, came to be unravelled; how the site on the voice-string from which the polypus had been snared was cauterised with unerring certainty by the electric cautery. He will tell how the pulse wrote its own diary with the sphygmograph. He will dilate on the complete revolution effected in our treatment and views of the nature and pathology of eye-diseases by the discovery and use of the ophthalmoscope. He will, no doubt, attribute the rich harvest of fruit gathered from some of these methods to the enlistment of the sense of sight as an aid to that of touch, or as the sole exponent of the characteristics of the diseased changes—a sense which affords ideas and impressions altogether different from those obtainable by any other means whatever.* While speaking of these triumphs in the art of diagnosis, I cannot here refrain the remark that the tendency of the inventions and methods of the surgery of our period, which has become largely mechanical, is to put more nearly on a level those of gifted intellect and those of moderate understanding. It cannot be doubted, I think, that this levelling process is effecting much good by diffusing the blessings of our art more widely amongst humanity.

Nevertheless, the division of labour, called Specialism, will be maintained, if not increased; and through such division the great body of the profession will, as heretofore, acquire more distinct views of the pathology and appropriate treatment of certain diseases. The public is every day becoming more firmly convinced that their interest lies in supporting this arrangement; and I believe the profession will ere long be quickened to recognise that a cordial co-operation between specialist and general practitioner should become an universal rule, because subserving in the most complete manner the beneficent and benevolent purposes for which the profession exists.

The death-rate in ovariectomy has been reduced, as I have said, to 20 per cent. In 333 cases of excision of the elbow-joint, 12 per cent. died; 4.2 suffered amputation; in 6.9 per cent. there was ankylosis; and in 256 cases, or 76.87 per cent., there was mobility of the joint. One hundred adult cases of stone in the bladder, occurring consecutively

* Although the touch has been habitually resorted to for ages as a means of diagnosis, the common existence of ulcerative and inflammatory hypertrophy was never even suspected until the speculum revealed their presence. When once, however, the eye has demonstrated the existence of these morbid states, the touch, with its assistance, generally acquires the power of distinguishing the most minute changes; and it then appears quite a marvel to the practitioner that the grossest morbid conditions should previously have always escaped his recognition. In illustration of these remarks, Dr. J. Henry Bennet quotes an instance where an eminent surgeon who had devoted his whole life to the study of uterine disease, mistook, thus relying on touch alone, advanced cancerous ulceration of the os uteri for a simple ulcer, and promised that the patient would soon recover.—*Inflammation of the Uterus*. By J. H. Bennet, 2nd edition. Pp. 494-6.

* Read before the Birmingham and Midland Counties Branch.

in the practice of Sir Henry Thompson, were all, without exception, submitted to operation. Of 84 who underwent lithotomy, 4 only died. The mean age of the whole number operated upon was 62½ years. There cannot be a doubt that this death-rate will be much diminished as the public obtain more correct information on the subject.

Humbly craving, Mr. President, your forgiveness, and that of the Society, for this digression, consisting of thoughts which presented themselves to me during my holiday in Ireland, I will, if your patience be not exhausted, request a few minutes' attention to the subject of the Ophthalmoscope and Optic Nerve.

The invention of the instrument—barely in the sixteenth year of its publicity—is due to Helmholtz. In consequence of the peculiar structure and arrangement of the parts composing the organ of vision, the ophthalmoscope confers the power of inspecting important living tissues which, prior to its existence, were hidden from our view; and not only this, but of examining them, if the expression may be allowed, in a state of complete nudity. By no other instrument is it possible, in the living man, to inspect, without preliminary dissection, a nerve, or the whole vascular system of an organ of special sense. With no other contrivance is it practicable to watch at all hours the departure from its normal state of a tissue which is a prolongation of the brain itself, the optic nerve. In no other organ, and by no other method, is it permitted to witness that most interesting of phenomena, embolism of an artery, with the grave nutritional consequences thereof. Almost the whole of the local vascular changes incident to the occurrence of a small cerebral apoplexy are seen here in miniature—the capillary or venous congestion; the pouring out of blood in minute spots, or, if from a larger vessel, in quantity; then the alteration of colour of the clot; its rapid, or may be gradual, absorption; and the final effect of the presence of it upon the choroid in the production of atrophy and pigmentation. Suffer me to remark, that effusions of blood on the fundus oculi, unlike cerebral effusions, do not terminate in cysts; they are absorbed; a blanched spot or a deposit of pigment, or both, as already said, marking their past existence.

It will be evident to you that, where impaired vision is traceable to these last named changes, improvement by drugs can seldom be anticipated; and that timely examination by a competent ophthalmoscopist will be the means of averting serious loss of time, expenditure of money, and perhaps constitutional strength, on the part of the patient.

Having regard to certain opinions that have been published respecting effusion of blood on the retina, it is incumbent on me to remark that its presence upon the retina, optic nerve, or choroid, can always be determined with perfect certainty. When the refractive media retain their transparency, there can be no ground for a reasonable doubt. As to the degree of impairment of vision arising from blood effused on the retina, that will depend upon its size and site. The statement made on oath, in an action for damages for railway injury, by a London oculist, that a man without legs was as well able to walk as a man to see who had a clot on his retina, would be received in well informed ophthalmic circles with ridicule. In certain diseases of the kidney, and in cases of near sight, I have often found small hæmorrhages, yet the patient could see to read small pica.

Anteriorly to the employment of the ophthalmoscope, weakness of sight occurring in the course of Bright's disease was attributed to the circulation of urea in the brain. Now it is acknowledged to be due, in the large majority of cases, to special changes in the optic nerve and retina.* Nay, more, I speak my own experience when I tell you that a knowledge of existence of kidney-disease has sometimes been first contributed by an oculist who has explored the fundus. It has been well said, "What might become of many other blood-diseases, if we could but watch the organs affected as we do here?"

The views of those surgeons who everywhere proclaim that secondary changes of syphilitic origin owe their existence to the administration of mercury, receive complete confutation from ophthalmoscopic investigation of the retina, upon which membrane sometimes are discovered signs of specific disease at an early state after the primary affection, and prior to treatment by mercury.

Notwithstanding these and numerous other facts, there remain still certain oculists who deny the utility of the ophthalmoscope for the successful treatment of eye-diseases. The scepticism of these gentlemen led them at first to challenge the possibility of clear definition of the fundus being obtained by this means. After a time, however, under an overpowering weight of evidence, this point was conceded; but now they

solace themselves in an ignorance of the practical application of the invention, by boldly declaring our treatment to be insusceptible of advance by its use.* The Society will perhaps condescend for a moment to examine with me this position. It amounts in many cases to this, that absolutely certain knowledge—ocular demonstration—of the diseased changes producing a malady, is of no more value to the prescribing physician than the general symptoms of such changes, even although those symptoms, as in the case of certain ophthalmic diseases by no means inconsiderable in number, represent no one disease in particular.

Let me take an ordinary instance of complaint, made by perhaps a dozen patients in a single morning at the service of the out-patient department of an ophthalmic hospital—"My sight, sir, is dim." Ophthalmoscopic observations, repeated hundreds of times by different observers, have proved that the cause may be spasm of the lens-muscle, hypermetropia, incipient cataract, a displaced lens, effusion into the vitreous humour, inflammation of the optic nerve or retina, apoplexy of the optic nerve or retina, blood on the vitreous humour, anæmia of the retina, atrophy of the optic nerve, or incipient glaucoma. Some of these conditions would demand no treatment, some immediate operative interference, some mercury, some purgatives, some steel, some strychnine, some change of air. You will, I am sure, agree with me that such instances as those referred to, however worthy of confidence the authors may have been at some period of their professional career, are now but the shifts of men who, having eyes, see not. Being themselves prejudice-blind, they would constrain others, who can see, to turn away from the greatest ophthalmic revelation of our period.†

I refrain from giving instances of serious blunders committed for want of competent knowledge of Helmholtz's method of investigation of the internal structures of the eye. The most dangerous enemies of ophthalmic progress are, I fear, surgeons who profess confidence in the instrument, but who, from some inaptitude or physical infirmity, have failed to master the elementary difficulties which beset all who take it up for the first time, and who yet profess to base their opinions in consultation upon ophthalmoscopic evidences, while in reality they merely make guesses founded on the general symptoms. To make the illusion complete, these persons tell their patients they will make written notes of what the fundus reveals. Such moral leprosy we can only deplore, and hope to lessen by exposure to the pure air of public opinion. "Medicine," says Dr. Latham, "claims a sort of moral respect in the handling; it calls upon the conscience as well as the intellect for more caution to avoid error, and more fearfulness of overstepping the truth." (*Clinical Lectures*, 1846.)

To those brethren who may feel disposed to take up the instrument, I would advise that one of the pupils of the surgeon's eye, as well as that of the patient, should be artificially dilated. I believe that then, for reasons explained by me in the *BRITISH MEDICAL JOURNAL* (June 1867, p. 628), no difficulty will be experienced in focusing the fundus. The normal colour of the optic nerve is not always the same in different persons. It may be pink, rosy, buff, brown, or grey, so variable is the colour of nerve-tissue. As regards inflammation of the optic nerve and retina, singly or combined, producing, when unrestrained, atrophy and blindness, I have witnessed many remarkable recoveries, which were entirely due to the priceless diagnostic precision got from ophthalmoscopic evidence. And I regret to say, cases continually present themselves where, for want of this, vision has been immediately impaired or lost, and the sad spectacle presented of a skilled artisan reduced to the level of a pauper, or made dependent on the sole exertions of wife and children for a maintenance.

I would fain believe enough has been adduced by me to impress my brethren, who have so courteously restrained any impatience provoked by my dulness, with the conviction that ophthalmoscopy is entitled to equal rank in their estimation with the stethoscope and speculum uteri. At the same time, I would not be understood for a moment to desire to disguise the fact that "the senses—the eye, the ear, the touch, however sharp and delicate they may naturally be, require a special course of training and education before their evidence can be trusted." (*Watson's Lectures*, 3rd ed.)

* "And, after all, however painfully and slowly the power of diagnosis has been acquired, it is still imperfect; an assertion well borne out by the comparison of the state of our knowledge previous and subsequent to the discovery of Laennec."—*Diseases of the Chest*. By Dr. Stokes. 1837. Pp. 11 and 12.

† "It has been asserted that, by studying the varieties in the nature, mode of occurrence, succession, and modification of these symptoms, we can, independently of the information derivable from physical signs, arrive at a diagnosis of thoracic disease. This is the common assertion of those few who are still opposed to the use of mechanical diagnosis." (Dr. Stokes on *Thoracic Diseases*.) "As diseases are better understood, and we possess surer signs of discerning their seat and progress and events, the records of past experience become obsolete, and so a necessity arises for a new course of clinical observation." (*Watson's Lectures*, 3rd edit., p. 295.) "The knowledge of the senses is the best knowledge." (*Idem*.)

* It is needless here to enlarge upon the importance of such distinction from a therapeutic point of view, or on how completely the pursuit of ophthalmic practice has bound up with it a competent knowledge of the pathological relation of diseases of remote, but connected organs; e.g., embolism of the retina as a result of valvular disease of the heart; exophthalmos, or protrusion of the eyeball, with a retarded cardiac venous circulation; inflammation of the retina, with pathological changes of the kidney-tissue or brain; spasm of the ciliary muscle, with concussion of the spinal column, as in certain railway accidents.