tion. He mentions also the bromides of mercury, and gives the dose of the subbromide (bromide) from one to two grains, and that of the perbromide (bibromide) from one-sixteenth to one-fourth of a grain; but I have usually found that the subbromide of mercury exerts a greater power than its corresponding salt, calomel.

Having been in the habit of prescribing the bromides of mercury during the past five years, and being impressed with their value as therapeutic agents, I wish concisely to point out their general effects, in the hope of enlisting others in this investigation; so that, by long and continued observation, we may together accumulate such knowledge of their powers as may enable us more accurately to judge of their merits, as compared with those of the chlorides.

I have found, with the protobromide or subbromide, that a less dose than of calomel is required; that it acts more agreeably and effectually, both as a cholagogue and purgative, without the depressing results so often observed to follow the exhibition of calomel. It seems to be useful in promoting the absorption of hypertrophied glandular and other morbid tissue with less constitutional disturbance than the chlorides. I have administered it, with apparent great relief, in cases of biliary congestion of the liver, in chronic hepatitis with enlargement, and in those skin-diseases which yield best to the influence of mercury. I might quote some interesting cases of large anomalous tumours of the abdomen sensibly diminishing, and in two instances wholly disappearing, under the use of these remedies—viz., the protobromide of mercury in half-grain doses, and the perbromide in the form of an ointment of four grains to the ounce of cold cream; also of uterine tumefactions and morbid deposits around the cervix uteri and in the connective tissue of the pelvis, treated with pessaries of the perbromide, as well as the ointment externally. I could cite other instances of acute and chronic inflammatory disease in which the bromides have been apparently of great use; but my observations have been as yet so limited, and the difficulties of making perfectly correct observations on the action of a new remedy are so great, that I think it best to forego such means of proving its value till a mass of evidence can be adduced which may better test the accuracy of my conclusions. With your permission, therefore, sir, I would here ask those members of the Association who may try these remedies to note their effects, and kindly to favour me with a report of their observations, so that I may, at a future time, publish the result of our united efforts; since to arrive at any definite ideas of the action of the bromides of mercury will, I feel sure, require the careful work of many observers.

This short paper, then, claims to be put forward, not as a complete account of a new remedy, but as a notice to the profession, with the view of stimulating others to take up and investigate the subject.

ON CARIES OF BONE.

By HOLMES COOTE, Esq.,
Surgeon to St. Bartholomew's Hospital, London.

I HAVE been long convinced that the results of operations undertaken for the removal of carious bone are unsatisfactory. We see case after case in which the surgeon gouges away as much as he dare of the diseased structure; and yet he finds, when the period for the cicatrization of the wound comes about, that the fistulous sinuses still remain open, and that the annoyances to the patient are as great as before. A second and a third operation may be recommended and performed, and still the relief is incomplete, until, finally, a cure is effected by natural processes.

These reflections lead naturally to the question, What is caries? Is the disease of such a nature as to admit of beneficial treatment by surgical operation?

We find, in reference to this question, no small difference of opinion among surgical authorities. Some mean by the term, simple ulceration of bone; others, chronic suppuration of the cancellous tissue. Mr. Stanley often defined it, in my presence, as "unhealthy ulceration of bone." Mr. Erichsen speaks of it as a disease characterised by increased vascularity, softening, and ultimate disintegration of the osseous tissue.

Perhaps the last definition expresses best the true nature of caries; but it is wanting in accuracy of detail.

Caries may be said to have its seat, with scarce exception, in the cancellous structure of bone, such as the bodies of the vertebrae, the bones of the carpus and tarsus, or the spongy extremities of the long bones. As already observed, the disease commences with increased vascularity (ostitis or osteomyelitis); and I think that I have never known it to be transmitted from the soft parts to the bone. It has always commenced in the bone itself. As the disease proceeds, it causes molecular disintegration of the osseous tissue, which comes away in the form of minute

fragments mixed with the pus. After a time, larger portions may perish and become detached; but, unlike that which is observed in simple necrosis, these carious sequestra do not represent the whole of the diseased part. One of the most intractable forms of caries is that termed *caries fungosa* (Volkman), which is characterised by both ulcerative destruction and the production of an abundant granulation. The bones become mouldering, osteoporotic and thinned, in parts absolutely destroyed, the medullary space being filled by a reddish-coloured marrow. The granulations lose their florid red colour and become pale; there is an abundant deposit of fatty matter in the neighbourhood; and the discharge becomes offensive and discoloured. It is this form of caries which especially attacks the bones of the carpus and tarsus. Portions of eroded bone may be detached, but the morbid action extends far beyond their sphere, and usually more bones than one are involved in different stages of the same disease. The pressure of the granulations is followed by absorption and alteration in contour, while the internal disintegration may terminate in leaving little more than an external shell of compact, but thinned and light tissue.

This is the disease, and such the conditions of the morbid parts, which we in modern surgery attempt to rectify by the operation of gouging away the diseased bone. The results have not realised my expectations, except in cases where bone, already detached, admitted of easy removal, or where an emptied cloaca, surrounded by healthy bone, could be scraped so as to relieve it of its soft and unhealthy lining.

Under the most favourable conditions, the prognosis is far from good; but I am inclined to doubt whether the patient's condition would not be better after a period of twelve months, and simply kept at rest and in good air, than if he had undergone the usual operation.

A CONTRIBUTION TO BABY-FARMING.

By HERMANN BEIGEL, M.D., M.R.C.P.,
Physician to the Metropolitan Free Hospital.

THE following interesting case illustrates some of the evils of the system of baby-farming, which have been so ably exposed in the reports which recently appeared in the columns of the BRITISH MEDICAL JOURNAL.

Rebecca Levy, now twenty-four years of age, was, soon after birth, possessed of a loud and clear voice. When she was one month old, her mother fell ill, and was obliged to become an in-patient at a hospital, and to give her baby in charge of a woman to be nursed. Very little care was, however, taken of the child, who was left by her nurse in a room with the door closed, with very little food, and who consequently cried nearly the whole day. This continued for nearly twelve months, when the mother returned home, and took the child back, the latter being but ill-fed and perfectly hoarse. The girl never recovered her voice, and only experienced a difference in the degree of hoarseness. During the last few months, this condition had become at times alarming; fits of suffocation sometimes setting in, interrupting sleep, which was always accompanied by a whistling, rattling noise. The patient could never lie on her back without experiencing one of these suffocating attacks; and was, therefore, obliged to remain in a sitting posture during the night. On the 27th of May last, she came to the Metropolitan Free Hospital in a state of suffocation, and was admitted as an in-patient under my care. Her breathing was at this time extremely laborious; the ordinary muscles of respiration, as well as the auxiliary muscles, were in powerful action; the alæ of the nose were working violently; the visible mu-

cous membranes were livid; perspiration covered her face and forehead; her speech was very hoarse and thick; and, from want of breath, she could only pronounce short interrupted sentences.

Laryngoscopic Examination.—The pharynx was red. The larynx and vocal cords were as illustrated in the accompanying woodcut. Both