ON OPERATIONS FOR CARIES OF BONE.

By F. LE GROS CLARK, Esq.,

Surgeon to St. Thomas's Hospital.

MANY years since, I discarded the use of the gouge in treating carious bone, and I am glad to find my convictions on this subject confirmed by Mr. Holmes Coote. The progressive disintegration of cancellous tissue is essentially identical, in all its leading features, with ulceration of soft parts. In some instances, this destruction, or death, is rapid, as in phagedæna; or slow, as in scrofulous subjects. The increased vascularity, when present, is really due to the reparative effort, and generally a measure of its activity; and is not a necessary accompaniment of the destructive stage. How do we treat abscess or ulcer, which are, in fact, but modifications, in their mechanical features and relations, of the same condition? We open the abscess, and take care that its contents shall have free vent; but we do not scoop it out, because it happens to be foul from the presence of disorganised tissue; nor do we scrape the surface of a foul ulcer. But we stimulate the sluggish parts with appropriate applications, and extend a narrow or sinuous opening, so as to give free vent to the discharge.

My practice, and what I regard as the rational treatment of these cases, is embodied in the following passage, which I extract from one of my Clinical Lectures, published in the *Medical Times and Gazette* for February 16th, 1861. "As regards operations for the removal of carious bone, I cannot say that my experience justifies my recommending the proceeding of gouging out the diseased texture. My belief is, that violence thus offered to the parts often involves the surrounding healthy bone in diseased action, and thus leaves the patient in a worse state than before. The condition is analogous to that of an excavated ulcer in soft parts, and seems to me to be more amenable to treatment adapted to such state. Sinuses may be freely laid open, and suitable dressing employed, to encourage a healthy action and spontaneous cure. But I am aware the weight of opinion, as regards the general practice, is adverse to this negative treatment."

Such has been my practice for many years; and I may add that I have found considerable advantage in the use of diluted acetic acid (which I have latterly substituted for phosphoric) as a bone-solvent, as well as a stimulant in carious cavities which have been previously laid freely open by external incision. Caries is usually primary; rarely secondary on ulceration of the skin surface, but not infrequently so, by extension of disease from a synovial surface. In the knee-joint, this secondary condition, destroying the cartilage in its progress, is often met with; and I think still more frequently in the tarsus, where the continuity of the synovial membrane favours and guides the extension of the disease.

THERAPEUTICAL MEMORANDA.

TREATMENT OF ECZEMA.

By W. H. AXFORD, M.B., Bridgwater.

APROPOS of Dr. Basham's case of Acute Eczema given in the *JOURNAL* of October 10th, the following case, which is of interest as being apparently due to suppressed menstruation, shows the great benefit to be derived in such acute cases from lotions containing gallic acid.

M. A. B., aged 18, was admitted as an out-patient at the Bridgwater Infirmary on September 21st. She had generally been a healthy girl, and had always been regular until the last period, which she missed a fortnight ago. A few days ago, a rash, consisting of very minute pimples, appeared, commencing at the upper part of the chest, and spreading rapidly to the arms, the whole front of the body, and upper part of the thighs. The rash was a bright scarlet, and exudes a watery fluid, especially on the left arm; in some parts, this fluid had dried, forming yellowish scabs. The itching was intense. There were some signs of indigestion. She was ordered a mixture, containing bismuth and hydrocyanic acid, three times a day, and a lotion.

R Sodæ sesquicarbonatis 3ij; glycerini 3ss; aquæ 3viij. M. Fiat lotio sæpe applicanda.

Sept. 26th. Somewhat better; less itching; but still considerable exudation of watery fluid.

R Acidi gallici 3j; glycerini 3j; aquæ 3viij. M. Fiat lotio sæpe applicanda.

The mixture was ordered to be repeated.

Sept. 28th. Much better; the itching was much less; and there was not nearly so much exudation. No part had yet, however, healed.

She did not again come to the Infirmary until October 3rd, when she appeared quite well, stating that the catamenia had appeared on Sept. 29th, and that the rash immediately began rapidly to disappear.

REPORTS

OF

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS OF GREAT BRITAIN.

THE MIDDLESEX HOSPITAL.

REMARKABLE CASE OF FRACTURE OF THE SKULL: TREPHINING: DEATH.

(Under the care of Mr. CHARLES MOORE.)

THE following case is one of not only great clinical interest, but of considerable importance in a medical point of view. That a man falling from a height, and receiving a blow which caused so extensive and dangerous a fracture, should be able to continue his work and walk a considerable distance with little if any evidence of mischief, is surprising. The case certainly loses none of its interest by the progress and result.

A. M., aged 40, was brought to the hospital in the evening of September 4th, said to have fallen, in the morning of the same day, from a height of ten feet on to his head. It seems that, after the fall, he got up, walked a short distance, and continued to work. He also entered into conversation with his fellow-workmen, and talked to them of his fall. They thought, from his manner and gait, that he was intoxicated. Late in the afternoon, he was found lying insensible in a cellar, and brought to hospital. His extremities were cold; the pupils were moderately contracted, but equal; the pulse 72. Half an hour later, the left pupil became contracted to the size of a pin-point, the right remaining as before. Temperature 98.6. At 9 P.M., Mr. Moore trephined. On making an incision over the right parietal bone, where the skin was broken, and to which the patient was observed to raise his hand, a fracture was found extending along the right parietal bone. A triangular piece of bone was removed, and a large clot discovered beneath the dura mater. The clot was scooped out, and two spouting points observed. They could not, however, be reached so as to apply ligatures. The wound was, therefore, left open, and ice applied to the head. The patient did not rally, but died during the night.

The *post mortem* examination was conducted by Mr. Henry Arnott, Surgical Registrar, to whom we are indebted for the following notes. The fracture of the skull was found to extend from near the vertex on the right side in front of the ear nearly to the temple. On removing the calvaria, some clotted blood was found between the bone and dura mater at the seat of operation; but the membrane was not torn at this point. Lower down, however, and anteriorly, was a rent in the dura mater; and, on lifting the upper part of the membrane, a quantity of black grumous blood, mixed with mashed brain, escaped. On further examination, it was found that nearly the whole right cerebral hemisphere, except about an inch in depth of the upper surface, was torn up, laying freely open the lateral ventricle, and rendering the parts in the base of the brain nearly indistinguishable. The cerebellum, pons, and left half of the cerebrum, were alone free from injury. Of the central parts of the base, the two optic nerves were alone recognisable. The right corpus striatum and optic thalamus were destroyed, and their place occupied by clot and torn brain-substance. The fracture ended in the right anterior fossa, crossing the groove in which ran the meningeal artery. Blood was effused into the posterior part of both orbits. The bone, however, was not broken here. With the exception of the lungs being congested, the other viscera were healthy.

ST. MARY'S HOSPITAL.

PETECHIAL TYPHUS FEVER IN A BOY: RECOVERY.

(Under the care of Dr. SIBSON.)

IN the *JOURNAL* of October 10th we published an interesting case of petechial typhus fever, which proved fatal on the eighth day, under most painful circumstances. We now give a short account of a similar case, which happily, however, terminated in recovery.

In his volume on *Continued Fever*, Murchison truly remarks, that "the ominous character of the livid and petechial eruption has been mentioned by all writers from the earliest period." The petechiæ in this case were unusually severe; they were literally black, which betokened a severe form even of petechial typhus. There were, however, certain indications in this case which gave a ray of hope that he might possibly pull through. These were especially dwelt on by Dr. Sibson in an interesting clinical lecture on the subject. In the first