campaign in 1864, in which Professor Langenbeck performed a subperiosteal resection of a large portion of the diaphysis of both tibia and fibula, and which was brought to a successful termination, with complete re-formation of bone.

Time would not permit me to discuss some other points connected with this subject, which I consider of the utmost importance. I shall content myself, therefore, with stating the following propositions, which are based on the physiological experiments and clinical experiences of M. Ollier especially, Langenbeck, Lücke, Moon, Wood, my own, and others who are interested in this subject. These are:

1. That, in subperiosteal resections, the reproduction of bone is more complete, and effected with greater rapidity, than after total removal of both bone and periosteum.

2. That the osseous reproductive properties of the membrane vary according as it is taken from the long or short bones, being greater in the former than in the latter. (Ollier.)

3. That the normal form of the joint is better preserved than when the precaution of leaving the periosteal covering is taken.

4. That the subperiosteal resection involves less danger than when conducted on the old principle. This proposition is grounded chiefly on the results of Ollier's experiments on the lower animals; the number of unfavourable results which followed when the membrane was removed being much greater than when it was left.

5. That the difficulties attending the separation of the membrane in the dead subject are not to deter us from attempting the operation on the living, inasmuch as the membrane is less adherent in the latter, and also much thicker, in the great majority of cases, on the diseased than on the healthy bone.

6. That the chances of much shortening of the limb are diminished by this method, as shown by the results of the ankle-joint resections during the late Schleswig-Holstein war.

7. That subperiosteal resections are more conservative, as, in a great many cases, they diminish the necessity for amputation.

Gentlemen, accept my best thanks for your kind attention during the reading of this communication on a comparatively novel procedure in operative surgery, and of my experience in which I am sorry I can only give so brief an account. But still I trust that what I have said will convince some present of the importance and applicability of a practice which is based on accurate observation of physiological phenomena, on which alone all systems of rational therapeutics, whether medical or surgical, can truthfully be founded.

ILLUSTRATIONS OF THE VALUE OF THE MICROSCOPE IN THE TREATMENT OF THE STERILE CONDITION.*

By J. MARION SIMS, M.D., Paris.

THE great Velpeau died, having no faith in the practical utility of the microscope; and I am ashamed to say that, even ten years ago, I was one of those who believed it to be merely a scientific toy with which to while away leisure hours. Fortunately, my ignorance was dispelled, and I now look upon the microscope as essential to the daily duties of the physician. While I would refer you to Beale and others for what it has done and is doing in the broad domain of medicine, I shall content myself with laying before you now a few illustrations of the value of the microscope in the treatment of the sterile condition.

In the investigation of a case of sterility, there are three questions that must be settled at the outset, if we expect to proceed understandingly.

1. We must be sure that we have semen with spermatozoa.
2. We must ascertain if the sperm enter the utero-cervical canal.
3. We must determine whether the secretions of this canal are favourable or not to the vitality of the spermatozoa.

The present advanced state of physiological knowledge warrants us in saying that conception is impossible without spermatozoa; that it is impossible if the spermatozoa cannot pass into the cavity of the uterus; and that it is equally impossible if they die, or are poisoned, on their passage to this cavity. These three points we must determine, it matters not what other complications there may be. Fortunately, they are all easily and quickly settled by the microscope. Without the microscope, it is utterly impossible to determine either one of them. Without the microscope, then, our treatment of the sterile state is simply blind empiricism. With it, our diagnosis becomes absolutely certain, and our treatment at least rational. What, then, are the first steps in

this investigation? How can we begin? Where shall we begin? We only need a drop of seminal fluid to determine whether it contains spermatozoa or not.

I am in the habit of examining the vaginal and cervical mucus for spermatozoa some hours after coition, and I have thus often determined the three propositions above laid down all at once. Gosselin and other French physicians direct the man to use one of the abominable things so common in France (and, I am sorry to say, now also in my own country), which are intended to protect the male against infection, and the female against both infection and conception. This, with its contents, is sent to the microscopist for the necessary investigation. But a nicer way of managing this disagreeable but important affair is that adopted by my friend Mr. Curling. He directs the man, immediately after intercourse, to squeeze a drop of mucus from the urethra on the object-glass; and from this the examination is made. This will settle the question, so far as the man is concerned; but it will still be necessary to examine the utero-cervical secretions, to determine the second and third propositions; and therefore I never put the man through the above ordeal till I am forced to do so. Fortunately, it is not often necessary. Given a case of sterility for investigation, the physician examines the state of the uterus and its appendages. His patient may have a frightful dysmenorrhœa, a contracted or flexed cervix, a malposition, or something that would probably prevent the passage of the semen into the cavity of the uterus. He may feel perfectly convinced that his patient can never conceive, unless some surgical operation be performed—perhaps incision of the cervix uteri. This operation is usually done, to permit the easy passage of the semen into the uterine cavity. But how can we tell that the semen does not pass into the uterine cavity? We must not take it for granted that it does not, simply because the os uteri seems to be small; for we know that cases are on record where conception has occurred where the os would barely admit a small sized probe; and we know also that spermatozoa occasionally pass along the Fallopian tubes, which barely admit the passage of a bristle. If the semen enter the cervical canal, we may lay it down as a rule, that an incision of the cervix is not necessary; but, if it do not, it may be necessary. We may perform any operation we please for the relief of suffering, and for the restoration of health; but I insist that we have no right to perform any operation, or to institute any treatment whatever, solely for the cure of the sterile condition, till we have first settled the three propositions above laid down touching the presence and viability of the spermatozoa. To find out all at once, and with the least delay and trouble, I usually say to the husband (or wife, as it may be), "It is very important, before instituting any treatment, to be sure that the semen enters the canal of the neck of the womb, and also to be sure that the uterine secretions do not neutralise the semen; for, under such circumstances, there could be no conception. Will you, then, have the kindness to send your wife to me, or come with her, some day when you have had sexual intercourse in the morning?" In nineteen cases out of twenty, the wife presents herself the next day. The speculum is introduced, and some vaginal mucus is removed with the syringe, and placed on an object-glass. Then some cervical mucus is drawn out, and placed on another object-glass. These are then examined under the microscope. If we find spermatozoa, well and good; but, if we find none, our fears are at once aroused. What then is to be done? I simply say that I am not quite satisfied with the examination, and would like to see the wife again under the same circumstances. But suppose we find none on the second examination: then I have been in the habit of saying, "I would like to see your wife in bed twenty or thirty minutes after sexual intercourse, for I fear her vagina holds no semen." The husband appoints the time; and, of course, I am punctual. If I then find in the vagina a fluid with the seminal odour, I am satisfied with the microscopic examination. But it will not do to say we have no spermatozoa, simply because we find the vagina empty; for I have occasionally examined the vagina eight or ten minutes after coition, and found it empty, and with no sign of spermatozoa whatever in the little mucus that would naturally stick to its walls; and that, too, where a normal quantity of semen, full of the fructifying principle, had been ejaculated. But, unfortunately, the vagina had ejected the whole of it, as a consequence of some anatomical configuration. If we fail to get any for examination, we are then forced to Mr. Curling's expedient. Suppose, at last, we are satisfied that the semen contains spermatozoa, then, as before said, we have to determine whether these pass into the cervix uteri; and, if so, do they find a fluid there favourable to their existence alive? And all this can only be done by the use of the microscope.

The question of the entrance of the semen into the cervical canal and of the effect of its secretion upon the spermatozoa, can be fully and satisfactorily ascertained only during a very brief period. We may make a mistake if the microscopic examination be undertaken just before the

* Read in the Midwifery Section at the Annual Meeting of the British Medical Association in Oxford, August 1868.

expected return of the menses; and why? Because there is always a certain amount of fullness of the uterus—of engorgement, so to speak—which precedes the menstrual flow; and the cervical canal may not admit the semen from mere turgescence of its walls. Besides, at this time, its secretions are almost sure to kill the spermatozoa even if they should happen to enter this canal.

Physiologists are generally agreed that conception takes place during the week following menstruation. Avrard says we have fourteen days of active uterine life and fourteen days of uterine hypnosis. He says that conception can occur at any time after menstruation up to the fourteenth day, counting from its commencement. For instance, if menstruation should last for three days, then we would have eleven days for the possibility of conception. But, if menstruation should last eight or nine or ten days, then we would have respectively but six or five or four days as the time possible for conception. After this time, the uterus, according to Avrard, lapses into the hypnotic state, when conception is impossible. While I am disposed to accept Avrard's dictum as the rule, I think I have seen exceptions to it, if we can always depend upon testimony seemingly reliable. Be this as it may, I am sure of this fact: if we wish to determine the effects of the cervical mucus upon the spermatozoa, we must make the experiment during the week that follows menstruation. About the fifth or sixth day after the flow is the best moment; for then the uterus is in the most favourable condition. The cervical mucus, which just before menstruation was perhaps thick and opaque, then becomes clear and translucent. If, by examination made at this particular period, we should find spermatozoa in the cervical mucus, we could safely say that it will not be necessary to incise the neck of the uterus. But if the sperm do not enter the canal, then the probabilities are in favour of the necessity of some surgical interference. The semen may enter the cervix in great abundance, and we may find the spermatozoa all dead, even but a few minutes after coition. Then, of course, we must find out the source of the poisonous secretion, and remedy it; for a vitiated secretion shows some organic condition requiring a special treatment.

As before said, when I wish to examine the action of the cervical mucus upon the spermatozoa, I order sexual intercourse in the morning—the dorsal decubitus to be retained for an hour afterwards; and I expect a visit from my patient four or five or six hours after coition. Sometimes I find spermatozoa in great abundance in the cervical canal, and not one living. [I have occasionally examined the mucus, six, eight, and ten minutes after coition, and found all the spermatozoa dead.] Sometimes, we find half of them dead; again, only about a third; again, two-thirds. Sometimes, in one portion of the mucus, every spermatozoon is dead; while, in another portion of the same sample with fewer epithelial scales, we may find them alive and in great numbers. Now and then, after treatment for a month or more, I have found the mucus drawn from the lower segment of the cervical canal full of living spermatozoa, and I have supposed that the case was cured; but when I came to examine that drawn from the upper segment of the canal, near the os internum, they were nearly all dead. This was evidently because the mucous membrane lining the lower segment of the cervix, being more easily reached and more thoroughly treated, had assumed a healthier character, and consequently its secretion was restored to a normal condition; while that higher up, and more difficult to reach, had not been so much improved, and hence its secretion was still abnormal—a condition requiring further treatment.

The vaginal mucus, by its natural acidity, kills very quickly every spermatozoon. I do not now remember ever to have found one alive in the vagina, except when the examination was made very soon after coition; when, indeed, the vagina was full of semen but slightly mixed with other secretions. Examined three or four hours after intercourse, the spermatozoa found in that portion of the mucus of the vagina adhering to its walls are always all dead. Indeed, the normal vaginal secretion seems to be a perfect poison for the superabundant spermatozoa.

When, after a month's treatment, I wish to know whether the case is cured or not—in other words, whether all possible recognised barriers to conception are removed—I order sexual intercourse (just after menstruation) at night, and examine the cervical mucus twelve or fourteen hours afterwards. If the majority of the spermatozoa be alive and active, I have great hope of conception. Before dismissing a case as cured, I think it necessary to examine the mucus thirty-six hours after coition; and, if it is then all right, of course I suspend the treatment, and patiently await the hoped-for result.

So much care is necessary in the removal of the mucus for microscopic examination, that I may be pardoned for referring to it again. The patient is placed in the left semi-prone position, as I have elsewhere so minutely described, and my speculum is introduced, and some of the vaginal mucus is drawn up with a small glass syringe, previously washed out with warm water. This is deposited on the object-glass; the vagina

is then cleared of all secretion, whether vaginal or cervical, the whole of the vagina and the os uteri being thoroughly wiped over with a pledget of cotton. This is for the purpose of guarding against the possibility of mixing vaginal with cervical mucus, which would, of course, spoil the whole experiment. The cervix is brought forward either by the depressor or a tenaculum, if necessary, which enables us to look directly into the cavity of the cervical canal. The syringe is then to be again thoroughly rinsed in warm water; its nozzle is passed into the gaping canal for half an inch, and the cervical mucus in its lower segment is drawn out. The instrument is emptied, washed out again with warm water, and re-introduced up to the os internum, and another portion of mucus is drawn out, provided there is any left after the first effort. Thus we have three specimens of mucus; *i. e.*, the vaginal, and two from the upper and lower portions of the cervix unmixed with the vaginal. The cervical secretion should be clear and translucent, and about the consistence of the white of egg. If it contain any little opaque specks of milky whiteness, it invariably poisons the spermatozoa, to a greater or lesser extent. But the portions of it without the milky-looking specks may be very favourable to spermatozoic life. We sometimes find the cervical mucus perfectly clear, and yet poisonous to the spermatozoa. Here we would naturally expect to find excessive alkalinity of the secretion; but I have not been able to detect it. In these cases, it has seemed to me that the spermatozoa were killed—drowned, as it were—by the very abundance of the secretion. I do not here allude to cases of uterine catarrh, where the secretion is very thick and albumino-purulent; for, of course, this is a deadly poison to the living principle of the semen. But I allude wholly to such cases as have been changed by treatment to a condition giving rise to a secretion seemingly normal, so far as an ordinary ocular examination is concerned. Here the microscope is our unerring guide. The mucus may be clear, and perfectly normal in appearance; but, if it kill the spermatozoa, then there is still some point in the canal of the cervix, or in the cavity of the uterus, that gives out a vitiated secretion; and this must be found out and corrected before the case is wholly cured. When we find living, active spermatozoa high up in the cervical canal thirty-six or forty hours after coition, we can pronounce the case cured, so far as it can be by surgical means, and not till then.

It is time for us to pause and consider if there is not something more to be done for the sterile condition, than to split up the cervix uteri. I look upon this operation as one of the greatest importance, as one of the most valuable in uterine surgery, but I think that we have followed too blindly the example and teachings of its illustrious author, Sir James Y. Simpson. For myself, I am now sure that I have cut open the cervix uteri, perhaps scores of times, when it was both useless and unnecessary; and I know that others have done the same thing. Do not misunderstand me. I speak here solely of the operation with reference to the sterile condition, when it would be wholly useless if the husband happened to be sterile, and certainly unjustifiable unless imperatively called for by considerations of health. Incision of the cervix for dysmenorrhœa is one thing; incision of the cervix for sterility, even if there be dysmenorrhœa, is another, and it behoves us to draw the line of distinction in every case, and not to take it for granted that every woman is sterile who may have dysmenorrhœa or feeble health, or that every man is prolific who may be vigorous and enjoy good health. I am sorry to say that I have had the misfortune to incise the cervix in half a dozen cases of sterility, where I found afterwards, to my great mortification, that the husbands were incapable of procreation, because their semen had no spermatozoa, and that too since I have known the value of the microscope. In each case the operation was called for to restore health, but was totally useless for the relief of its incidental accompaniment, sterility, and would not probably have been submitted to for considerations of health alone, had it not been for the hope of offspring afterwards. I made the mistake of operating on these cases, because the social position, moral character, and appearance of health in the husband, conjoined with the excessive dysmenorrhœa and utter prostration of the wife, led me to operate without the preliminary step of ascertaining whether there were spermatozoa or not. I wish others to profit by my mistakes; and I am less ashamed to tell you of them, than I am to own them to myself. However, this can never happen to me again, and should not, after this warning, happen to any of my brethren. I know many men who have no spermatozoa and cannot therefore become fathers. They are all strong active men in the prime of life, and all perform the sexual function with vigor and at will. The very fact of their natural vigor and strong passions had been their ruin, for most of them had contracted urethritis during their early and unmarried life, and had suffered from its unlucky sequence epididymitis. This condition is said to be sometimes remedied, but I have not as yet seen a case of it cured.

[To be concluded.]