religion, that the sanction of government support should not be given to doctrines the removal of which, supposing them ultimately to be found erroneous, would have political as well as intellectual obstacles to contend with. And as, in the present vexed state of all economical questions, I should regret to see the sanction of the Privy Council given to any elementary treatise containing, for instance, an attack or defence of trade-unions, so, in such a sanitary catechism as I am speaking of, it would obviously be improper to put forward crude assertions on any such doubtful question as that, for instance, of the use of alcoholic drinks. But it would be very easy to pick out a dozen elementary propositions with respect to air and water and the human body, about which no one could raise any dispute, and the practical inferences from which would be obvious and certain. Any child can be made to breathe into lime-water, and can see the difference produced by the air which comes from his lungs and the air which comes from a bellows. Any child can be shown the effect of carbonic acid on a candle, and will readily believe, half rationally, half on trust, that what is incompatible with light is incompatible with life also. Elementary dogmas connected with ventilation will follow in due course; the importance of windows opening at the top; of a free vent through the chimney; of sufficient space in the sitting-room and bedroom; of a through current from front to back, etc.—would naturally follow. Water might be dealt with in the same way; and perhaps some simple facts connected with the skin and the lungs might be added.

But, I will again repeat, the essential matter is neither that the knowledge thus given should be complete, nor that it should be fully and rationally apprehended. The object is to call the universal attention of the ratepaying public to health-questions from the earliest period of life; to implant, if I may be permitted the word, an universal prejudice in favour of health; to place it, in fact, upon the same footing, in the ratepaying mind, as reading, writing, and arithmetic. Anything short of this will not meet the necessities of the case. Desultory popular lectures are useful in their measure; but they are attended, of course, by the more thoughtful of our working classes—that is, by those to whom they are least necessary. What is essential is to touch the unthinking mass; and this can only be done by penetrating them from childhood with an organic instinctive dislike of poisoned air and water, of dirty skins and overcrowded rooms. Thus, and thus only, will you create a public to whom the application of such sanitary measures as may perhaps always be necessary in our complicated civilisation will not be distasteful, will be met halfway, and will, therefore, be tenfold more effectual.

ON THE ADMINISTRATION OF NITROUS OXIDE.*

By J. T. CLOVER, F.R.C.S.Lond.

THE inhalation of nitrous oxide as the means of preventing pain in surgical operations was suggested by Sir Humphry Davy at the close of the last century. We have, however, no record of any serious attempt to introduce the practice. Probably the gas was experimented with, but abandoned from its not uniformly producing anaesthesia. The apparatus at that time would fail to produce satisfactory results, if they were made use of by the most skilful administrator. In 1844, Dr. Colton successfully administered nitrous oxide to Horace Wells, an American dentist, who then had a tooth extracted without feeling pain. Wells tried to introduce the practice of using it in dentistry; but, after the adoption of ether inhalation, by Morton and Jackson, it was rarely used till about five years ago, when Dr. Colton, having improved the means of storing and using it, successfully revived the subject. When lately in London, he said he had records of twenty-eight thousand cases of inhalation without a single fatal result, and the only administration of nitrous oxide, he had heard of, which was followed by death shortly after, was one of extensive phthisis; he had never known a person made so ill by it as to need a carriage to ride home.

Mr. Rymer, a London dentist, published some experiments with the gas, in 1864, but they were not so successful as to induce any one to adopt the practice of giving it. In March last, Dr. Evans, of Paris, assisted by his nephew, administered nitrous oxide to some patients at the Dental Hospital, London, in the presence of several medical men who were invited to witness its effect. He brought with him a large India-rubber bag and an India-rubber tube (without any spiral wire inside it). The tube was connected with a wooden pipe supplied with valves for preventing the re-inhalation of the gas. Dr. Evans obliged the patient to breathe through the pipe by pressing his lips against it, whilst an assistant closed the patient's nostrils. Most of the cases were

so successful as to render it undeniable that the gas, when properly used, afforded complete immunity from pain, and enabled the patients to walk away without any unpleasant effect in a few minutes. But some of the patients struggled a good deal before they were put to sleep, and some appeared to feel the pain during the whole time of the operation. These partial failures were the results either of the supply-tube having collapsed, or from air having been sucked in accidentally. The following modification of an instrument which I have used for the last seven years, for giving four per cent. of the vapour of chloroform, will be found to obviate the objections to the plan previously adopted. The bag is filled with nitrous oxide, and I hang it at my side by means of a loop of ribbon; a non-collapsible tube leads from it to a face-piece which is made of sheet lead, so as to be easily moulded to the face, and edged with India-rubber tubing, so that nose and mouth may be covered with an air-tight cap; two valves prevent the gas from being breathed a second time. This plan supersedes the necessity for pinching the nose; but when a patient breathes in a rapid and forcible manner, it does not always prevent a little air from being drawn in through the beard, or by the cheek falling in, during inspiration. To obviate this I have added a supplemental bag, holding about two hundred inches and connected

The woodcut represents the face-piece applied, with the supplemental bag, M, attached. A. Soft pad of India-rubber. C. The valve opening during expiration. D. The mount upon which the stop-cock, F G, fits; the spindle part of the inspiring-valve is seen projecting. E. Opening in the stop-cock of inhaling-tube, which permits air to be breathed whilst the face-piece is adjusted. When the little knob is turned down to G, the opening, E, is closed, and the supply-tube, H, leading from the bag of gas, is opened. M. The supplemental bag to be used or not by turning the stop-cock, L.

by a three-quarter inch tube and stop-cock with the face-piece; the use of this, in preventing the access of air, and in saving gas, will be seen presently. As patients are apt to wake too soon, I have sometimes prolonged the anaesthesia whilst the mouth is still open, by giving the gas through the nose-cap which I have been in the habit of using with chloroform. Most of the patients to whom I have given it have had no

* Read in the Medical Section at the Annual Meeting of the British Medical Association in Oxford, August 1868.

preparation, but I would not advise it to be taken just after a full meal; and the dress of the patient ought not to be very tight. It is well to warn the patient that a ringing noise and general sense of pulsation may be experienced, and that he should not make the fitting of the mouth-piece difficult by attempting to talk. When teeth are to be removed, a gag should be put between the jaws, so that there may be no loss of time in getting the mouth open. Whilst fitting the mouth-piece, the patient should breathe air only, and the gas should not be turned on till the patient breathes steadily. As long as he breathes calmly the supplemental bag should be empty; but when he begins to *pant*, the stop-cock should be opened. The gas received into this bag is so readily yielded during inspiration, that there is not any air sucked in under the face piece. When the breathing has become so calm as scarcely to raise the expiring valve, the supplemental bag should be compressed every fifth or sixth respiration and allowed to refill with fresh gas; by this means the gas will be sufficiently free from carbonic acid to allow its elimination from the lungs. Mr. Coleman has invented an ingenious apparatus for absorbing the carbonic acid of the expired gas by means of hydrate of lime. I have had no reason to think that any harm results from the small amount of carbonic acid breathed in using my apparatus; but Mr. Coleman has certainly obviated one great objection to the use of a simple bag, in and out of which the patient is made to breathe without any exit for the respired gas. After inhaling twenty or thirty seconds, the patient begins to assume a livid appearance, this must not be taken as a sign of anaesthesia. I have often breathed the gas before a looking glass, and observed a decided lividity in my lips, whilst my sense of touch was perfect. In about fifty seconds the expression of the patient suddenly alters, his eye becomes unsteady and his hands slightly convulsed; I usually continue the inhalation ten or fifteen seconds after this, unless the respiration become very slow or stops, or unless the pulse become very quick or unsteady. The sudden dilatation of the pupil is another sign which should cause us to remove the gas. If general muscular rigidity have not come on during the inhalation, it often does so when the operation begins, and then the breathing may cease for ten or fifteen seconds. This should excite no alarm, if the pulse can be distinctly felt; the free inspiration which will follow will quickly restore a florid colour to the lips if the heart act efficiently; and, as has been stated, the inhalation should be immediately discontinued if the pulse threaten to stop. The state of complete insensibility does not usually remain more than half a minute after the removal of the mouth-piece, provided the patient breathe regularly; but there is a period after this when, although there is a sensation of something being done, it is not felt as pain. I had a tooth extracted a few weeks ago after inhaling the gas. I had no knowledge of the first application of the forceps, but before the tooth was extracted I felt that it was wriggled about, yet with no more sensation of pain than as if I had held the tooth with my fingers. Just before recovery, men and women alike often manifest hysterical symptoms. The will for a short time seems very weak, but I have never seen a prolonged fit of hysteria, as I have after chloroform; these symptoms always disappear in two or three minutes. Since the middle of last April I have recorded the cases in which I have given the gas, and they amount to 384. In all these cases there has been produced an amount of lividity, which in other circumstances I should have thought dangerous to life; in the majority, convulsive twitchings of the hands have occurred, in a few, dilatation of the pupil. In several, the pulse has become quick and weak, but never imperceptible at the wrist, and the recovery has been, in all cases, rapid and complete, so that the patient required no assistance in walking away. I am aware of only three cases of sickness where vomiting occurred within a few hours of taking the gas, and it was even then attributable to other causes.

My cases have been chiefly those of teeth extraction; two patients have had as many as seven teeth painlessly removed in the sleep of one inhalation; but, when more than three teeth have required extraction, I have generally noticed signs of returning consciousness, and allowed the patient to awake and rinse his mouth before repeating the inhalation; this is sometimes important, on account of the accumulation of blood in the pharynx which would cause a difficulty to the passage of air, should any necessity arise for artificial respiration. Patients seldom shew any reluctance to begin inhaling a second time. A patient of Mr. Rogers had ten teeth removed in one day and, a few days afterwards sixteen more were removed painlessly in three inhalations. The lady laughed hysterically every time on awaking, but only for a minute or two. I have applied the gas for several cases of iridectomy; in some of these the convulsive movement of the eye-ball delayed the operation, but it was completed without pain. I have also used it in operating for strabismus, and for wrenching an ankylosed knee. It is well suited for reducing dislocations, for removing the toe-nail, and opening fistulae, boils, and abscesses of all kinds. Two of my patients were subjects of epilepsy, but the gas did not affect them worse than others.

Whether nitrous oxide will ultimately displace chloroform and similar bodies, in operations which may last ten minutes, and leave much smarting, is improbable. The rapidity of recovery and absence of unpleasant consequences contrast favourably with chloroform; but when it is important to keep a patient quiet for as long as ten or fifteen minutes, chloroform is the best agent with which I am acquainted. I have never had a fatal accident from the administration of chloroform or of any other anaesthetic; and I think, if the air contain no more than 4 per cent of chloroform vapour, that a subject who had been allowed to inhale this till his breathing ceased would be restored by artificial respiration. In my experiments with nitrous oxide on dogs, cats, and rabbits, I have always found that respiration stopped before the circulation, and that, if the gas were removed when the animal was apparently dying, and at its last gasp, or if air were admitted by artificial respiration, it always recovered. If dangerous symptoms should arise from the inhalation of nitrous oxide gas being continued too long, artificial respiration would be the proper remedy. The chest and abdomen should be forcibly compressed six or eight times, at intervals of a second, whilst the patient is sitting or lying; if this produce a florid colour in the face, it will suffice, if not, the patient should be placed horizontally, and Silvester's method resorted to. It is to be regretted that the mode of making pure nitrous oxide on a small scale is difficult. Mr. Fox, the dentist, who has obligingly supplied me with gas, has exhibited to the members of the Association the apparatus which he uses. The gas is sold in a condensed form, by Mr. Barth, of Duke Street, Bloomsbury.

ILLUSTRATIONS OF THE VALUE OF THE MICROSCOPE IN THE TREATMENT OF THE STERILE CONDITION.*

By J. MARION SIMS, M.D., Paris.

To illustrate the necessity of the microscope in this department of surgery, I shall append a few cases, drawn up as succinctly as possible.

No. 1 had consulted two of the most eminent physicians in England and remained under the care of one of them for many weeks. She said that during the time the neck of the uterus was repeatedly cauterised. She got impatient and went to another physician who told her that the caustic treatment that she had submitted to was worse than useless; and that a surgical operation was the only thing to be done. She consented to it and he incised the cervix bilaterally. She did not conceive, and two years afterwards came to Paris to see me. I found the uterus normal in all its relations, the os tincæ and cervical canal sufficiently patulous. I explained to both husband and wife the importance of examining the cervical mucus four or five hours after coition. They returned the next day; the cervical mucus contained spermatozoa; therefore there was no necessity for any further surgical operation. But the spermatozoa were all dead; therefore there was a necessity for a treatment to rectify the vitiated cervical secretion. She remained in Paris a few weeks under my care, was cured and became a mother in nine months after her dismissal. Now, if the first physician had used the microscope, as I direct, he would probably have found that the semen never entered the cervix at all; and, if the second one had done the same thing, he would certainly have found that the mucus of the cervix poisoned the spermatozoa.

No. 2, a lady, in the highest ranks of life, was sterile. The cervix uteri was incised bilaterally. She had pelvic cellulitis afterwards. Two years after this I saw her, she was still childless. The microscope showed that the cervical mucus, examined four hours after coition, killed all the spermatozoa. While this condition exists conception is impossible.

No. 3, sterile, was treated for sterility in America for a long time (two or three years). She came to Europe; had the cervix cut open, and was sent away with the promise of offspring. I saw her some time afterwards. The microscope proved that the husband was sterile. Therefore, the previous treatment in America and the operation in Europe were useless. I could relate several other cases like the above. But, as I have often made the same mistake before I fully understood the value of the microscope, I compassionately forbear.

No. 4, married four years; sterile. She had dreadful dysmenorrhœa, followed by discharge of a bloody brownish mucus of an offensive odour. The uterus was ante flexed; anterior wall hypertrophied; os uteri small. I was in doubt, at first, whether to recommend an incision of the cervix or not. The husband was a distinguished physician. I told him that an operation would be necessary if the semen did not enter the canal of

* Concluded from page 466 of last number.