halves separated from one another by a space of about 3 inches in their whole length. This gap allows the spine to rest on the split mattress without pressure upon the spinous processes of the vertebrae. In the middle of the bed the gap between the mattresses consists of a hole 8 inches in diameter, into which the special bedpan fits. This hole for the bedpan is suitably protected by rubber sheeting to prevent soiling the mattresses. The bedpan (A) is held by a lever which works in such a way that when pushed into place under the patient's buttocks the bedpan rises close under the hole in the mattress. When the pan has to be emptied the lever is pulled into position, shown in the illustration, and it drops so as to clear the bed frame. The case in which the patient lies is large enough to take a Thomas spinal frame or a hip abduction frame in a moderate degree of abduction. If this is used the lower mattresses would be discarded.

To the sides of the cage are hinged two flaps, on to which are strapped half-mattresses (B). Except when the patient is to be turned over these flaps lie as shown, flat against the sides of the bed cage. When the patient has to be turned over, the pillow is removed from under the head, and one pillow placed in front of and at the sides of his body, so as to pack him tightly in the cage. The flaps are then turned over so as to cover in the front of the patient, and are secured in this position by a catch and screw at the head, and another at the foot of the cage (C in Fig. 3). The front mattresses of the cage, when in position in front of the patient, have a 4-inch gap between them, which gives ample space for the face (D in Fig. 2).

The patient, being now snugly encased between the two sets of mattresses and the side pillows, can be turned over. This is done by releasing the catch (C) at both ends of the bed frame, and Fig. 2 shows the patient in the cage being turned over. When the cage has been reversed by being turned through a half-circle it is fixed in this position by the catch, and the back halves of the cage are open. In Fig. 3 one half (E) is seen being lifted up, and the other (F) has been dropped down by the side of the cage. The patient's back is now fully exposed for the purposes of dressing or attention to the skin of the buttocks.

The cases for which this device has proved to be of the greatest service have been: fractures of the spine, with or without paraplegia and paralysis of the bladder; tuberculous disease of the spine, especially cases in which a bone graft operation has been done; spinal cord tumour with paraplegia, or after a laminectomy; bad cases of fracture of the pelvis; special cases of fracture of both femora; and bad cases of burns involving the back.

The bed has been made for me by Messrs. Nesbitt-Evans and Co. of Wednesbury (Staffs), to whom I am greatly indebted for their trouble in carrying out my design.

THE RELATION OF ORTHODOX RHEUMATIC INFECTION TO MULTIPLE INFECTIVE ARTHRITIS.

BY

VINCENT COATES, M.C., M.A., M.D.CANTAB.,
M.R.C.P.,

PHYSICIAN, ROYAL MINERAL WATER HOSPITAL, BATH; CARDIAC SPECIALIST,
WILTS COUNTY COUNCIL SCHOOL MEDICAL SERVICE.

It is manifestly impossible from a statistical point of view to assess the relation, if any, between the acute and the more chronic forms of rheumatic disorder unless such terms as "acute rheumatism" and "chronic rheumatism" have a standard significance. As, however, the term "acute rheumatism" in the generally accepted sense means rheumatic fever, and the term "chronic rheumatism" is indiscriminately applied to many dissimilar affections of both muscle and joint, it is proposed in this paper to deal only with orthodox rheumatic infection and with multiple infective arthritis. By the former is meant frank rheumatic fever, and by the latter that type of subacute or chronic arthritis which is non-suppurative and is generally conceived to be due to infection of a non-specific character—in other words, which is not gonorrhoeal, pneumococcal, tuberculous, or syphilitic in nature. "Rheumatoid arthritis," as opposed to osteo-arthritis of

the hypertrophic type, and "atrophic arthritis" are other terms for the disorder in question. Any arthritis which can be classified as being due to, or associated with, disordered metabolism, such as climacteric arthritis, is thus excluded from the series under discussion.

The possible relation of orthodox rheumatic infection to multiple infective arthritis was investigated by the following methods.

1. The evidence afforded by the family history of 50 consecutive cases of multiple infective arthritis in regard to orthodox rheumatic infection.

In 16 instances rheumatic fever was known to have occurred in a parent, brother, or sister.

2. The incidence of a clear history of orthodox rheumatic infection in 300 consecutive cases of multiple infective arthritis. By a clear history is meant the patient's positive assertion that rheumatic fever had occurred earlier in life and was not immediately connected with the present disability. This evidence was frequently borne out by the individual's own private practitioner.

Rheumatic fever is believed to have occurred in 14 instances, without, however, the presence of discernible complications of a cardinal nature (for example, carditis or large subcutaneous nodules) at the time of examination. This series does not include 13 additional cases in which rheumatic fever was alleged by the medical practitioner in charge of the case to have immediately preceded the onset of infective arthritis.

3. The occurrence in the aforesaid 300 cases of cardinal signs of orthodox rheumatic infection—namely, cardiac lesions of the orthodox rheumatic type and large subcutaneous nodules of the same histology^{1,2} as those which occur in rheumatic fever of childhood. The millet seed granules³ described by me as occurring in subacute rheumatic infection of childhood and in multiple infective arthritis are not included in this category.

The findings of the cardinal signs of orthodox rheumatic infection in these 300 cases of infective arthritis can be tabulated as follows.

	No Rheumatic Fever.		Rheumatic Fever.	
	No Nodules.	No Nodules.	No Nodules.	Nodules.
Mitral stenosis	3	2	—	—
Mitral regurgitation	2	—	1	2
Aortic regurgitation (in young subjects with negative W.R.)	2	—	—	—
Nodules	No carditis ... 13		No carditis ... 1	

Conclusions from the Above Data.

1. The incidence of orthodox rheumatic infection in the immediate family history of 50 consecutive cases of multiple infective arthritis is 32 per cent. This is a very high percentage, and accords with the popular conception that "rheumatism" has a familial or hereditary basis.

2. A low estimate of orthodox rheumatic infection in 300 consecutive cases of multiple infective arthritis is nearly 5 per cent. The attack rate of orthodox rheumatic infection among the under 15 population in Norway⁴ is given as 0.8 per 1,000, and the highest figures published are from the Royal Naval School at Greenwich⁴—namely, 6 per 1,000. The attack rate was 2 per 1,000 in the 16-24 age period of the Ministry of Health's inquiry⁴ into "rheumatism" in insured adults of both sexes.

3. The cardinal signs of orthodox rheumatic infection—namely, carditis and/or subcutaneous nodules—were present in 300 consecutive cases of infective arthritis in the following percentages:

	Per cent.
Cardiac lesions only without a history of rheumatic fever	2.33
Subcutaneous lesions only without a history of rheumatic fever	4.33
Cardiac lesions in the whole series	4.00
Subcutaneous lesions in the whole series	6.00

These cardiac figures are much higher than those given for organic cardiac disease after routine examination of school children⁴ in England and Wales, 0.7 per cent.; in Scotland, 0.52 per cent.; and in New York,⁵ 1 per cent.

In regard to subcutaneous nodules, Carey Coombs⁶ has stated that of the principal phenomena of orthodox rheumatic infection—namely, carditis, polyarthritis, the

subcutaneous node, and chorea—the subcutaneous node is the most “rheumatic” of all. The above figures are therefore, to say the least of it, significant.

REFERENCES.

- ¹ Vincent Coates: *Med. Journ. and Record*, October, 1926.
- ² Vincent Coates and Carey Coombs: *Arch. of Dis. in Childhood*, August, 1926.
- ³ Vincent Coates: *Proceedings of Rheumatic Conference at Bath*, May, 1928.
- ⁴ Quoted by Ministry of Health's Report No. 44, 1927.
- ⁵ R. H. Halsey: *Journ. Amer. Med. Assoc.*, April, 1923.
- ⁶ Carey Coombs: *Rheumatic Heart Disease*, 1924, p. 6.

SOME EFFECTS OF WARM IMMERSION BATHS UPON THE CIRCULATION.*

BY

GEOFFREY HOLMES, M.B.CANTAB.,
PHYSICIAN, HARROGATE ROYAL BATH HOSPITAL.

SPA physicians are aware that some of their professional brethren are a little sceptical about the efficacy of bath treatments, and it is probably not generally realized what profound effects such a simple procedure as a warm immersion bath may exert on the bodily functions, and especially upon the circulation. Some of these effects can be quite accurately measured. First, and most obvious, is an alteration in the pulse rate. Secondly, variations in blood pressure, both systolic and diastolic. Thirdly, and beyond the scope of ordinary clinical observations, there is a change in the amount of blood flowing from the heart per minute, as well as in the stroke volume or output per beat. Fourthly come various less accurately measurable effects, such as sweating, cutaneous hyperaemia, etc.

The following observations were made during a series of experimental baths taken by three normal subjects with a view to ascertaining the effect of immersion baths on the circulation and to determine whether the consequent alterations in pulse rate and pulse pressure were in any close degree correlated with alterations in the heart's output. The effects on the circulation rate will be the subject of a separate paper, but I wish to draw attention to some rather striking effects on pulse rate and blood pressure revealed during these experiments. Typical examples are here shown in tabular form.

It is common knowledge that an increase in the body temperature is accompanied by a proportional increase in pulse rate. Our experiments show this effect, but, in addition, two other features appear. (1) When a normal subject is placed in a bath at a temperature of approximately 103° F. there is an immediate increase in the pulse rate long before any significant rise of body temperature occurs. This increase is so sudden that it would appear to be due to some reflex cause, and a subsequent greater rise is superimposed. (2) In certain of the experiments there was definite hyperpnoea and consequent alkalosis (excessive CO₂ elimination). This alkalosis was associated with still further increase in the pulse rate.

It is just thirty years since Edgcombe and Bain¹ published papers on the effect of baths, massage, and exercise on the blood pressure, and these papers are still regarded as classics. They contain no reference to diastolic arterial pressure, but they were written before present-day methods of measuring diastolic blood pressure had been devised. The writers do, however, state the effects on mean arterial pressure, which they define as the reading taken when the indicator of the haemodynamometer gives its maximum excursion. It is probable that this point of maximum excursion would approximate somewhat closely to the diastolic pressure.

Our own observations show that in normal persons entering a bath at 103° F. there is an immediate fall in both systolic and diastolic blood pressure, approximately 10 mm. of mercury in each case. This initial fall would appear to be due to simple hydrostatic causes. When the blood pressure is taken with the subject on a couch the arm is lying by the side. If the arm be elevated to an extent

such as occurs when taking the blood pressure of a patient immersed in a bath, a similar fall is observed. Subsequent to these immediate changes the systolic pressure may show a slight rise, which in none of our experiments exceeded 16 mm. of mercury, but was usually rather less. The most striking feature with regard to blood pressure was a marked fall in diastolic pressure, together with an alteration from the normal characters of the Korotkow sounds—that is, the sounds heard in the artery during decompression when determining blood pressure by the auscultatory method.

Wiggers² describes five phases in the Korotkow sounds:

“*First phase*: The sudden appearance of a clear sound, lasting for a fall of approximately 14 mm. of mercury. This has already been described as a criterion of systolic pressure.

“*Second phase*: The acquisition of a murmurish character, lasting while the pressure falls approximately 20 mm. of mercury more. It is attributed to the stenosis of the vessel produced under the cuff at this stage.

“*Third phase*: The replacement of the murmur by a sound becoming progressively louder and lasting during the next 25 mm. of pressure fall.

“*Fourth phase*: The muffling of the sounds, lasting while the pressure falls 5 to 6 mm. more. The cause of this muffling has been variously interpreted as due: (a) to the absence of diastolic collapse; (b) to the slowing of the stream under the cuff; (c) to the decreased resonance in the deflated cuff; (d) to a lack of flattening of the arterial wall.

“*Fifth phase*: The disappearance of all sound.”

The pressure in the cuff at the commencement of the fourth phase is generally accepted as being the diastolic pressure. We found that although normally at rest the transition from the third to the fourth phase was well defined, and gave a sharp reading of diastolic pressure, after a short time in a bath at 103° F. this was no longer so. Almost immediately the sounds during the third phase, instead of becoming progressively louder throughout the whole of the phase, became at first louder and then gradually softer, and during this period of diminuendo tailed off almost imperceptibly into the fourth phase. Hence it was not possible to recognize the usual clearly defined criterion of diastolic pressure. In addition, the fourth phase became prolonged, and, not infrequently, after the subject had been in the bath for thirty minutes or longer, the fourth phase was found to continue right down to zero pressure—that is to say, the fifth phase was absent altogether.

In column three of the table the left-hand figure is, so far as could be ascertained, the transition from third to fourth phase. The right-hand figure, where only two are given, is the pressure at the end of the fourth phase—that is, disappearance of all sounds. Where this figure is zero it indicates that sounds were still present in the artery when the sphygmomanometer cuff was completely deflated. Where three figures are given, the front two show pressures at which clearly defined softening or muffling of the sounds occurred. So far we are unable to interpret these points with reference to Wiggers's phases, but it seems at least possible that in the later stages of a hot immersion bath the diastolic pressure falls very near to zero. These changes—namely, the fall in diastolic pressure and alteration of the Korotkow sounds—closely resemble those observed in severe cases of Graves's disease (hyperplastic exophthalmic goitre), and possibly also in cases of aortic incompetence and severe anaemias.

The significance of some of these observations is as yet obscure, but one point of interest may be briefly mentioned. It has generally been considered that an increase in pulse pressure associated with an increase in the pulse rate is indicative of an increase in the circulation rate. This has been shown to occur in exophthalmic goitre. Our observations on the circulation rate in baths have shown that, as a rule, no significant increase in circulation rate occurs even though both pulse rate and pulse pressure are increased. Indeed, in certain experiments the circulation rate under these conditions has been found to be actually lessened.

In the present communication it is impossible to give all the evidence in detail, but the most reasonable interpretation would be that there is constriction of the arteries and arterioles, together with a dilatation of the capillaries. This dilatation gives rise to diminished peripheral resistance, as well as a certain amount of peripheral stasis.

* A preliminary communication summarizing an address to the Yorkshire Branch of the British Medical Association at Harrogate on October 19th, 1929, and based on observations made in collaboration with Professor Whitridge Davies and a number of the senior students from the Leeds Medical School. The work was carried out at the Royal Baths and the Royal Bath Hospital, Harrogate.