We have, then, our mental picture of the structure of the erythrocyte demonstrated—the condensation of lipin molecules on the surface which acts as a limiting membrane enclosing a spongy stroma in the interstices of which haemoglobin is contained.

CONCLUSIONS.

1. The normoblast nucleus disappears by karyolysis or by pycnosis and karyolysis, with or without karyorrhexis, and when the erythrocyte enters the blood stream no trace of the nucleus remains.

2. The erythrocyte does not possess a histological cell membrane, but has on its surface a molecular condensation of lipins which acts as an envelope. Whether or not this is a bimolecular layer, or what physico-chemical relationship between the lipin covering and the protein of the stroma exists, are fields for further investigations.

3. The basophilic conditions of the erythrocyte—punctate basophilia, reticulation, and diffuse polychromasia—are expressions of the same process and vary only in degree. The staining substance is not chromatinic.

4. The presence of these conditions, when obtained by the usual staining methods, would appear to indicate a defect in the lipins on the surface of the erythrocyte.

5. During the course of the work it was noticed that not all blood films behaved in the same manner. Accurate standardization of the reagents and the time, or the employment of other methods on similar lines, might lead to the elucidation of certain problems in connexion with the genesis and life-history of the erythrocyte, and also with its behaviour in disease.

The photomicrographs are by W. E. Cooke and C. F. Hill.

BIBLIOGRAPHY.

- Bütschli, O.: *Protoplasm and Microscopic Foams*, English Trans. E. A. Minchin, 1894.
 Cooke, W. E.: *British Medical Journal*, 1928, ii, 790.
 Idem: *Le Sang*, 1929, iii, 10.
 Idem: *Amer. Journ. Med. Sci.*, 1929, 4, clxxvii, 537.
 Ehrlich, P.: *Berl. klin. Woch.*, 1881, xvii, 43.
 Idem: *Charité-Annalen*, 1885, x, 136.
 Görter, E., and Grendel, F.: *Journ. Exper. Med.*, 1925, 41, 439.
 Gruner, O.: *Biology of the Blood Cells*, 1913, 75.
 Hardy, W. B.: *Journ. of Physiol.*, 1899, xxiv.
 Hawes, J. B.: *Boston Med. and Surg. Journ.*, 1909, clci, 493.
 Key, J. A.: *Arch. Int. Med.*, 1921, xxviii, 511.
 Idem: *Amer. Journ. of Physiol.*, 1924, lxx, 86.
 von Knauff-Lenz, E.: *Arch. ges. Physiol.*, 1918, 171, 51.
 Langmuir, J.: *Journ. Amer. Chem. Soc.*, 1917, xxix, 1848.
 Pockels, A.: *Nature*, 1891, 43, 437.
 Rayleigh, Lord: *Phil. Mag.*, 1899, 48, 331.
 Rohonyi, H.: *Kolloid-Chemie*, Beihefte, 1916, 8, 337, 391.
 Walker, C. E.: *Proc. Roy. Soc.*, 1928, B, 103, 397.
 Walker, C. E., and Allen, M.: *Proc. Roy. Soc.*, 1927, B, 101, 468.

TYPES OF ACUTE MASTOID INFECTION.*

BY

A. J. WRIGHT, F.R.C.S.,

SURGEON TO THE EAR, NOSE, AND THROAT DEPARTMENT, BRISTOL GENERAL HOSPITAL.

WHILE, in what may be described as the classical textbook cases of acute mastoiditis, the diagnosis is easy, experience has taught me that atypical cases are frequent and that, in these, diagnosis is often extremely difficult. It is probable that in all cases of acute middle-ear inflammation the lining of the mastoid antrum and cells participates in the inflammatory process. It is only, however, in those cases in which the inflammation spreads to the bone, owing either to defective drainage or to the intensity of the infection, that the term "mastoiditis" should be properly applied. The early recognition of such a condition is of the utmost importance, both from the point of view of preservation of the hearing and also of the general health.

As an aid to diagnosis, I have tried to classify cases into groups, but, in dealing with them in this way, I wish it to be clearly understood that cases are not infrequently met with showing characteristics of two or more groups in combination.

The Classical Type.

Dealing first with the classical type, there is present a history of recent earache with discharge, and tenderness and swelling over the mastoid process. Fever and constitu-

tional disturbance are present. Characteristically, the discharge has either suddenly ceased with the onset of the mastoid swelling or has been distinctly intermittent in type, being sometimes profuse and sometimes scanty. The periods of diminution in the discharge are generally accompanied by an increase in the pain. In contradistinction to such intermission, the discharge may be continuously very profuse. This profusion indicates involvement of a wide area, and therefore of the mastoid antrum and cells. The character of the temperature chart is of some assistance. Most usually, after a few days' fever, the temperature falls, and then, after a short period, commences to rise again. While this remission, if present, is very significant, not infrequently it does not occur, the temperature remaining high from the commencement of the acute otitis.

A characteristic factor in the tenderness over the mastoid process is that it is confined to one particular spot, and is deep and not superficial. In testing for this the finger should be firmly passed down over the mastoid process while the eye is being watched. The presence of tenderness is always evidenced by a flinch. This obviates, to a large extent, the personal factor of the patient.

Having thus detailed the characteristic symptomatology, it may be said that any, or all, of these symptoms or signs may be absent.

Cases Without Aural Discharge.

Cases presenting signs of inflammation of the mastoid process, but with a normal middle ear, are not infrequently seen. In these a history of previous aural discharge may be present. The explanation is simply that while the middle-ear inflammation, suppurative or catarrhal, has cleared up, the infection in the mastoid process has progressed.

Latent Type.

The characteristic of cases in this group is the triviality of the signs and symptoms, and I can best illustrate this class by a description of a case recently dealt with.

The patient, a clergyman aged 52, following a "cold," had earache with discharge for twenty-four hours only. For two months afterwards he suffered from a vague ache in the ear, with considerable diminution in the hearing and a general feeling of unfitness. During this period the temperature was normal and he was able to continue his duties. The diagnosis of a suppurating focus in the mastoid was made on this history, combined with the presence of a localized tender spot towards the apex of the process, with some opacity and injection of the tympanic membrane. Operation showed pus under tension in the mastoid cells, and drainage produced rapid restoration of the hearing, with return to health.

The difficulty in such cases is to persuade the patient that he has a condition sufficiently severe to need an operation. If left, the suppuration will probably eventually reach the surface and produce manifest symptoms, but complications may supervene.

The points that I have found of most help in diagnosis in this group have been: the presence of a dull ache in the ear; a point of localized tenderness, which may be situated at any portion of the mastoid process; if discharge is present, either its profusion or intermission; the persistence of a defect in hearing of a conductive type; some bulging of the posterior superior meatal wall; and, finally, a haziness in outline of the mastoid cells, in a skiagram, as compared with the other side.

Fulminating or Osteomyelitic Type.

While in the majority of cases of acute mastoiditis the disease will progress for a considerable period without any great risk to life, in the class which I am now trying to describe I believe that early operation is of paramount importance. The cases are associated with a streptococcal infection, frequently haemolytic in type, and, in them, the infection from the beginning widely invades the bone and produces severe constitutional disturbance. They are most commonly met with in association with epidemics of acute sore throat and are due to a spread of infection up the Eustachian tube. There is high fever from the commencement, and the discharge from the ear is unusually profuse, watery, and often blood-stained. Widespread tenderness over the mastoid occurs early. Such cases not infrequently

* Read at a meeting of the Bath and Bristol Branch of the British Medical Association, January 29th, 1930.

develop septicaemia, and the mortality is probably 10 per cent. The infection in these cases does not, however, seem prone to invade the respiratory tract.

The diagnosis is based on the coexistence of an acute sore throat, the high temperature, rapid pulse, furred tongue, profuse bloody discharge, and widespread mastoid tenderness. Early operation is indicated with provision of the freest possible drainage. Scarlatinal antitoxin sometimes seems to be helpful in septicaemic cases.

Mastoiditis in Infants with Enteritis.

It has been pointed out recently that many infants dying of an acute gastro-enteritis show, at necropsy, the presence of an acute otitis media and mastoiditis. It has been suggested that this latter infection is the primary one, and that the gastro-enteritis is a secondary result of the acute toxæmia of the middle-ear infection. I am not prepared to offer any opinion on this, but would only emphasize the importance of examining the ears of infants suffering from any acute febrile infection.

Parotid Mastoiditis.

Cases are occasionally met with in which an inflammatory swelling occurs in front of or above the ear as a result of mastoiditis. Such swelling is often associated with some oedema of the eyelids on the same side, and errors in diagnosis are common. The explanation of such cases is that the infection reaches the surface through cells at the root of the zygoma. If this possibility is recognized the diagnosis can be established by the signs or history of the middle-ear disease, together with the presence of some tenderness over the mastoid process.

Bezold's Mastoiditis.

This is a similar type, in which, however, the swelling occurs in the neck, the explanation being that the suppurative bursts through a cell at the apex of the mastoid process and on the deep side of the sterno-mastoid insertion. The remarks just made as to diagnosis apply here also.

Tuberculous Mastoiditis.

Tuberculosis of the middle ear and mastoid process is usually met with in the infant. In a typical case, it is characterized by occurring in a child of under 1 year with painless and watery aural discharge, enlarged glands around the ear, and facial paralysis on the same side. The infection has been introduced via the Eustachian tube by infected milk.

Spurious Mastoiditis.

Two conditions may simulate an acute mastoid infection—namely, a furuncle of the posterior meatal wall and the so-called herpes oticus.

A furuncle of the posterior meatal wall not infrequently produces a tender swelling behind the auricle. Owing to the meatal swelling, the membrane is invisible and diagnosis may be difficult. If the hearing is relatively good in the affected ear, otitis media is unlikely. The slight degree, or absence of fever or constitutional disturbance is also helpful. In addition, tenderness on moving the auricle is suggestive of a furuncle rather than of a bone infection. Sometimes, however, it is impossible to decide whether one is dealing with a meatal furuncle or a mastoid suppuration, except by the performance of an exploratory incision. If, as a result of such incision, it seems reasonably probable that the soft parts only are involved—that is, that one is dealing with a meatal condition—it is important to avoid any interference with the bone, owing to the risk of introducing infection. I can remember a case of infection of the soft parts in which a fatality ensued as the result of opening the bone.

Herpes oticus is the name given to a rare condition, in which herpetic inflammation of the ganglia of the seventh and eighth cranial nerves gives rise to intense pain in and around the ear, with a herpetic eruption on the pinna and in the meatus, facial paralysis, and nerve deafness. The combination of earache, aural discharge from the ruptured vesicles, deafness, and facial palsy, not infrequently leads

to a mistaken diagnosis of acute otitis media and mastoiditis. If it is realized that such a condition may occur, it will be recognized.

The diagnosis of an acute infection of the mastoid process, therefore, depends on the recognition of one or more of the many possible signs and symptoms. Pain may be slight or absent. Tenderness is usually present and often strictly localized. Fever and aural discharge are not infrequently absent. Some defect in hearing is more or less constant, and swelling, when present, may be situated behind, in front of, above, or below the ear.

In conclusion, a word or two as to the type of operation to be performed. The object of such operation is to provide drainage. Care should therefore be taken to open up all infected cells. Since, however, the middle ear will recover if not interfered with, it is unwise to pass a bent probe into the aditus to make sure that one has opened the antrum. This proceeding, although commonly done, is extremely liable to dislocate the incus, with resultant permanent damage to the hearing.

It was formerly the practice to pack the operation cavity with gauze and to continue such packing until the cavity had become obliterated. This, in addition to being painful, is unnecessary. In latent cases the incision can be sewn up except for the provision of a small rubber drain at the lower end. In the more acute cases, I find that the best results are obtained by leaving wound and cavity widely open. If, however, instead of filling the cavity with gauze, either thin rubber dam or a strip of oil-silk is used, the pain associated with subsequent dressings is entirely obviated. As soon as the cavity is granulating and showing signs of closing, I believe that there is nothing to be gained by attempting to keep it open with any form of packing.

BRAIN ABSCESS WITH UNUSUAL FEATURES.

REPORT OF TWO CASES.

BY

E. G. COLLINS, M.B., F.R.C.S. Ed.,

HONORARY SURGEON, EAR, NOSE, AND THROAT DEPARTMENT, NORTHERN INFIRMARY, INVERNESS.

BRAIN abscesses may be classified under the three headings of adjacent, traumatic, and metastatic. The two cases which are here recorded belong to the last group, for, although the second case is of aural origin, yet pathologically and clinically it falls into the metastatic class. In both instances the main cerebral infection was contralateral to the original focus.

CASE I.

A man, aged 24, was admitted to the Northern Infirmary, Inverness, on May 25th, 1929. Ten days before admission he had received a punctured wound of the right lower eyelid from a garden stake. The wound oozed a little, and he had some headache, especially on the right side, for a few days afterwards. His vision was not affected. He had had a similar accident to the left upper lid ten years previously.

Examination revealed a punctured wound of the inner half of the right lower eyelid extending upwards and inwards for half an inch, but apparently not perforating the conjunctiva. Both lids showed some inflammatory oedema, and pus was present in the conjunctival sac. The bulbar conjunctiva showed chemosis and generalized congestion. The remaining structures of the eye were normal.

Progress Notes.

May 26th to June 3rd: The chemosis and discharge disappeared, but there was some bleeding from the nose, and severe frontal headache, worse at night. He vomited twice. The general condition improved.

June 4th: Pressure below the supraorbital margin revealed considerable tenderness on both sides. No pus was seen in the middle meati, and no oedema or redness of skin over the frontal sinuses. Transillumination and x-ray examinations showed relative obscurity. The diagnosis made was double acute frontal sinusitis, and conservative treatment was ordered.

June 6th to 12th: Headaches not so severe; less tenderness on pressure over the sinuses. He complained of stiff-neck and tenderness of the muscles of the nape of the neck, giving pseudo-head retraction. There was some pain in the epigastrium.