Experiments were made with many methods of prevention. The Schultz-Charlton test merely corroborated what was usually known—namely, that a bright rash indicated scarlet fever; the Dick test was found to be of little value in the positive cases where the immunity had not had time to develop or where there was susceptibility to the disease. If the Dick test is repeated at the end of a week useful information is obtained, but susceptible patients are meanwhile exposed to infection. Active immunization by means of scarlet fever prophylactic develops too slowly to be of any real value; and barrier isolation, though employed to some purpose, was ineffective in wards where all the patients were not confined to bed.

The production of passive immunity by means of scarlet fever antitoxin was then considered. A small dose was said to give a passive immunity for ten to fourteen days. This period was not considered to be long enough, as "peeling" in many mild cases often does not occur until about the fourth week. If a dose is repeated within ten days there appears to be no danger of anaphylaxis, as sensitization to a foreign protein does not begin until after that period. The risk was lessened by administering only 2½ c.cm. at first.

The following routine method was therefore gradually adopted. All patients with indefinite symptoms received 2½ c.cm. of scarlet fever antitoxin on admission, a second dose of the same amount on the tenth day, and a third dose on the twentieth day. This was all that was necessary as a rule, but some received a fourth dose, and a few a fifth dose at the same intervals.

The doses have recently been increased to 5 c.cm., on the advice of the makers, though the smaller doses appeared to

give good results. The number of patients that received treatment was 108; of these, 30 patients received two doses; 56 received three doses; 20 received four doses; and 2 received five doses. When typical desquamation occurred no further injections were given, and patients who showed no sign of desquamation by the fourth week were discharged. For various reasons some were detained for longer periods.

Classification of these 108 cases showed that 34 of them were probably not scarlet fever; of these 34 patients only one developed the disease, though all of them were exposed in the wards during the whole time. This patient was detained in hospital for a certain reason for forty days, but received only three doses of the serum, the last dose being on the twentieth day. Two days after discharge, on the twenty-second day after the last prophylactic dose, there developed a typical attack of scarlet fever. Henceforth all patients without definite signs received a prophylactic dose every tenth day until discharge.

Signs of Serum Reaction.—Signs of serum sickness have been almost entirely absent. Two patients developed a very mild urticarial rash on the thirteenth day after the first dose or the third day after the second dose; one developed a similar rash on the sixth day after the third dose; one had a rise of temperature of two degrees the morning after the third dose, and one had a rise of one degree on the day of injection of the third dose. None of the other cases showed adverse signs of any kind.

The method no doubt requires further testing, but in my application of it during the past three years it appeared to have been very successful, as ward infections have almost disappeared.

THALLIUM ACETATE AS A DEPILATORY IN RINGWORM OF THE SCALP.

BY

H. McCORMICK MITCHELL, M.B., C.M.ABERD.,
CONSULTING DERMATOLOGIST, CANCER AND SKIN HOSPITAL, LIVERPOOL;
DERMATOLOGIST, WEST DERBY UNION HOSPITALS.

COMPLETE epilation of the scalp is generally considered to be the best and most expeditious form of treatment for ringworm of the scalp. Epilation by x rays has for many years been the method employed, and has given very satisfactory results, but it has some drawbacks which cannot be overlooked.

1. If there is any inflammation or pustular condition of the scalp x rays are, in my opinion, contraindicated as a depilatory agent. Moreover, in a small number of cases a troublesome dermatitis follows their use, although the scalp may appear quite healthy, apart from the tinea, before radiation.

2. X-ray epilation is unsuitable for children under the age of 3, since they cannot be kept still during the application.

3. The most serious drawback is the permanent baldness which follows in some cases unless the treatment is given by someone highly skilled at the work.

I have been particularly struck with the variation in the output of x-ray tubes from day to day, and since the margin of error is small on either side the exact output of the tube should be carefully checked each day before work is commenced. The use, therefore, of any other method of producing depilation is worth careful consideration.

The depilatory action of thallium acetate has been known for many years. More than a quarter of a century ago Sabouraud used this drug in his clinic for the treatment of tinea capitis, but he seems to have abandoned it on account of its toxic effects on the patients. Recently its employment for the treatment of ringworm of the head was brought into prominence by A. Buschke, and it is almost entirely due to his experiments on animals that the method has been made practical and safe.

The success and advantages of thallium acetate depend entirely on correct dosage. Buschke, after much research and in collaboration with L. Peiser, established the minimal dose for complete epilation of the scalp to be 8 to 9 mg. per

kilo of the body weight. One dose is, in the majority of cases, quite effective in producing complete depilation, and can be administered either in sweetened water or mixed with food.

The number of patients treated by me has been 74; of these, 51 were under the age of 4; 9 were aged 4; 3 were aged 5; 7 were aged 6; 3 were aged 7; and 1 was 8 years old.

I have used throughout the preparation made by Kahlbaum of Berlin, who supplies the drug in tablet form containing 1 decigram, 1 centigram, and 1 milligram. I have varied the dose from 8 to 9 mg. per kilo of the body weight, and have found that the cases of failure, to which I shall allude later, were among those to whom I gave 8 mg. All that is required is to convert the weight of the child into kilograms and then multiply by 8, 8.5, or 9 milligrams, according to the dose intended, administering the required number of tablets dissolved in sweetened water or mixed with food.

The result of a single dose of thallium acetate is very striking. On the sixth or seventh day the hairs begin to loosen; by the fifteenth day they are coming out freely; and by the twentieth day depilation is almost complete, except for those fine hairs at the margin of the scalp which, in my experience, never fall out, and the same applies to the eyebrows and eyelashes. I have never administered the drug to an adult, so cannot say what the effect is on the hairs of the face, the pubes, and the axillæ, but the experience of other workers seems to be that a dose sufficient to cause depilation of the scalp does not cause the hairs on the above parts to fall out. The infected stumps, as is the case after x-ray depilation, do not fall out so readily as the non-infected hairs, and this fact should be carefully borne in mind in the after-treatment. Removal of the affected hairs and stumps is most important in guarding against the reinfection of the new hairs, which appear much earlier than after x-ray epilation.

The routine treatment is as follows.

The urine in all cases is examined for albumin and casts; the weight of the child is converted into kilograms, and 8.5 or 9 mg. of thallium acetate per kilo of body weight is given in a little sweetened water or mixed with food. From the first day after the thallium is given the scalp should be washed daily with sp. saponis kalinus, or a 10 per cent. sulphur soap, and a fungicide ointment well

rubbed in. When the hair begins to fall freely Buschke recommends the use of tincture of iodine and a fungicide ointment on alternate days. I have followed this practice carefully in all my cases with great success.

After the eighteenth or twentieth day, when epilation is almost complete, a careful examination of the scalp is made with the ringworm lamp for the detection of infected stumps; these are removed as far as possible with the epilation forceps, and by adhesive strapping. It is advisable to use local treatment for ten or fourteen days after a careful examination with the ultra-violet beam has failed to detect the presence of infected hairs or stumps. If care is not taken to remove the infected hairs as quickly as possible the new hairs are liable to become reinfected, owing to the rapidity with which they appear after depilation.

Of the cases treated, 65 were cured after one dose of thallium acetate. In 10 cases the hairs were infected after the regrowth, and a second dose was necessary; in only one case had a third dose to be given.

Most of my failures occurred among my early cases, and I attribute my non-success to two causes: (1) At first I gave 8 mg. per kilo of body weight; this dose, in the light of my experience, is not sufficient to produce complete depilation, and I now invariably give 8.5 or 9 mg. (2) As the ringworm lamp was not available during the first few months for examination purposes, a number of infected stumps were not detected and removed, with the result that reinfestation of the new hairs occurred. Since giving 8.5 or 9 mg. per kilo of body weight and conducting a daily examination for infected stumps with the ultra-violet beam after epilation is apparently complete, my results have been much more satisfactory.

One or two further points in connexion with this treatment are, I think, worth recording. (1) Children with dark hair are more difficult to depilate than those with fair hair; it is therefore advisable to give 9 mg. per kilo to the former class of case. (2) The hairs on the crown fall out less readily than those on the other part of the scalp, so that for ringworm in this situation it is also better to give the full 9 mg. dose.

Toxic Effects.

I have not seen any alarming toxic symptoms in any one of the cases I have treated; this was due partly to the fact that I insist on all patients remaining in bed for forty-eight hours after the thallium acetate has been administered, and partly to the fact that most of my patients were under the age of 4 years, as it is well known that young children are less susceptible to the toxic effects of this drug than older children.

The toxic effects most noticeable were apathy, loss of appetite, and drowsiness during the first two or three days, and pain in the joints, particularly in the lower limbs, appearing from the sixth to the tenth day. All these symptoms gradually disappeared without leaving any ill effects in any of the patients treated. All the children who showed toxic symptoms in any marked degree were more than 4 years old.

Summary.

As the outcome of my clinical investigation the following points might be emphasized:

1. Thallium acetate can be safely used as a depilatory agent where, on account of the age of the patients, x-ray treatment is contraindicated.
2. Accurate weighing of the patient and correct dosage are most important.
3. The after-treatment is all-important in obtaining satisfactory results; the patients should be kept in bed for forty-eight hours after the initial dose of the drug, and from the first day of treatment the scalp should be washed daily with sp. saponis kalinus, or with 10 per cent. sulphur soap, and a fungicide application be well rubbed in.
4. Careful examination with the ultra-violet beam should be made after depilation is apparently complete, so as to detect any infected stumps which have failed to come out and which would reinfest the new growth of hair.
5. The use of thallium acetate should, as far as possible, be limited to children under the age of 5, owing to its toxic effects on those above that age.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

HAEMATOMETRA WITH HAEMATOSALPINX IN ONE HORN OF A UTERUS BICORNIS UNICOLLIS.

THE rarity of this condition prompts us to record the following case.

The patient, aged 16, was sent to hospital on October 9th, 1929, by Dr. Crawford, for pain and swelling in the perineum.

History.—Menstruation began at 14 years of age; regular cycle of twenty-eight days, lasting five days, loss scanty. Last menstrual period September 10th, 1929. At the onset of menstruation the periods were free from pain, but during the last year there had been dysmenorrhoea of progressive severity. This pain, which started two days before the period, and continued throughout, was referred to the lower abdomen and perineum. On the second day of her last period the patient suddenly fell in a sitting posture, and a few hours later complained of severe pain in the right lower abdomen and perineum, which was worse than before. This pain persisted throughout the following month, making walking painful and sitting difficult; she also suffered from dysuria. At the onset of her next period the pain became so severe that she was sent to hospital.

Examination.—The patient was well developed but anaemic. In the right iliac region there was felt a sausage-shaped swelling, soft in consistency, irregular in outline, and tender to pressure, but with no rigidity of abdominal parietes. The right vaginal wall was bulged inwards by a globular swelling, which extruded from the vaginal orifice upwards into the pelvis, and was definitely cystic. At the vault of the vagina the os externum could be detected, but no true cervix could be felt.

Bimanual Examination.—It was difficult to arrive at any conclusion, owing to the cystic swelling, the size of a large orange, occluding the vagina. As far as could be ascertained, there was a mass in the position of the uterus, continuous above with sausage-shaped mass below with the cystic swelling.

While awaiting operation, the patient suddenly developed acute abdominal pains with vomiting, the lower abdomen became rigid, and the patient took on the aspect of an abdominal crisis.

Operation.—Under general anaesthesia the peritoneal cavity was opened, when there was an escape of a large amount of altered blood. The uterus was enlarged to the size of a three months' pregnancy, and on further examination was found to be bicornute; the left horn was the size of a normal virgin uterus, with a healthy tube and ovary. The right horn was slightly larger, and merged into a large globular swelling, which had burrowed down, lateral to the right wall of the vagina. The tube on this side was greatly dilated, and altered blood was seen issuing from the fimbriated end. The whole, except the left tube and ovary, was removed in one mass. Great difficulty was experienced in separating the cystic swelling from the deep part of the lateral wall of the vagina and surrounding tissues. The specimen showed the left horn to contain a normal cavity and cervical canal opening into the vagina. On the right side the large cavity, which contained altered blood, appeared to be a greatly distended cervical canal, above which was the enlarged right horn of the uterus with the haematosalpinx.

J. E. G. CALVERLEY, C.M.G., M.D., B.S.Lond.,
Surgeon,

J. W. D. BUTTERY, F.R.C.S.Ed.,
Assistant Surgeon,
Royal Victoria Hospital, Folkestone.

DRY FRACTURE OF NECK OF FEMUR.

I was particularly interested in Mr. Blundell Bankart's description (*Journal*, January 4th, p. 8) of a case of intracapsular fracture of the neck of the femur, in which he had obtained bony union after injection of blood into the hip-joint. I must admit that I was sceptical of the absolute absence of bleeding between the ends of the fracture and into the hip-joint, especially as I had often noticed an apparent fullness over Scarpa's triangle, and considered that it was due to effusion into the hip-joint. I wish, therefore, to place on record the following case, which showed a dry fracture of the neck of the femur investigated at necropsy.

W. M., aged 81, was admitted to the Radcliffe Infirmary with retention of urine and uraemic symptoms due to senile enlargement of the prostate. His retention was relieved by catheterization, but two days after admission he got out of bed one night and fell, breaking the neck of his femur. He was treated