FIG. 1.—Case i. ($\times 90$.)

FIG. 2.—Case ii. ($\times 25$.)

FIG. 1.—Showing triangular shadow within the cardiac shadow.

FIG. 2.—Triangular shadow absent. Lipiodol showing no evidence of bronchiectasis.

FIG. 3.—Showing the triangular shadow with bronchiectasis within it.

FIG. 4.—Showing bronchiectasis within the triangular shadow.

FIG. 5.—Showing triangular shadow at right base.

Dr. Grace's Report.

CASE I.—Subperitoneal layer thickened and packed with lymphocytes. Number of lymphatics also full of lymphocytes. Muscular layer normal. Submucosa enlarged, with several small vessels packed with lymphocytes. Histological character consistent with diagnosis of syphilitic infection.

CASE II.—Chronic inflammation scattered throughout submucosa and underlying layers.* There is a predominance of lymphocytic infiltration, and the findings support a diagnosis of syphilis, but further than this it is not possible to go.

There was no evidence of tubercle in either specimen. Efforts to show up the spirochaete by Levaditi and other silver methods were unsuccessful, but, according to Warthin,² this is frequently the case in undoubted syphilis.

REFERENCES.

¹ *Semana Med.*, 1929, i, 301. ² *British Medical Journal*, August 10th, 1929.

ATELECTATIC BRONCHIECTASIS.

BY

A. J. SCOTT PINCHIN, M.D., M.R.C.P.,

PHYSICIAN AND PHYSICIAN IN CHARGE OF RADIOGRAPHIC DEPARTMENT,
CITY OF LONDON HOSPITAL FOR DISEASES OF THE HEART AND
LUNGS, VICTORIA PARK; PHYSICIAN, WEST LONDON AND
HAMPSTEAD GENERAL HOSPITALS;

AND

H. V. MORLOCK, M.C., M.D., M.R.C.P.,

PHYSICIAN TO OUT-PATIENTS, CITY OF LONDON HOSPITAL FOR DISEASES
OF HEART AND LUNGS; ASSISTANT PHYSICIAN, HAMPSTEAD
GENERAL AND MILLER GENERAL HOSPITALS.

(With Special Plate.)

PERHAPS it is true to say that in no branch of medicine has radiology been more useful as a diagnostic aid than in disease of the chest; nevertheless, though the skiagram may show a condition which is definitely abnormal, yet the abnormality present is not diagnostic of the underlying pathological lesion; and, in order to ascertain the nature of the abnormal shadow, the aid of clinical medicine and pathology must be sought.

Such a radiological abnormality is a triangular shadow which is seen at the base of the lung. The shadow is a rectangular triangle, one side of which rests on the diaphragm, the other along the mediastinum; the hypotenuse runs from the mediastinum to the diaphragm. This shadow may appear on either side of the mediastinum, and when it occurs on the left it may lie within the cardiac shadow, and thus tends to be obscured and may be unrecognized at first.

In 1911 Dieulafoy¹ described the clinical findings in cases of mediastinal effusions; not long afterwards the radiological appearance of this condition was reported as being a shadow similar to the triangular shadow described above. The following case illustrates this.

H. W., aged 11 years. Four weeks before admission to hospital the patient had felt "off colour"; slight cough was present, with occasional traces of sputum. Physical examination on admission showed impaired percussion note and absent air entry at the left base behind. Temperature 99.8° on admission. The skiagram showed a triangular shadow behind the heart shadow (see Fig. 1). After a few days in hospital the temperature fell and the patient felt quite well; during the four ensuing weeks the physical signs gradually disappeared. A second skiagram showed that the shadow had gone, and that lipiodol could not demonstrate the presence of any bronchiectasis (see Fig. 2).

One brother of the patient had recently died of pulmonary tuberculosis, and another brother was suspected of the disease. The effusion was probably tuberculous in origin. Wallgreen² has described a similar case in which diagnosis was confirmed by exploratory puncture.

Recently the occurrence of massive atelectases of the lung has attracted considerable attention, from both the clinical and radiological sides, particularly the massive atelectases occurring as a post-operative complication, or as a result of haemoptysis; but massive atelectasis may occur in other pulmonary conditions—for example, trauma to the chest wall, pneumonia, and pleurisy. The atelectatic lobe, or part of the lobe, gives a dense shadow on the

skiagram, and the mediastinal shadow is displaced towards the side of the lesion. The shadow of the atelectatic lobe may be similar to the triangular shadow now under discussion.

D. S., aged 8 years, came to hospital complaining of a cough which had been present for some time, but which recently had become worse. Physical signs revealed an area of dullness and bronchial breathing at the right base behind. The heart was displaced slightly towards the lesion. The skiagram shows the triangular shadow at the right base with the cardiac shadow drawn towards it. In view of the past history of pneumonia the diagnosis of bronchiectasis was made, and the patient was sent away to a convalescent home. On return the physical signs had disappeared, and lipiodol injection of the lungs showed no bronchiectasis; the shadow had gone from the skiagram and the heart had returned to its normal position.

In the following series of cases the skiagrams of the patients' chest showed the triangular shadow. In their histories these patients admitted to the raising of much sputum, or dated their cough from their recovery from pneumonia; which facts suggest the possibility of a bronchiectasis; the lung was therefore injected with lipiodol, and bronchiectasis was demonstrated within the shadow.

W. A., female, aged 17 years. Had pertussis when she was 4 years old; since then bronchitis, and during the last four years repeated haemoptysis. Patient had been notified as suffering from pulmonary tuberculosis. Physical signs: dullness and added sounds at left base. A skiagram showed a triangular shadow behind the heart (see Fig. 3), and lipiodol injection demonstrated bronchiectasis within the shadow.

B. P., aged 11 years, complained of cough which had been present for "years." The amount of sputum raised varied from 1 to 2 ounces. There was a history of patient having had pneumonia at 3, 5, and 10 years of age. Physical signs at left base; dullness, bronchial breathing, and rales. A skiagram showed a triangular shadow behind the cardiac shadow, and lipiodol injection demonstrated bronchiectasis in this shadow.

F. R., aged 14, had suffered from cough for "years." There was very little sputum. Past history of measles. Physical signs were present at the left base, and the fingers showed slight clubbing. A skiagram showed a triangular shadow behind the heart (see Fig. 4); lipiodol demonstrated bronchiectasis within the shadow.

E. S., aged 9 years, complained of cough which had been present since the last attack of pneumonia two years previously. Patient had had pneumonia three times. The sputum raised was offensive and varied from 2 to 3 ounces daily. Physical signs at right base; behind dullness and bronchial breathing, added sounds. Fingers clubbed. A skiagram showed the triangular shadow at the right base. Repeated attempts at lipiodol injection were unsuccessful (see Fig. 5). The facts that the amount of sputum was large and that the physical signs and skiagram remained unchanged for two years leave little doubt that the condition was one of bronchiectasis.

In 1926 Singer and Graham³ described a case the skiagram of which presented the triangular shadow, and in which bronchiectasis was demonstrated by lipiodol. At operation an atelectatic lung with bronchiectasis was found to correspond to the shadow. Rist, Frocine, and Jacob⁴ collected a series of seven such cases, which at first they suspected as being purulent mediastinal effusion, but the lipiodol injection showed the conditions to be bronchiectatic. Wallgreen reported five such cases, and discussed the pathology. He believes that the primary condition in these cases is an atelectasis and bronchiectasis.

It is a matter for speculation how much influence local areas of collapse exert in the etiology of bronchiectasis generally, though it is probable that it may have some bearing in relation to the disease occurring after chronic pneumonic conditions. The clearing up of a collapsed area of lung is dependent on the free mobility of the chest and diaphragm, and occurs least readily in those parts where there is a deficiency of movement.

This condition obtains particularly in that portion of the lung under discussion. Lying as it does in the recess formed by the spine and ribs and the angle of the diaphragm, it is shielded from the more forcible respiratory movements, and is a spot where one would expect to find a residual atelectasis. In such manner we account for the comparative frequency of the condition described: we consider, from the number of cases we have met with, that

* The appearance at operation was anything but chronic.

it is a not uncommon event, and may account for some of those cases of continual attacks of haemoptysis, with no definite signs of tuberculosis or gross disease, which are met with fairly frequently.

In this communication we have called attention to this triangular shadow, and we have discussed its significance; in our experience the most common lesion with which it is associated is bronchiectasis, and we are in agreement with Sergeant and Wallgreen that the presence of this shadow should always raise the suspicion of a bronchiectasis unless other clinical evidence is forthcoming to clinch the diagnosis of a mediastinal effusion.

Our best thanks are due to Dr. F. G. Chandler for his kind permission to use the notes and skiagram of the case F. R.

REFERENCES.

- ¹ Dieulafoy: *Manuel de Pathol.*, interne 1 (1911).
- ² Wallgreen, A.: *Beit. z. Klin. d. Tuberk.*, 1928, lxxviii, 641.
- ³ Singer and Graham: *Amer. Journ. Roentgenol.*, 1926, xv, 54.
- ⁴ Rist, Frocine, and Jacob: *Annals of Medicine*, 1927, xxi, 144.
- ⁵ Serger and Bodet: *Bull. Soc. Méd. des Hôp. de Paris*, 1927, 742.

EXPERIENCES WITH REGIONAL ANAESTHESIA.

BASED ON 208 CASES IN WHICH IT HAS BEEN USED.

BY

H. W. L. MOLESWORTH, F.R.C.S.,
ASSISTANT SURGEON, ROYAL VICTORIA HOSPITAL, FOLKESTONE.

It is probable, if the discovery of novocain had preceded that of cocaine and of chloroform and ether, regional anaesthesia would by this time have been firmly established as the standard method of averting the pain of operations. Events having occurred as they did, regional anaesthesia has had to struggle against adverse conditions, but has steadily gained ground as safer drugs and improved technique have been evolved. One of the chief obstacles to its wider adoption is the entrenched belief that for major operations general anaesthesia is essential. Early fatalities, due to the use of cocaine, or to technical errors in administration, have done the method much harm. Injections of cocaine or of high titres of novocain (1 or 2 per cent.) have been the greatest errors; death has resulted from intravenous and intraspinal injection. According to Finsterer intravenous injection is easily avoided by adherence to two rules: (1) in regions free from large blood vessels injections should be made only with a moving needle point; (2) in dangerous areas it is necessary to aspirate before injection and repeatedly during injection.

The number of cases (208) here dealt with is too few to suggest conclusions about the relative safety of regional and general anaesthesia, but in this series there has been no death directly attributable to the anaesthetic, and one only in which it is possible that the anaesthetic (spinal) played a part. A survey of the series can, however, give some idea of the value of the method to a general surgeon who is prepared to give some time and trouble to the acquisition of a technique which, though not easy, is fascinating.

Drugs and Solution Strengths.

I have tried novocain, borocain, tutocaine, quinine-urea, and eucupin, and though the first three of these have given satisfactory service, I prefer tutocaine, for the reasons presently to be stated.

Novocain (with Adrenaline).—Novocain has a low toxicity. Solutions of 1 in 400 may be used for infiltration anaesthesia; 1 in 200 is adequate for field-block procedures, and 1 per cent., and in rare instances 2 per cent., for blocking the large nerves. The stronger solutions should be used with extreme caution.

Borocain will give similar results; it is said to be superior for producing surface anaesthesia and to be less toxic than novocain. I can state from personal experience that 10 c.cm. of a 2 per cent. solution may produce toxic symptoms in a patient of lowered resistance. Solutions of borocain have the disadvantage that they are extremely sensitive to slight changes in reaction.

Tutocaine is, strength for strength, more toxic than novocain, but it appears to be much quicker in its action,

and in 1 in 500 solutions has proved a thoroughly reliable anaesthetic. A 1 in 500 solution of tutocaine appears to be as effective for field block as a 1 in 200 solution of novocain.

Quinine-Urea.—I have not given this drug an extensive trial. Used alone it causes initial pain on injection, and it has a tendency to produce oedema, which persists for some time.

All these solutions are made up with normal saline and are sterilized by boiling; adrenaline is added (the synthetic variety will withstand a few seconds' boiling). The solution becomes pinkish; it may be used so long as the colour is not a mahogany; 1 c.cm. of 1 in 1,000 adrenaline is added to each 250 c.cm. of solution.

Classification of Results.

Many difficulties arise in attempting to assess the results of regional anaesthesia, two of the chief being the following: (1) the patients vary greatly in their reaction to a painful, uncomfortable, or an emotional stimulus; (2) it is difficult to assess the quality of anaesthesia where inhalation anaesthesia has been used in combination with regional anaesthesia. All cases in which it has been intended from the first to combine the two methods are classified in the following table as "intended combined." The quality of anaesthesia is classified as "poor" in

Table showing Results in 208 Cases in which Regional Anaesthesia was Employed.

Operations.	No. of Cases.	Intended Combined.	Quality of Anaesthesia.			Technique.
			Good.	Fair.	Poor.	
Major operations on stomach and gall-bladder	26	2	14	—	—	Anterior splanchnic anaesthesia (Braun).
Ditto	5	1	2	1	1	Posterior splanchnic anaesthesia (Kappis).
Ditto	4	—	2	2	—	Mesenteric anaesthesia (Finsterer).
On abdominal wall	15	3	12	—	—	Field block.
Appendicectomy ...	6	—	4	2	—	Field block and mesenteric (Finsterer).
Hernia: radical cure ...	15	—	14	—	1	Field block.
Jejunostomy and caecostomy	5	—	5	—	—	Field block.
Various abdominal operations	50	1	31	12	6	Spinal anaesthesia.
Suprapubic cystotomy	11	—	11	—	—	Field block.
Perineal prostatectomy...	4	—	4	—	—	Trans-sacral.
Suprapubic prostatectomy	3	—	—	3	—	Field block and epidural.
Operations on testicles ...	2	—	2	—	—	Field block.
Urethrotomy	3	—	3	—	—	Epidural 30 c.cm. 1/5% tutocaine.
Cystoscopy in males ...	15	—	14	—	1	Ditto
Excision of breast ...	3	—	3	—	—	Field block.
Empyema	4	—	1	3	—	Field block.
Thyroidectomy: excision of glands of neck	5	—	5	—	—	Paravertebral cervical block.
Perineorrhaphy repair of cervix (D. and C.)	4	—	4	—	—	Field block; broad ligament block.
Haemorrhoids	5	—	4	1	—	Field block.
Operations on upper extremity	6	—	4	1	1	Brachial plexus block (Kulen-kamp).
Hallux valgus; Mayo's operation	3	—	3	—	—	Field block.
Sympathectomy, femoral	1	—	1	—	—	Field block.
Excision of tongue ...	1	—	1	—	—	Field block.
Operations on foot ...	2	—	—	1	1	Anterior tibial and posterior tibial block.
Saphenous varix	1	—	1	—	—	Field block.
Circumcision	3	—	2	1	—	Field block.
Varicocele	2	—	2	—	—	Field block.
Amputation of penis ...	1	—	1	—	—	Field block.
Hydrocele	2	—	2	—	—	Field block.
Operations on hand ...	1	—	1	—	—	Median and ulnar block.