become too stereotyped. A distrust of tinkering and a desire to be thorough is characteristic of British surgery, but the mortality rate of total prostatectomy is still too high to allow of our being content with our present position.

Another argument, in addition to that of "tinkering," has been urged against operations that do not entail the complete removal of the prostate—namely, that some 14 per cent. of seemingly innocent enlargements contain foci of malignancy. But this argument presupposes that prostatectomy eliminates the possibility of subsequent malignant disease, which it does not. Enucleation leaves behind sufficient prostatic tissue to allow of a recurrence of malignant disease, and although the risk may be diminished by the operation, it is never eliminated. Moreover, the increased danger of prostatectomy more than counterbalances any slight addition to the risk of malignant disease that a partial operation may have entailed.

PUERPERAL SEPSIS.

AN INVESTIGATION REGARDING CONTAGION, AND THROAT CARRIERS OF THE STREPTOCOCCUS HAEMOLYTICUS.

BY

E. FARQUHAR MURRAY, M.D., F.R.C.S.,
ASSISTANT GYNAECOLOGIST, ROYAL INFIRMARY, NEWCASTLE-ON-TYNE.

THE respective parts played in the production of puerperal sepsis by contagion and infection from throat carriers of haemolytic streptococci have been investigated at the Princess Mary Maternity Hospital, Newcastle-on-Tyne. Outbreaks of pyrexia and definite sepsis occurring from time to time among "clean" cases, caused much anxiety and demanded investigation.

In addition to its own booked cases, the hospital receives patients from a very large area, sent in by doctors on account of puerperal sepsis, difficulties arising during or immediately after labour, abortions, and so on. Many of these cases, when admitted, are "suspect" as regards infection, and many are frankly septic. The figures for 1929 were: puerperal sepsis, 108; "suspect," 302; abortions, 280. Patients requiring Caesarean section are not included, nor are those patients suffering from pyelitis and other medical conditions which were admitted during pregnancy from the ante-natal department, or sent in by doctors. Many of the "abortion" cases were septic or developed temperatures after admission.

The existing system had certain obvious defects. (1) All cases, whether suspect or septic (except Caesarean sections) were delivered in the labour ward. (2) The isolation department, consisting of 9 beds, was unequal to the demand made upon it. In consequence septic cases occasionally, and suspect cases as a rule, were nursed in the general wards. Precautions were taken to isolate them as much as possible from the other patients.

The following changes were effected on May 1st, 1929. (1) A room was set apart where each patient, on admission, was examined and classified as "clean," "suspect," or "septic." (2) The operating theatre was converted into a labour ward for suspect and septic cases, the labour ward being reserved for clean cases only. (3) The isolation department was increased by 15 beds, making a total of 24, by including one of the general wards in that department. The isolation of suspect and septic cases, their nursing staff, baby room, and annexe, was complete. Improvement was noticed at once.

The terms "septic," "suspect," and "clean" must be clearly defined.

Septic.—This term is too limited in its application, for

Conclusions.

In the foregoing paper I have attempted to make a broad survey of the mechanism of prostatic obstruction and to pick out some of those points which have appeared to me to be of importance. I have tried to show that whilst a dynamic theory may account for the retention of a hydronephrosis, spasm of the vesical sphincter is a contributory rather than a primary factor in the causation of a prostatic retention. The explanation of retention that I have found most satisfying is that there exists a constant mechanical obstruction acting along the lines suggested by Swift Joly which is supplemented by the variable factors of congestion and spasm.

In discussing treatment I have emphasized the importance of treating each type of prostate on its own merits, and have called attention to the value, in selected cases, of such operations as the removal of a middle lobe and of perurethral diathermy.

inflammation may be afebrile and extragenital; therefore such conditions as pyelitis, inflamed breasts, vaginal discharges, bronchitis, discharging sinuses, boils, and all patients admitted with high temperatures, must be regarded as infective.

Suspect.—This term is usually applied to patients admitted in labour who have been examined, or on whom attempts have been made to deliver with forceps before admission; it may be used too loosely. Many patients sent in by doctors have not been examined before admission and are not suspect. The responsibility for their safety rests with the hospital.

Clean.—These patients have no infective process, either local or general. The majority quoted in the results had been admitted from the ante-natal department, but a number had been sent in by doctors. They consist of two groups:

Group A patients are normal medically and obstetrically, and have had the routine vaginal examinations and possibly a forceps delivery or perineal repair.

Group B patients are handicapped by one of the following conditions:

Medical: Disease of the heart, lungs, kidneys, ductless glands.

Surgical: Loss of blood, ante-partum or post-partum.

Obstetrical: Induction of labour, internal version, plugging, manual removal of placenta.

Anatomical: Slight degrees of pelvic contraction.

Group A cases are the most reliable, and *Group B* cases the most sensitive index of the efficiency of a hospital routine.

A study of cases which are already suspect or septic has very little value in such an investigation. The following standards were established before analysing the results.

1. Cases in Groups A and B were delivered in hospital after the twenty-eighth week of pregnancy, and, with only a few exceptions, at term.

2. Only notifiable temperatures were recorded—for example, 100.4° F., sustained for twenty-four hours, or occurring twice in twenty-four hours (excluding the day of delivery). Temperatures were recorded up to the twenty-first day if residence in hospital was prolonged after the routine tenth day.

3. The diagnosis of sepsis was made either by direct clinical evidence or by a process of exclusion; thus all temperatures not definitely diagnosed as due to some extragenital lesion were attributed to sepsis.

The accompanying charts show the incidence of septic temperatures for Groups A and B, independently and combined. The results are based on two-monthly records, the average number of cases in each record being as follows. In 1928: Group A, 167; Group B, 37. In 1929: Group A, 173; Group B, 36. The percentage sepsis among the two groups during 1928 is shown in Charts 1 and 2. On May

CHART 1.—Percentage sepsis among cases in Groups A and B during 1928.

CHART 2.—Percentage sepsis among Groups A and B combined during 1928.

1st, 1929, the new routine was put into operation. This included, in addition to the foregoing, the immediate isolation of any hitherto "clean" case which showed the least evidence of temperature or genital or extragenital inflammatory process. The percentage sepsis for Groups A and B during 1929 is given in Charts 3 and 4. In Chart 5 is shown a comparison of Groups A and B combined

CHART 3.—Percentage sepsis among cases in Groups A and B during 1929.

CHART 4.—Percentage sepsis among Groups A and B combined during 1929.

CHART 5.—Comparison of incidence of sepsis among Groups A and B combined during the periods May-December, 1928 and 1929.

for the periods May to December, 1928, and May to December, 1929.

The beneficial results can be given numerically as follows: for the periods May to December, 1928 (old routine), and May to December, 1929 (new routine).

	1928.		1929.	
	Total.	Sepsis.	Total.	Sepsis.
Group A cases	679	24 = 3.5%	657	6 = 0.9%
Group B cases	155	23 = 14.8%	135	7 = 5.2%
Group A and B combined ...	834	47 = 5.6%	792	13 = 1.6%

Suspect Cases.—It is obvious that "clean" cases benefit by being strictly isolated from suspect and septic cases; the latter two also benefit by getting more intensive supervision and nursing. This is proved by the observations recorded in Charts 6 and 7, which show the incidence of septic temperatures among the suspect cases. Analysis of a series of typical suspect cases, the majority being sent in by doctors, affords further proof of this contention. These included cases of accidental haemorrhage, placenta praevia, contracted pelvis, failed forceps, prolonged

CHART 6.—Percentage sepsis among suspect cases during 1928 (continuous line) and 1929 (broken line).

labour, and malpresentations. These showed improvement under the new routine as follows:

	Total.	Sepsis.
May-December, 1928	209	26 = 12.4%
May-December, 1929	204	14 = 6.9%

It is interesting to note that a new routine was established in the extern department. Prior to July 1st, 1929, nurses, irrespective of their hospital duties, were called to attend district cases. From July 1st, 1929, a special district staff was organized, under the direction of a district sister. The results as regards notifiable pyrexias believed due to sepsis were:

	Total.	Sepsis.
July-August	113	4 cases.
September-October	114	1 case.
November-December	99	0 case.

Conclusions.

1. Outbreaks of pyrexia and sepsis in maternity hospital practice are due to contagion.

2. These can be prevented by (in addition to ante-natal care) the following:

(a) Examination and classification of cases on admission to hospital.

(b) Rigid isolation and intensive nursing of suspect and septic cases.

(c) Immediate isolation of any "clean" case which shows the least evidence of a genital or extragenital inflammatory process.

(d) Provision of a separate nursing staff for the extern department.

THROAT CARRIERS OF THE HAEMOLYTIC STREPTOCOCCUS.

The precautions regarding the prevention of contagion had been in force for two months when this second investigation was started. This was fortunate, since otherwise any observations made would have been most misleading and inconclusive. From July 1st to October 31st, 1929, some general observations were made which helped to decide the way in which the matter could be most effectively investigated.

1. Cultures were taken from the throats of residents, students, or nurses who had delivered "clean" cases. One nurse in the labour ward and two students were reported positive, and were at once suspended, although the nurse was allowed to do dressings, provided she wore a mask. A further series of positive reports on the indoor staff, and the fact that the entire outdoor staff were reported positive, made it clear that wholesale suspensions would entail either closing the hospital or seriously curtailing its work. It was also obvious that the wearing of masks by the district staff would produce alarm and arouse comment. It was therefore decided not to wear masks, but to take the following precautions: (a) all carriers were warned against coughing on their hands, instruments, dressings, or patients; (b) they gargled with boric lotion thrice daily, and their throats were painted once daily with 90 grains of resorcin in 1 ounce of glycerin.

2. The result of this treatment was fairly satisfactory. After five days treatment was suspended for two days, and a second culture was taken. If this was positive treatment was resumed, while if it was negative a confirmatory report was made after fourteen days. The second and third reports were usually negative; several, more resistant, were positive at the second report, had further treatment, and were negative at the third, while in two cases the reports were persistently positive, in spite of a continuance of treatment.

3. A patient developed a septic dermatitis on the buttocks. The student who delivered her was positive and had a bad throat. The nurse in attendance was also positive. Cultures taken from the pustules showed the presence of *non-haemolytic streptococci*. The patient's throat was negative, and she had no sign of genital sepsis. The student may have been the infecting agent, but the fact that the patient was an epileptic suggested other possible means of infection.

4. False deductions may be made unless every possible source of error is eliminated. A nurse (positive) was associated with a series of cases of non-notifiable temperature and mild genital sepsis. Certain of her colleagues who were interested in these observations indicated privately that it was not her throat but her careless technique which was responsible for the bad results.

5. One nurse, known to be positive, developed an acute throat, and the culture which was taken at that time was negative.

6. The possibility of attendance on septic cases being the means of infecting a nurse's throat was considered. Two nurses who had been in constant attendance in the isolation department for six months were negative, and the sister in charge, who entered the department positive, was after three and a half months' residence found to be negative. In neither instance was antiseptic treatment of the throat carried out.

CHART 7.—Comparison of incidence of sepsis among suspect cases during the periods May-December, 1928 and 1929.

7. These preliminary observations showed that throat carriers of haemolytic streptococci are extremely common, and that the organism in that locality is very susceptible to treatment and other hazards of its existence. This last is proved by a later observation, which showed that fifteen nurses reported positive in November were negative in January without any local treatment being used, while six nurses reported negative in October were positive in January.

The known prevalence of the organism, and the rarity and trivial nature of the temperatures which occurred among the clean cases following the established precautions regarding contagion, made it clear that a conclusive test must be made which would leave no vestige of doubt regarding the reputed danger of throat carriers to parturient women.

There were two weak links in the chain of evidence to date: (1) positive carriers had been submitted to an antiseptic throat routine; and (2) only those who delivered the patients had been examined—not the dressers.

To make the test conclusive it was necessary: (1) That all the nursing staff should be examined. (2) That all antiseptic treatment of the throat should be discontinued. (3) That only clean cases, Groups A and B, should be investigated. (4) That no masks should be worn.

The test lasted from November 1st to December 31st. The entire staff—doctors, students, and nurses—were examined. Of the 50 persons from whom swabs were taken, 34 (68 per cent.) were positive.

A record was kept of the work done by each person reported to be positive. This was easy as regards deliveries, but it was a little difficult to keep a complete list of all dressings done by the nurses. The officially recorded number of patients dressed by "carriers" is certainly an underestimate.

Clean cases were delivered or dressed by "carriers" as follows:

Deliveries.—Hospital: total deliveries 183; by carriers 102. District: total deliveries 99; by carriers 47. The district sister-midwife (positive) assisted at nineteen of these deliveries, while the district medical officer (positive) delivered with forceps, or repaired lacerations, on twenty-six occasions.

Dressings.—Hospital and district 478. This number represents individual patients dressed by carriers, but as only 282 patients are under review, it will be clear that a number of patients were dressed by more than one carrier.

Results—Indoor.

Chart 4 shows that the incidence of sepsis in hospital during November and December was the same as during the previous four months. The detailed statistics are as follows:

Group A cases:				Sepsis.
September–October, 1929	...	...	...	1.1 %
November–December "	...	...	...	2.1 %
Group B cases:				
September–October, 1929	...	...	...	2.8 %
November–December "	...	...	...	nil
Combined Groups A and B:				
September–October, 1929	...	...	...	1.4 %
November–December "	...	...	...	1.6 %

The three cases of sepsis among the 183 clean cases delivered in hospital were analysed. All three were trifling, and the temperatures were notifiable on 1, 1, and 2 days respectively.

Case 1.—One day's pyrexia. Delivered by a positive and dressed by negative.

Case 2.—One day's pyrexia. Delivered by a negative and dressed by two positives.

Case 3.—Two days' pyrexia. Delivered by a negative; dressed by two negatives.

Throat and vaginal cultures were taken from the patients in Cases 1 and 3. It is significant that all were negative except the vaginal culture in Case 3, which was positive.

The total district deliveries for September–October, 1929, numbered 114; sepsis 0.8 per cent. The total for November–December was 99; sepsis nil.

It will be seen that the excellent record resulting from the precautions regarding contagion was not in any way affected by this exposure to presumed gross potential infection.

The suspect department afforded a certain amount of material for this investigation. During November and December 19 suspect cases were delivered by carriers and showed no ill effects.

Conclusions.

It is possible that an individual with an acutely septic throat may cause puerperal sepsis by coughing on his or her hands, dressings, instruments, or patients. Any attempt to obtain positive proof of this would be on a par with observations regarding the results of vaginal examinations made with fingers suffering from whitlows, or of surgeons with septic throats coughing into abdomens at laparotomies.

The sum total of the combined investigations indicate:

1. That outbreaks of fever or definite sepsis in maternity hospital practice are due to contagion, and are preventable.

2. That throat carriers of the *Streptococcus haemolyticus* are very prevalent, and that apart from a negation of surgical and aesthetic principles—for example, coughing on hands, instruments, dressings, or patients—they are not a danger to parturient women.

In carrying out these observations special thanks are due to the matron, Miss Ferlie, and her nursing staff, and to Drs. Ind, Cover, and Blacklock for their valuable and most willing assistance.

THE MECHANISM OF REFERRED PAIN IN DYSPEPSIA:

ITS REFLECTION ON THE RESPIRATORY AND
CIRCULATORY SYSTEMS.*

BY

A. T. TODD, M.B.ED., M.R.C.P.LOND.,

HONORARY ASSISTANT PHYSICIAN, BRISTOL ROYAL INFIRMARY.

THE term "dyspepsia" is here employed literally to mean a group of symptoms associated with abnormal digestion produced by any cause. Pain is a symptom in many cases, by no means in all. Though the cause of the dyspepsia may vary within rather wide limits the pains tend to conform to a few types or mixtures of types. It is well known that the pains of ulcer, inflammation, or ulcerated cancer may strongly resemble each other; as a result the nature of the pain and its site have little or no value in differential diagnosis. The timing of the pain, too, is of limited significance, early and late pain being regarded as due to a higher and lower lesion respectively. This similarity of pain brought about by so very different lesions raises the thought that the nature of gastric pain is not understood.

There has been controversy on the mechanism of gastric and other visceral pain for many years; several theories have been advanced, but no single one has met with general acceptance. This article describes a modification of the theory propounded by the late Sir James Mackenzie.¹ The observations on which it is based have extended over some years, and their accuracy is proved by the finding that the signs can be predicted from the symptoms.

The current theories to explain pain in gastric disease fall under two main heads:

1. *That the pain is felt in the affected viscus.* This is the popular theory; physician and patient speak of a pain in the stomach. The pain is regarded as being due to: (a) the action of hydrochloric or other acid on a sensitive mucosa or exposed submucosa; (b) abnormal contraction of the smooth muscle of the wall of the viscus; (c) stretching of the sensory nerves in the wall of the viscus; (d) stretching of the nerves or muscles of the supporting structures. There is abundant clinical and experimental evidence for each of these being a factor.

2. *That the pain is not felt in the viscus but is referred to some other structure.* It is of interest to us that one of the earlier propounders of this theory was a professor of physiology at the Bristol Medical School,² but his views were never widely known. The main upholders of this theory were Mackenzie and Head. Mackenzie considered that stimuli passed from the stomach via the lower dorsal sympathetic fibres to the corresponding spinal segments;

*The substance of a lecture delivered to the January meeting of the Bristol Medico-Chirurgical Society.