N. H. LINDSAY: FOUR CASES OF FRACTURED PELVIS.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

FIG. 1.—Case I. Symphysis pubis wired. Nothing else to note. The dark shadow to right and above symphysis is due to a drainage tube.

FIG. 2.—Case II. Right acetabulum driven into pelvis. Fracture of right pubic bone below the symphysis.

FIG. 3.—Case III. Fracture of left ischium in three places.

FIG. 4.—Case IV. Double fracture of right pubic bone with upward displacement of right side of pelvis at symphysis pubis and right sacro-iliac joint; also fracture of right ilium adjacent to sacro-iliac joint.

ARNOLD K. HENRY: THE HINGED BONE-GRAFT.

FIG. 1.

FIG. 2a.

FIG. 2b.

FIG. 1.—Child's foot hinge-grafted two and a half years previously for extreme calcaneo-valgus deformity due to paralysis. The arrow points to the hinge in the lateral graft.

FIG. 2a.—Adult foot hinge-grafted fourteen months previously for "dangle-foot." The arrow points to the hinge in the medial graft. The position of the lateral hinge is shown by the broken line.

FIG. 2b.—The same foot, seen from the side, showing the strong development of the two grafts. The hinges are not seen.

CASE 4.

A boy, aged 17. Sudden onset in July, 1929, of left renal pain, with temperature 101°. Albumin, blood, and pus were present in the urine. Culture showed staphylococci. Blood urea 28 mg. The urine from the right ureter was normal; that from the left contained pus cells. No tubercle bacilli were seen. The bladder was normal. Indigo-carmin—good concentration from the right in five minutes; none seen from the left in twenty minutes. He was given 40 grams of uroselectan intravenously. Half an hour later a definite picture of the right renal pelvis and calyces was obtained. Fig. 4 shows the appearance after one hour. Exposures were taken at one hour, two hours, and four hours; at no time was any shadow seen on the left side. Cystoscopic pyelography showed a large pyonephrosis on the left side, completely replacing the renal tissue. Fig. 4 agrees very closely with a right pyelogram obtained by the cystoscopic method. It is significant that the uninfected side excreted the drug after half an hour in sufficient quantity to give a shadow, while the infected side failed to show a shadow in any of the radiograms, thus illustrating the depressing effect of sepsis on the renal function.

CASE 5.

A man, aged 28, who for three years has suffered from right renal colic. No frequency; never haematuria. Examination shows tenderness in the right loin only. The urine is straw-coloured, acid, and the specific gravity is 1012; albumin is present. Microscopic examination showed a few pus cells and red blood cells. Cultures were sterile. Blood urea was 31.5 mg. per cent. X-ray examination showed a pyriform oxalate calculus in the right renal pelvis.

After the usual radiological preparation, 40 grams of uroselectan were injected intravenously. Fig. 5 shows the condition a quarter of an hour later. The left renal pelvis and renal outline are shown clearly to be normal. The right pelvis is hydronephrotic, and in the negative the calculus is seen lying within the shadow of the pelvis. The depth of renal tissue and the early appearance of the drug in the right pelvis indicate that the stone has so far caused little interference with excretory function. Pyelolithotomy confirmed the above findings. During the excretion of the drug the specific gravity of the urine rose to 1035, but subsequent examination showed no change in the abnormal constituents already present.

The advantage, in the cases of uninfected calculi, of obtaining all the necessary evidence without recourse to instrumentation, is self-evident.

CASE 6.

A man, aged 30, who had recently undergone excision of a urethral stricture. The cystoscope could not be passed owing to a fold. There was no obstruction to micturition. Intravenous pyelography showed normal renal pelvis half an hour after the administration of the drug.

CASE 7.

This patient had nephrostomy performed in October, 1929, for infected right hydronephrosis and incipient uraemia, the left kidney being congenitally undeveloped. The left kidney excreted only two to three ounces per diem, and concentrated urea only to 0.4 per cent. Nephrostomy reduced the blood urea from 95 mg. per cent. to 44 mg. per cent. A ureteric catheter could not be passed into the right pelvis from the ureter, and a satisfactory pyelogram was not obtained via the nephrostomy tube. The patient was given 40 grams of uroselectan intravenously. An irregular ill-defined shadow began to appear after half an hour, and was more intense in two hours, but no shadow was seen on the left side. Unfortunately the patient (who was an out-patient) could not stay for further x-ray examination.

CASE 8.

A boy, aged 7, who for twelve months has had attacks of symptomless haematuria. Cystoscopy shows no vesical cause for the haematuria. Indigo-carmin—good colour from right in five minutes, poor from left in fifteen. The urine collected from each ureter contains many red blood cells with epithelial cells. A few red blood cell casts were occasionally seen. X-ray examination was negative. Blood urea was 45 mg. per cent. The blood count shows very marked eosinophilia (28 per cent.). A blood Wassermann test was positive (+++); the father's Wassermann reaction was also positive (+++). The stools and urine showed no evidence of paracytic cysts or ova. The condition is considered to be one of chronic syphilitic nephritis, as cystoscopic pyelograms show no abnormality. The patient was given 16 grams of uroselectan in 40 c.cm. of distilled water. A film taken a quarter of an hour later shows a little drug in the bladder only. A film at twenty-five minutes shows the right pelvis outlined, whilst the left pelvis began to show only after an hour. The right pelvis still showed faintly after four hours. The findings were similar to those obtained by cystoscopic methods.

Two normal patients were used as controls—one a youth aged 19, the other a boy aged 12. In each case the pelvis were seen after fifteen minutes.

CASE 9.

Fig. 6 shows the result obtained after thirty minutes in a woman, aged 28, whose right ureter has been transplanted into the rectum for epispadias, with vesical incontinence.

In conclusion we wish to express our thanks to Dr. Wylie, radiologist to the hospital, for the use of his department, and to Miss Kreckeler, our radiographer, for her very able co-operation. We would also like to thank Dr. H. V. Morlock, assistant physician to the hospital, for his helpful interest and advice.

BIBLIOGRAPHY.

- ¹ Roseno, A., and Jekins, H.: *Fortschr. a. d. Geb. d. Röntgenstr.*, May, 1929.
- ² Osborne, Sutherland, Rowntree, and Scholl: Roentgenography of Urinary Tract during Excretion of Sodium Iodide. *Journ. Amer. Med. Assoc.*, 80, February, 1923, 268-276.
- ³ Roseno, A.: Intravenous Pyelographie. *Klin. Woch.*, No. 25, 1929, S.1165.
- ⁴ Howard, R. C.: Intravenous Injections of Sodium Iodide in Large Doses in Obstinate Syphilis. *Amer. Journ. Syph.*, July, 1928, 550.
- ⁵ Da Matta: *Brazil-Med.*, October 14th, 1916.
- ⁶ Ziegler and Köhler: Perorale Pyelographie. *Med. Klinik*, No. 1, 1930, S.10.
- ⁷ Rosenstein and Lichtenstein: *Zentralbl. f. Chir.*, 51, 1169-1170. Pneumoradiography. *Zeit. f. Urolog. Chir.*, 12, March, 1923, 268-298.
- ⁸ Swick: Darstellung der Niere und Harnwege im Röntgenbild durch Uroselectan. *Klin. Woch.*, No. 45, 1929, S.1087.
- ⁹ Von Lichtenberg and Swick: Klinische Prüfung des Uroselectans. *Klin. Woch.*, No. 45, 1929, S.2089.
- ¹⁰ Kielluthner: *Munch. med. Woch.*, No. 7, 1930, S.276.
- ¹¹ Volkmann and Hyntschaks: Zur röntgenographischen Darstellung der Harnwege durch intravenöse Verabreichung schattengebender Mittel. *Deut. med. Woch.*, 50, 2, 1924, S.1413.

Since this paper was written Mr. Frank Kidd has published examples of the use of this drug in the current number of the *British Journal of Urology*.

THE HINGED BONE-GRAFT.

BY

ARNOLD K. HENRY, M.B., B.Ch.DUBL.,
F.R.C.S.I.,

PROFESSOR OF CLINICAL SURGERY, FACULTY OF MEDICINE, CAIRO, AND
DIRECTOR OF THE SURGICAL UNIT, KASR-EL-AINI HOSPITAL, CAIRO.

(With Special Plate.)

THREE years ago I described in this *Journal* a method of splinting unstable feet by means of hinged grafts, which combined stability with a degree of useful movement.¹ The results have been so entirely satisfactory that a further description is desirable.

The grafts are made from osteo-periosteal strips implanted into bony grooves, cut in the back of each malleolus, and extending on either side of the foot to a point just beyond the mid-tarsal articulation. Connivance during convalescence at slight movements of plantar flexion and dorsiflexion of the foot secures the formation of a hinge in each graft just distal to the tip of the malleolus. An internal living splint—like the hinged cage-splint used externally for loose knees—is thus incorporated with the ankle.

I have employed the method in two kinds of case: (1) for patients with calcaneo-valgus deformity due to paralysis; (2) for complete foot-drop or "dangle-foot." Before grafting, any fixed deformity in the paralysed foot must, of course, be corrected by manipulation or tenotomy.

Paralytic Calcaneo-valgus Deformity.

Apart from hinge-grafting, the operations devised to cure this deformity have either failed to stand the test of time or have spoilt the shapeliness of the part. A tenodesis will stretch; transplanted tendons are inadequate, and astraglectomies make unlovely ankles. Fig. 1 is an x-ray picture of a foot hinge-grafted two and a half years previously for an extreme calcaneo-valgus deformity. The girl, when shown in the Surgical Section of the Congress held during the centenary celebrations of the Cairo School of Medicine, could stand and hop on the grafted foot, and could move it in a vertical plane. There was no lateral instability of any kind. (One of the two hinges is clearly seen in the radiograph.) A second patient was operated on shortly after the first, with a satisfactory result.*

"Dangle-foot."

Many operations have been devised for this disabling condition. Some are unsatisfactory; Jones and Lovett regard arthrodesis as a method "yielding at best but an

* A third patient, but with equinus instead of calcaneus deformity, has recently been hinge-grafted after correcting the fixed equinus component by tenotomy.

imperfect functional result," and "contraindicated in young children." The end-results, too, of tenodesis that I have seen have been mediocre, while other procedures are either mutilating or less simple of performance than the operation of hinge-grafting.

Makers of artificial limbs realize the disadvantage of a foot fixed immovably to the leg, and they provide limbs with a slight degree of movement at the ankle-joint. I believe that some mobility is equally desirable in the living foot, and that while bony fixation at right angles to the leg is good, a stable foot with a few degrees of movement in the vertical plane is better. This can easily be got, in the most unstable feet, by means of hinged grafts.

Figs. 2a and 2b show x-ray pictures of a foot hinge-grafted fourteen months ago; the whole lower limb was paralysed by a stab wound of the cauda equina. The foot is at a right angle to the leg, and the hinge grafts give a range of about 5 degrees of movement in the vertical plane, which allows the leg to rock naturally on the foot when the patient walks. The rest of the limb is stabilized by means of a skeleton splint. It is particularly important in "dangle-foot" to bring the grafts across the mid-tarsal joint.

The Operation.

The following points are worth noting.

The Position of the Patient for Operation.—The prone position is best; this gives access to the backs of both malleoli, and allows the graft to be cut from the opposite tibia after flexing the knee (Fig. 3).

FIG. 3.

Skin Incision.—Time may be saved by making the three skin incisions—one extending to the foot from behind each malleolus, and one over the opposite tibia—in succession at the beginning of the operation. The scalpel is then discarded, and a second scalpel is used for the deep dissections. Side-curtains are fixed to the neighbouring wounds behind the ankle by means of Michel clips, to avoid a double encumbrance of towel forceps.

The Graft Bed.—Medially the tibialis posterior, and laterally the peronei, are displaced from their grooves, which are deepened by a gouge to form beds for the grafts; each groove must cross the mid-tarsal joint. The grafts are held in position by stitching the tendons over them; four stitches suffice for each graft. I do not permit either my own skin or the patient's to come in contact with the outer surface of my gloves. By this precaution I avoid the elaborate game of skill in which wound, graft, and sutures are treated as untouchable, and time is saved.

The Grafts.—I cut the osteo-periosteal strip, which is outlined by a scalpel, with a chisel of the same width as

FIG. 4.

the subcutaneous face of the tibia—entering the chisel, bevel up, to the required depth, and then using it, bevel down, to shave off the strip (see Fig. 4). Then, holding the strip with tissue forceps by the four corners, I divide it longitudinally with straight scissors into two equal grafts. It does not seem to matter whether the graft has the pliability of wet paper or the spring of a hazel twig; the result is the same—a hinge, just distal to the malleolus, is ultimately formed in either case. The patient whose

radiograph is shown in Fig. 2a, where the hinge is clearly seen, was grafted with tough, springy grafts, while in the girl's foot (Fig. 1) the grafts were flaccid.

Skin Closure.—Time is again saved, and contamination from skin is avoided by postponing the closure of all three wounds till the end of the operation. Michel clips are used instead of sutures, since the clips provide no path for infection. Rutherford Morison's invaluable thick layer of sterile boric powder is applied as a dressing.

After-Treatment.

The foot is bandaged for a week at right angles to the leg in an ordinary soft bandage; this permits the small degree of movement that produces the hinge in each graft. The Michel clips are then removed and the bandage reapplied. If there is any tendency to foot-drop or lateral deviation, or if the patient is unlikely to co-operate by remaining in bed, the limb is fastened to a long back splint, with the foot at right angles to the leg. A month after operation a plaster casing that extends above the knee replaces the bandage or the splint, and holds the foot in the same position. The patient is made to stand beside his bed on the plaster casing for increasing periods of time. The casing is removed two months after the date of operation, and the patient is encouraged to walk.

The Advantages of Hinge-Grafting.

1. The stability given by these bony hinge grafts improves with use; they do not stretch like tendons that are fixed or transplanted.
2. The after-treatment is simple: final and satisfactory function is obtained within three months—a time which compares favourably with the long period of care and re-education required after transplanting tendons.
3. If relevant muscles can still contract, the hinges allow movements of dorsiflexion and plantar flexion.
4. Hinge-grafting does not—like astragalectomy—disfigure the ankle.
5. It gives excellent results in "dangle-foot."
6. For talipes calcaneo-valgus it is the method of choice—not, I submit, another poor best among a disheartening array of forlorn hopes; but a thoroughly satisfactory treatment of the deformity.

REFERENCE.

Henry, A. K.: *British Medical Journal*, April 9th, 1927, p. 663.

FOUR CASES OF FRACTURED PELVIS.

BY

N. H. LINDSAY, M.B., B.Ch., B.A.O.,

HOUSE-SURGEON, ROYAL UNITED HOSPITAL, BATH.

(With Special Plate.)

THE interesting features about the following cases are the manner in which the lesions were produced, illustrating the different types of fracture caused by varying types of violence, and the fact that all four patients were admitted to hospital within a fortnight of each other.

CASE I.

T. D., a railway guard, aged 48, was admitted to hospital on January 17th, 1930, with a fracture through the symphysis pubis with rupture of the bladder. It was an incomplete fracture produced by compression violence. The patient fell from a station platform, and was caught between the platform and the running board of a moving train.

The following symptoms were exhibited. Extreme pain on compressing the iliac crests; inability to move the lower limbs without causing severe pain, but no evidence of nerve lesion; urgent desire for micturition; passage of a catheter withdrew a few drops of blood-stained urine.

The same day, after x-ray examination, the bladder was explored and the pubic symphysis exposed. A linear tear, a quarter to half an inch in length, at the junction of the anterior surface of the neck of the bladder and urethra was found and closed. The left superior ramus of the pubis was displaced downwards and backwards, being separated from the right superior ramus by about two inches. The displacement was corrected by manipulation and the fragments wired in position. A catheter was passed and tied in. Micturition was normal within forty-eight hours.

Further treatment consisted in the application of a stout pelvic binder and nursing on fracture boards.