the muscle by a paroxysm of coughing. Palpation of the lower abdomen during such a paroxysm will demonstrate a surprisingly sustained and continuous contraction of the rectus muscle, particularly in its lower half. The actual strain of the muscle is not itself productive of the haemorrhage.

With regard to the diagnosis of such cases, stress must be laid on the symptoms of peritoneal irritation that occur. Such symptoms have been noted in most of the recorded cases, and were present in both of the cases just described, leading to the initial diagnosis for each of an acute intra-abdominal lesion. The ecchymosis of the external genitals in either case proved a valuable indication of the nature of the swelling within the rectus sheath, forming as it did quite soon after the first appearance of the haematoma. The chief value of exploration of the swelling would seem to be the establishment of the diagnosis, which must in many cases be doubtful without it. Unless the swelling is very large or some complications arise, there is little to be done at operation.

Summary.

Two further cases of spontaneous haematoma of the rectus abdominis muscle are described, and the etiology and diagnosis of such haematomas are discussed. Their relationship to systemic conditions, and in particular to the haemorrhagic diathesis, is emphasized.

REFERENCES.

- ¹ Andrews, H. R.: *Proc. Roy. Soc. Med.*, Obstet. and Gynaecol. Sec., 1915-16, 9, 98.
- ² Cole, C. E. C.: *British Medical Journal*, 1919, ii, 566.
- ³ Culbertson, C. J.: *Journ. Amer. Med. Assoc.*, 1925, lxxxv, 1955.
- ⁴ Fothergill, W. E.: *British Medical Journal* 1926, i, 941.
- ⁵ Grignani: *Boll. di Soc. Med.-Chir.*, Pavia, 1926, 901.
- ⁶ Halperin, G.: *Surg., Gynecol. and Obstet.*, 1928, xlviii, 6, 851.
- ⁷ Perman, E.: *Acta Chir. Scand.*, 1922, 54, 434.
- ⁸ Stoeckel, W.: *Zentralbl. f. Gynäk.*, 1901, x, 241.

MYOSITIS FIBROSA PROGRESSIVA.

BY

I. I. PRICE, M.B., B.S.LOND., F.R.C.S.ENG.,
MEDICAL SUPERINTENDENT, TILBURY SEAMEN'S HOSPITAL.

MYOSITIS FIBROSA can be defined as a disease in which the various muscles of the body undergo a fibrous change. The literature of the disease is very sparse, and it is not mentioned in Boyd's *Surgical Pathology*. The condition is very rare; it is said to be more common in females than in males, and most of the cases reported have been in children.

Two types have been described: the localized and the generalized.

The Localized Type.—In this type one or more muscles have undergone a fibrous change, and the patient has not suffered any considerable disability. Deformities such as torticollis and talipes have been an end-result, but these deformities have not been severe.

The Generalized Type.—Here many muscles are involved. The main part of the musculature of the body may be transformed into fibrous masses which may be so disabling as to cripple the patient seriously or render him an easy prey to pneumonia. The disease apparently attacks the voluntary muscles only, those of the back and upper limbs being the most commonly affected.

Clinically the disease is indicated by the appearance of swelling in a part or in the whole of a muscle. This swelling occurs generally near one of the attachments of the muscle. It is hard, diffuse, and painless; and although within a few days it decreases in size and feels harder, there is left a permanently indurated area in the muscle concerned. Sometimes a swelling which is decreasing in size suddenly increases again and then decreases. This change in the muscle corresponds to a formation of fibrous tissue in the musculature, and is unaccompanied by any rise of temperature or any feeling of malaise. Examination of the blood reveals nothing abnormal, and the Wassermann reaction is negative. Microscopically there is a general increase of fibrous tissue among the muscle fibres. The fibrous tissue appears to surround and invade the muscle fibres, which degenerate and disappear. Round-celled infiltration is noticeably absent. The fibrous tissue appears to be well supplied with blood vessels.

The cause of the disease is unknown, and treatment has little effect on its progress. In some cases massage and electrical treatment has produced some improvement, but the most that such treatment can be expected to accomplish is to prevent deformities from occurring as a result of the contraction of the fibrous tissues.

Myositis fibrosa can be compared with the following diseases causing a change in muscles: (a) myositis ossificans progressiva, (b) so-called congenital torticollis, (c) Volkmann's ischaemic contracture, (d) true chronic metritis, and (e) acute polymyositis. Such conditions as tuberculous, hydatid, pyogenic, syphilitic, and others of known pathology affecting muscles will not be considered.

Myositis ossificans progressiva bears a very close resemblance to myositis fibrosa. In both conditions muscle undergoes degeneration. In one the muscle is replaced by fibrous tissue; in the other the increased formation of connective tissue is followed by a deposit of bone in the muscle. It would appear at first sight that myositis ossificans is but a later stage of myositis fibrosa; this, however, is not borne out by clinical evidence. Stunting of the big toe has been noted in myositis ossificans progressiva, but this is absent in myositis fibrosa. Probably both diseases are a result of an inherent degenerative process of muscle which, in one case, stops short at the connective tissue stage, and in the other goes on to bone formation.

Both Volkmann's ischaemic contracture and so-called congenital torticollis have been regarded as due to vascular obstruction causing a degenerative fibrous process in muscle which finally ends in contracture, but apart from the pathological condition they appear to have no relationship with myositis fibrosa. It is curious that in the case reported in this article torticollis occurred at one stage and gave a clue to the diagnosis.

Fletcher Shaw has described three types of chronic metritis, one of which he calls "true chronic metritis." In this condition the main part of the uterine musculature is replaced by fibrous tissue, and the condition has been regarded as a result of chronic nephritis or arterio-sclerosis. It appears that the disease is due to some unknown toxin, and the toxin theory might explain the process undergone by the muscles in myositis fibrosa.

In acute polymyositis there is an acute inflammatory condition resulting in degeneration of the muscle fibres, though the responsible toxin is unknown.

The patient whose case is here recorded came under my care at Tilbury Seamen's Hospital.

A girl, aged 3½ years, was admitted as a patient suffering from Pott's disease of the dorsal spine. The child had been perfectly well, but the mother had noticed for about a month a swelling about the size of a five-shilling piece on the right side of the spine in the region of the scapula. This swelling was hard, painless, not attached to the skin, and had no palpable edge. It was movable across the direction of the fibres of the rhomboid muscle but not along them. Movements of the spine were quite free, and the child appeared quite happy. X rays revealed nothing abnormal. No focus of sepsis was discovered. In the lumbar region on the right side there was a similar swelling of a smaller size, apparently involving the erector spinae muscle. Further examination revealed that the movements of the left shoulder were considerably diminished. Abduction was limited to 20 degrees. External and internal rotation were also limited. The scapula was freely movable. The insertions of the muscles around the shoulder-joint felt harder and more tendinous than usual. The Wassermann reaction was negative.

The child was sent to see several surgeons, who all diagnosed the condition as sarcoma. She was watched, and the swellings appeared to enlarge and then to diminish in size; they did not disappear, but left a hard indefinite mass in the muscles involved. Later a swelling appeared in the left sterno-mastoid in its upper quarter and went through the same changes. The child then acquired a left torticollis. A fortnight later the right sterno-mastoid became involved and contraction occurred. The left pectoralis major at its origin then became involved and showed a large swelling about 4 by 5 inches in size. Both shoulder-joints now showed restricted movements. The right brachialis anticus in the region of the elbow-joint became affected, and the condition in this part closely resembled myositis ossificans traumatica. There was limited extension of the right elbow-joint.

A portion of the pectoralis major was excised under anaesthesia and was sent for microscopic examination. Dr. Arthur Davies, pathologist to the Seamen's Hospital, reported as follows:

"Sections show a considerable excess of fibrous tissue replacing and surrounding bundles of muscle fibres. In one part of the

tissue the musculature has been completely replaced by fibrous tissues, prolongations from which spread throughout its musculature. The fibrous tissue is well supplied with blood vessels of considerable size and there is no indication of a true sarcoma."

It is noticeable that the sections do not show any inflammatory process, but suggest rather the appearance of a neoplasm.

Endocrine disturbances have been described as a cause of myositis fibrosa progressiva, but probably some non-specific toxin of intestinal origin is an important factor.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

A CURIOUS COMPLICATION OF MEASLES AND WHOOPING-COUGH.

THERE has been for some time evidence of a changing character in our common infectious diseases and the complications arising. This has been noticeably so in cases in which measles and whooping-cough have occurred in conjunction. The case here recorded, however, presented features I have never encountered before.

A girl, aged 7 years, of poor physique, but previously healthy, developed a cough in March. A typical whoop came on two weeks later, and at the same time the child commenced a severe cold in the head, with a rise of temperature to 101°. As there was a considerable epidemic of measles also about, I warned the parents that an attack of this disease might be beginning. A typical rash developed on the morning of April 9th, and the progress of the case was as follows.

April 10th. Child very ill; temperature 103°; rash fully out; some headache, cough troublesome.

April 11th. Severe headache; temperature 103°-104°; rash fading on face.

April 12th. Pain in head intolerable; child delirious; became unconscious in the evening; temperature 104°; pulse 140.

April 13th. Child unconscious; in the morning she ceased to move her left leg, and by evening her left arm was also to all appearances paralysed. On the left side the knee-jerk was exaggerated, and ankle clonus was present. There was no definite Kernig's sign, and no tache cérébrale.

April 14th. Temperature 100°; pulse 110; child seemed slightly conscious, but appeared, to my horror, to be blind.

April 15th. Child conscious, but blind; can move left leg, but not arm; temperature 99°; pulse 100.

April 17th. Child can now move left arm very slightly and leg better. Complaints of severe headache, but can distinguish between light and dark; can ask for the bedpan, and is sleeping for two or three hours at a stretch. The ankle clonus is gone, but the knee-jerk is still more marked than on the right side.

From this stage the child made a gradual recovery. The eye-sight, after considerable photophobia, became normal in a fortnight. The leg first, and then the arm, entirely recovered. The girl is now (June 15th) perfectly normal; she is unusually bright mentally (which, incidentally, I had warned the parents might not be the case), and has had no ill health since.

Dr. Edmondson of Lancaster, who saw the patient with me, agreed that it was obviously a case of cerebral irritation, and suggested the possibility of the cause being a secondary rash on the meningeal membrane.

The treatment, on her return to consciousness, was simply urotropine, 5 grains every four hours.

Bentham.

J. F. Dow, M.D. Vict.

Reports of Societies.

PSYCHIATRIC LESSONS FROM AMERICA.

AT the quarterly meeting of the Royal Medico-Psychological Association held at the House of the British Medical Association on May 22nd, with the president, Dr. NATHAN RAW, in the chair, a discussion took place on American psychiatry and its possible bearing on the application of recent local government and mental treatment legislation.

Dr. A. A. W. PETRIE, who opened the discussion, gave an account of a recent visit to American general and mental hospitals, psychiatric clinics, reception units, observation wards, and mental deficiency institutions, as well as to mental hygiene centres and various forms of child guidance organizations and associated nursery schools. Observations were also made of psychiatric work in connexion with juvenile courts. Each State, said Dr.

Petrie, had its own method of dealing with the problems involved. In the progressive Eastern States it was usual for the mental hospital service to be a subsection of the state welfare services under medical commissioners. Concentration of executive authority in the State rendered progressive development easier, and made it more possible to obtain legislative and other changes. Well-organized State social services were being developed in New York and New England. The larger unit provided by a State allowed a co-ordination of activities that was not possible in smaller units. New York State, for instance, dealt with over 63,000 patients, who were drawn from a population of 44,000,000. The recent Local Government Amendment Act, however, did make one authority in an administrative area responsible for the treatment of the earliest stages of mental disorder. Well co-ordinated and efficient reception hospitals should take the place of the present wasteful division into observation wards of doubtful efficiency. Dr. Petrie also hoped that the Mental Treatment Bill when it became law would allow the development of voluntary clinics on the same lines as in the Maudsley Hospital and the famous Phipps Clinic in Baltimore, which consisted of sixty beds and an out-patient department, education and research being prominent features. Notable also was the clinic at Philadelphia, a hospital somewhat resembling Bethlem in London. The Psychiatric Institute which had been established in New York in connexion with the medical unit of Columbia University would be one of the most notable in the world when fully developed. Reception hospitals were increasing in number, which was one of the most hopeful signs in the United States. He doubted whether such clinics and reception hospitals should be an actual part of the hospital to which they were attached. The out-patient department at the Boston Psychopathic Hospital, which was the clearing house for Boston area, attracted as many patients as the corresponding out-patient department at Boston General Hospital. The work of these reception units was largely determined by the source of the admissions. In the Chicago clinic, with its 175 beds, about 100 beds had to be cleared weekly in order to deal with new admissions. These units had not usually sufficiently large staffs to deal with the great amount of material. A unit with 600 beds was projected in New York. The chronic patients in the large State hospitals in America were not any better dealt with, in Dr. Petrie's opinion, than they were in England; in America overcrowding occurred at many places. Much difference of opinion existed in America as to what was the best size of mental hospital. In Massachusetts, a very progressive State, 2,000 patients was the usual limit. He found in the United States a tendency, similar to that in England, to modernize old hospitals by building new and up-to-date units for recently admitted mental patients, and infirmary blocks for those suffering from physical diseases. He also saw good examples of the villa or colony hospital. Consultants were usually appointed on a liberal scale in American mental hospitals. In the progressive States the medical staff was in larger proportion to the number of patients than in England; in Massachusetts there was one assistant physician to 150 patients. A useful appointment, common in the States, was that of clinical director or clinical psychiatrist, who was equal in rank to the deputy medical superintendent and the pathologist. He directed and co-ordinated treatment, and was usually a teacher in a medical school.

The laboratory facilities were similar to those in England; frequently there was a more elaborate equipment. Lectures to psychological and theological students and social workers in general were encouraged in the States. Much could be done in this way to break down the prejudice and mystery which were generally associated with mental hospital work. A sharp division existed between nurses and attendants, based upon the preliminary education required. Attendants had no special training. The line of cleavage would become more marked in the future when a complete secondary education for nurses was demanded. Occupational therapy, said Dr. Petrie, was more broadly developed in a general way in the States than in this country. At St. Elizabeth's Hospital, Washington, sixteen occupational therapy officers were engaged