Other factors than dysfunction or disease of uterus and ovaries are responsible for some cases of uterine bleeding. Cases occur in which it is to be attributed to a pathological condition of the blood.

The mechanism of haemorrhage in these cases is not clear, but it may depend on disordered calcium metabolism—for example, increased calcium output or low blood calcium.

In cases of severe uterine haemorrhage without obvious pelvic or local cause, a systemic investigation and a detailed blood examination (preferably by a haematologist) should be made at the earliest opportunity.

I am indebted to the surgeons of the Royal Samaritan Hospital for Women, Glasgow (Drs. Shannon, West, Gardner, and McIntyre), and to the medical superintendents of the Glasgow Royal and Victoria Infirmarys (Drs. Grant and McCall Smith respectively), for access to records and notes in some of these cases.

REFERENCES.

- ¹ Shaw, W.: *Journ. Obstet. and Gynaecol. of the British Empire*, 1929, i, p. 60.
- ² Schmitz, H.: *Journ. Amer. Med. Assoc.*, September 29th, 1928, p. 955.
- ³ Linton, Marks, and Smith: *Ibid.*, March 23rd, 1929, p. 966.
- ⁴ Munro Kerr, J. M.: *Clinical and Operative Gynaecology*, London, 1922, p. 87.
- ⁵ Osler and McCrae: *Principles and Practice of Medicine*, D. Appleton and Co., 1920, pp. 748, 704.
- ⁶ Meigs, J. V.: *Amer. Journ. Obstet. and Gynecol.*, 1927, xiv, p. 225.
- ⁷ Eden and Lockyer: *Gynaecology*, London, 1920, p. 133.
- ⁸ Price, F. W.: *Text Book of Practice of Medicine*, Oxford Press, 1929, p. 728.
- ⁹ Berkeley, Comyns (and others): *Diseases of Women* (by Ten Teachers), London, 1926, p. 73.
- ¹⁰ Eden and Lockyer: *Loc. cit.*, pp. 426, 430.

PRODUCTION OF PERNICIOUS ANAEMIA (BARTONELLA ANAEMIA) IN RATS BY DEFICIENT FEEDING.*

BY

LUCY WILLS, M.A.CANTAB., M.B., B.S.LOND.,

AND

MANECK M. MEHTA, M.A.BOMBAY, PH.D., D.SC.LOND.

(From the Nutritional Research Laboratories, I.R.F.A., Coonoor, South India.)

THE dietetic survey of various classes of women in Bombay suggested that the disease occurring in pregnant women known as "pernicious anaemia of pregnancy" was closely connected with a relative deficiency of vitamins A and C in their diet. In this connexion a series of experiments was carried out at Coonoor in which diets relatively deficient in these vitamins led to the production of a severe anaemia in intact albino rats, especially in the female animals; it was further shown to be a true "Bartonella anaemia." The term "Bartonella anaemia" is used to describe the well-known anaemia of rats which occurs after splenectomy in infected stock, and is associated with the appearance of Bartonella "organisms" in the red cells; it is used without prejudice as to the exact nature of these small bodies.

DIETETIC EXPERIMENTS.

The diets used had one common fault—a relative deficiency in vitamins A and C.

Series I.

These experiments have been reported before by McCarrison,¹ in his work on the production of urinary calculus by faulty feeding. In them two types of diet were used, the one based largely on oatmeal, and the other on atta (whole wheat flour). These diets contained no vitamin C and only a little vitamin A, the latter being obtained from the small quantity present in the grains used. Vitamin B was judged to be present in adequate amounts.

Cases of a severe, and frequently fatal, anaemia occurred among the rats fed on these diets; among those fed on the oatmeal diets there was a death rate from this disease of 13.8 per cent.; among those fed on the atta diet there was a death rate of 12.4 per cent. Milk added to the oatmeal diet appeared to afford the rats protection against this anaemia. The disease was much more frequent in

females; out of a total of seven pregnant animals in one experiment five died of this anaemia.

The diagnosis of Bartonella anaemia was made on an examination of the blood slides. The characteristic appearances in the blood will be described in the account of the next series of experiments; in these the blood examination was much more complete.

Series II.

Six diets, two natural and four synthetic, were used in this series of experiments. The two natural ones were based on diets in common use among the hospital class in Bombay; one resembling that used by Mohammedan women and the other based on a diet used by Hindu women. Of the four synthetic diets the first was a basic one without vitamins A and C except such traces of vitamin A as might be found in the olive oil used (very

FIG. 1.—Blood from splenectomized anaemic rat showing "Bartonella organisms" in the red cells. (×1,000.)

little if any present, as the animals went downhill rapidly); the others were the same as the first with the addition of orange juice to the second, cod-liver oil to the third, and both orange juice and cod-liver oil to the fourth.

Cases of Bartonella anaemia occurred spontaneously among the rats fed on the Mohammedan diet and among those fed on the basal synthetic diet. Pregnant animals living on the Hindu diet were, however, susceptible to inoculations of Bartonella-infected blood; in this respect they were unique, as we failed to produce an anaemia after inoculation either in well-fed rats, whether pregnant or not, or in pregnant rats when they were fed on the Hindu diet to which vitamins A and C were added in the form of cod-liver oil and tomato juice.

The amount of vitamin B and salts and also of fat varied in the different diets. The Mohammedan diet, on which the animals developed the anaemia most rapidly, was markedly deficient in vitamin B (poor growth from the beginning of the experiment) and salts, and at the same time had a high content of vitamin-A-free fat. The Hindu diet, on the other hand, had more vitamin B and a very low fat content. The synthetic diets had what has been shown to be an adequate amount of vitamin B and salts. The animals on complete synthetic diet grew well. The common factor in the diets which led to the production of the anaemia was, therefore, a relative deficiency of vitamins A and C, or very possibly some other dietetic fault associated with this deficiency.

* Summary of a paper appearing in the *Indian Journal of Medical Research*.

The anaemia was severe, as shown by the extreme pallor of all the organs and by the blood counts. Haemoglobinuria was common. The most striking post-mortem feature, apart from the general bloodlessness, was the fatty, mottled liver. This organ appeared to be enlarged; but when its weight was charted against body weight it was found that the organs obtained both from intact and from splenectomized rats fell within normal limits. In cases exhibiting haemoglobinuria the bladder was distended with dark-coloured urine and the kidneys were large and very dark. The bone marrow showed no macroscopic change.

The blood showed the typical picture of a macrocytic anaemia. The red cell count fell in certain cases from 10 to 1.35 million, and the haemoglobin from 13.2 to 4.9 grams per cent. This fall was associated with a marked degree of anisocytosis; the average diameter of the red cells was increased from 5.9μ to 8.2μ . As the anaemia progressed large numbers of normoblasts appeared in the blood stream. But the finding that clinched the diagnosis of Bartonella anaemia was the presence of the characteristic rod-like bodies in the red cells (Fig. 1).

SPLENECTOMY EXPERIMENTS.

In order to confirm the diagnosis arrived at in the previous experiments, and to show that all the stock and experimental animals were infected with Bartonella, and in addition to test various lines of treatment, a series of representative animals were splenectomized. Sixty-six animals were operated on by Colonel McCarrison. Of these one died of haemorrhage from the pancreas in the first twelve hours; the others survived the operation well. Post-operatively all the animals were fed on the stock diet with the addition of unlimited whole milk; 25 had no further treatment, 17 were treated with liver extract, 12 with fresh spleen, and 11 with vitamin A concentrate (radiostoleum). With a few exceptions all the animals developed the anaemia; liver, spleen, and vitamin A extract afforded no protection against the "infection" in splenectomized rats.

The clinical, pathological, and post-mortem appearances were identical in the splenectomized rats and in those developing the anaemia as a result of faulty feeding, save that the former had no spleens and their blood picture showed an enormous polymorphonuclear neutrophil leucocytosis and a much heavier incidence of red cells "infected" with Bartonella. The leucocytosis is probably a direct result of the splenectomy as it occurs immediately after the operation; but the reason for the number of "infected" red cells being so much greater, although the anaemia is no more severe, is not clear.

INOCULATION AND CULTURAL EXPERIMENTS.

With the exception of the above-mentioned experiments with pregnant animals fed on a Hindu diet all attempts to produce anaemia by the inoculation of infected blood into intact animals were unsuccessful. We were also unable to confirm Ford's² work with young rats. All attempts to culture the "organism" failed.

DISCUSSION.

The interest of these experiments lies in the fact that it has been possible by dietetic means alone to produce a severe Bartonella anaemia in intact animals, the anaemia being of the same order as that occurring after splenectomy in infected rats. The diets used have one common fault—a relative deficiency in vitamins A and C. There were many other experimental animals in close contact with those which developed the disease, but although all were infected, as shown by the splenectomy experiments, only those fed on diets with this dual deficiency developed the anaemia. It seems safe to conclude, therefore, that the factor that renders intact rats liable to this anaemia is a dietary one. The present series of experiments is not sufficiently conclusive to determine the exact nature of the deficiency; the results suggest that either a relative deficiency of vitamins A and C or of some other closely associated dietetic factor, perhaps a specific anti-anaemia one as found in liver, is most likely to be responsible. The presence of the characteristic "organism" in the red cells clinches the diagnosis of Bartonella anaemia.

It is suggested that the deficiency, whatever be its exact nature, may have some effect on the reticulo-endothelial system comparable to that of vitamin A deficiency on epithelial structures; and that it is by virtue of this action that the natural resistance of the intact rat is overcome, even as it is overcome in infected stock by removing a large bulk of this tissue—that is, the spleen. In view of the importance of the reticulo-endothelial system in the defence of the body against infections, particularly protozoal ones, the experiments reported here may not be without significance in connexion with the incidence of these diseases in India, where a large section of the population exists on diets having the faults described above.

SUMMARY.

1. A severe anaemia has been produced in rats by feeding them on diets low in vitamins A and C.
2. The anaemia was more frequent and more severe in females than in males.
3. Blood examination revealed the presence of *Bartonella muris* in the red cells of the anaemic rats.
4. Splenectomy experiments showed that all the Coonoor rats were infected with Bartonella.
5. Liver, spleen, and vitamin A concentrate failed to protect the rats from this anaemia after splenectomy.

We wish to thank Colonel McCarrison for placing the material from his earlier experiments at our disposal, and for his generous help in this work.

REFERENCES.

- ¹ McCarrison, R.: *Indian Journ. Med. Research*, 1927, vol. 14, p. 895.
- ² Ford, W. W., and Eliot, C. P.: *Journ. Exper. Med.*, 1928, vol. 48, p. 475.

TREATMENT OF ANAL FISSURE AND SPASM BY DIATHERMY.

BY

FRANK HOWITT, C.V.O., M.D., M.R.C.P.,
PHYSICIAN IN CHARGE, PHYSIO-THERAPEUTIC DEPARTMENT,
PRINCE OF WALES'S GENERAL HOSPITAL.

THE use of high-frequency currents in the treatment of haemorrhoids and fissures and other painful and ulcerative conditions of the anus was first advocated and practised by Professor Doumer of Lille in 1897. Doumer introduced into the anus a conical metal electrode attached to the Oudin resonator of the original high-frequency apparatus. A detailed report of eight cases treated by Doumer's method was published by Seeuwen¹ in 1906.

The advent of the modern diathermy apparatus rendered possible a greater and more sustained production of local heat, and Bensaude and Meyer,² by employing this current, were able to improve upon Doumer's original technique. They published a report of 114 cases so treated with markedly beneficial results. They made use of a modification of Doumer's electrode in their more chronic cases, but in the very acute ones they did not attempt to penetrate the anal canal even with a fine electrode, confining themselves initially to a saline pad pressed against the orifice of the canal. The indifferent electrode was a pad placed upon the lumbar region. Finally J. H. Marchand³ reviewed these various methods in the *Journal Médical Français* in 1929.

Our own experience of the treatment of certain ano-rectal cases by diathermy leaves no room for doubt that it is a curative agent of considerable value. Cumberbatch has drawn attention to the efficiency of this form of treatment, but it is remarkable that it has been so little employed in this country. The types of cases most suitable for its employment may be roughly subdivided under two headings—acute and chronic.

Acute Cases.

These include: (1) simple anal fissure and other ulcerations of the ano-rectal mucous membrane; (2) inflammatory and ulcerative conditions of the anus associated with haemorrhoids; (3) cases following operations on haemorrhoids and other operations in this area, in which the mucous membrane has not healed. Such cases are always attended with a considerable degree of spasm of the sphincter ani. The spasm cuts off the blood supply to the underlying mucous membrane, and the resultant tissue