MALIGNANT DISEASE OF THE LUNG*

BY

A. TUDOR EDWARDS, M.D., M.Ch., F.R.C.S.

SURGEON, BROMPTON HOSPITAL; SENIOR SURGEON TO OUT-PATIENTS,
WESTMINSTER HOSPITAL

(With Special Plate)

Malignant disease in the chest—and especially in the lung—has occupied the mind of the profession to a much greater degree during recent years than formerly. Whether there has been a definite increase in the incidence of malignant disease in the lung during the last ten or twelve years has been, and still is, much debated. Of one thing there appears to be no doubt: the incidence of cases diagnosed is much higher to-day than even as recently as five or six years ago.

Primary malignant neoplasms in the chest—and more especially those arising in the lungs—are extremely insidious in their onset, and may have been present for many months before their presence is suspected. Unfortunately in a large proportion of them dissemination into the glands of the mediastinum has resulted by the time the symptoms are sufficient to cause the patient to seek medical advice, and all hopes of successful surgical removal of the disease is gone. In the lung, carcinoma appears to arise primarily in two situations—the bronchi and the parenchyma. In the bronchi, carcinomatous changes may originate—

(1) in the epithelial lining of the tubes, comprising (a) the squamous-celled group which result from anaplasia or metaplasia of the normal epithelial lining cells, and (b) the so-called cylindrical-celled carcinoma in which there is very wide variation in structure; or

(2) in the mucous glands, in which growths—generally alveolar in structure—myxomatous degeneration is often seen.

In the lung parenchyma the histological appearances vary somewhat according to the type of tumour. In the diffuse pneumonic type, involving varying proportions from part of a lobe to the whole lung, the alveoli are occupied by masses of cuboidal, cylindrical, or flattened cells. The localized carcinomata in the parenchyma may be single or multiple, and consist of vesicles partially filled with papillary projections of cylindrical cells. Endothelioma of the lung has been described, the growth being localized to a greater or lesser degree, and consisting of masses of cells of endothelial type. These are sometimes confused with the squamous-celled carcinomas. Sarcoma is generally of the spindle-shaped type, and varies considerably in size, occasionally filling the hemithorax, eroding ribs and even chest wall. Sarcomata are much rarer than has been thought, and unquestionably a number of cases formerly described as sarcoma would now be classified as carcinoma.

SYMPTOMS AND SIGNS

The onset of malignant disease is generally very gradual and insidious, but in a number of cases is initiated by a pyrexial attack commonly labelled influenza. For weeks, and occasionally for months, the history in others may disclose nothing except increasing lassitude without localizing symptoms. This symptom is later accompanied by dyspnoea or non-productive cough, both persisting and tending to become worse. The general appearance is often good, and weight may even be gained at this period. Haemoptysis occasionally occurs early. Irregular pyrexia may be an early symptom, but the temperature is rarely high until the later stages, when there is retention of secretions, or when necrosis has occurred in the centre of the growth. A sudden attack of haemoptysis, associated with the expectoration of a quantity of purulent

sputum, occurs occasionally in malignant disease, and tends to confuse the condition with pulmonary abscess, especially as it is often preceded by a rise, and succeeded by a fall, of temperature. Pain is certainly rare in the early stages, but until the pleura or mediastinum is involved is never severe. Dysphagia occasionally results from the pressure of secondary deposits in mediastinal glands upon the oesophagus, and hoarseness from pressure upon, or involvement of, the left recurrent laryngeal nerve.

The signs vary according to the type of the disease. In the bronchial growths the physical signs are those of pulmonary atelectasis, bronchial breathing of a blowing type, some dullness, and often increase of voice sounds when the obstruction is incomplete, with dullness and weakness or absence of breath and voice sounds when complete bronchial obstruction is present. In the central or alveolar type physical signs may be absent, or may consist of localized dullness associated with weakness of breath and voice sounds. Extrapulmonary signs should also be sought for, such as enlarged supraclavicular glands and paralysis of the homolateral diaphragm due to phrenic nerve involvement. Evidence of venous obstruction, such as dilatation of the superficial veins of the chest and the veins of upper limb and neck, is comparatively rare in pulmonary neoplasms. Effusions of the pleura are usually late in their incidence, and are often haemorrhagic. Bronchiectasis secondary to the obstruction results in a proportion of cases, and its symptoms may mask those of the initial lesion.

DIAGNOSIS

The fact that the signs are atypical of any of the common lung diseases should immediately suggest the possibility of malignant disease of the lung. Diagnosis may be confirmed by further investigations. These comprise *x*-ray examination, before and after pneumothorax, and the introduction of lipiodol into the bronchi, bronchoscopy, thdracoscopy, and, finally, exploration of the thoracic cavity.

X-ray Examination

This examination should consist of observation of the patient under the screen to judge the range of diaphragmatic movement and the manner in which the lighting up of the lung field varies during inspiration and expiration. The different types of pulmonary growth give rise to different radiological appearances.

(a) *Growths Obstructing the Bronchus.*—A characteristic appearance is given in skiagrams where a neoplasm is occluding a bronchus sufficiently to prevent the passage of air into and from the area of lung supplied by that bronchus. The air in the occluded portion of lung is gradually absorbed, and the lung tissue appears opaque in the skiagram. Generally the area is somewhat wedge-shaped, with the apex at the occluded bronchus and its base at the periphery. When the main bronchus to a lobe is occluded the whole lobe becomes atelectatic and shows a shadow. When the obstruction has been present for some time the mediastinum may be displaced to the affected side. *X*-ray examination, following the introduction of lipiodol, will often show an irregularity in the shadow of the opaque oil which has collected above the obstruction, corresponding to the irregularity at the surface of the growth. Naturally, the oil will not enter the obstructed part of the lung. Other causes of bronchial occlusion, such as foreign bodies, can generally be eliminated by the history, but where there is any doubt bronchoscopy should be performed.

(b) *Central Growths.*—In localized growths in the parenchyma of the lung the skiagrams show a definite opaque shadow corresponding in outline with the tumour. There is complete absence of any pulmonary reaction

* A paper read at the joint meeting of the Sections of Surgery and Tuberculosis at the Annual Meeting of the British Medical Association, Winnipeg, 1930.

around, as is seen in infective conditions such as abscess, and this usually results in a very clear definition of the margins of the growth. Pleural and subpleural tumours often give similar appearances in skiagrams to those of central pulmonary neoplasms. Where pneumothorax is possible these can be differentiated, the lung collapsing partially over the intrapulmonary growths, whereas extrapulmonary growths can be seen outside the area of the collapsed lung. In a certain proportion of cases the centre of the growth necroses, and should infection spread in from the bronchi an infective abscess surrounded by new growth will result, whereon the x -ray appearances will be altered and the opacity be much more diffuse and ill-defined owing to the inflammatory reaction in the surrounding lung. This complication accounts for those cases of new growth which are diagnosed as pulmonary abscess and treated as such for some time before the underlying cause is suspected.

Bronchoscopy

In all cases in which there is evidence of bronchial obstruction, bronchoscopy should be undertaken. Not only these cases, but those of patients in whom haemoptysis is a prominent symptom and in whose sputum no specific organisms, such as tubercle bacilli, can be found after several examinations, should be submitted to bronchoscopy. During this examination it is often found difficult to distinguish a malignant growth from oedematous granulation tissue, and it is therefore essential that a small portion of suspicious tissue should be removed for histological examination.

Thoracoscopy

Following the production of an artificial pneumothorax or the replacement of an effusion, either haemorrhagic or serous, by air, thoracoscopy may be of value in a limited number of cases. In cases where the growth or growths are superficial in the lung it may enable a definite diagnosis to be made. It may also be of value in the differentiation of growths of the lung itself from growths of the pleura or chest wall.

Thoracic Exploration

It cannot be questioned that the enormous advance made in diagnosis and treatment of abdominal disease in the past is largely due to the frequency of exploration for obscure lesions. In our comparative ignorance of the early pathology of chest disease it is surely reasonable to expect that exploratory thoracotomy should be much more frequently undertaken than is the case if progress is to be made. Exploratory thoracotomy as carried out under modern conditions and, if possible, preceded by the induction of artificial pneumothorax, is certainly no more serious a procedure than abdominal exploration, and in general the patients have much less post-operative discomfort. Intercostal incision, with retraction of the adjoining ribs, gives sufficient exposure, and rib resection is unnecessary, except in older subjects in whom the costal cartilages are ossified. In each of two cases of suspected malignant disease submitted to exploration I have been able to remove successfully a teratoma, one containing six pounds of semi-solid sebaceous material.

TREATMENT

Apart from the various forms of x -ray therapy, no definite attempts have been made until recently to treat malignant disease of the lung in an adequate way. Owing to its insidious onset, and therefore the comparatively advanced stage which is reached before diagnosis is made, and owing to its position with regard to vital structures, the difficulty of complete removal or adequate irradiation of malignant disease of the lung is obvious. As cases have

been classified into two main groups—the parenchymatous and the bronchial—in x -ray diagnosis, similarly for treatment they may be grouped as: (1) central growths, (2) bronchial carcinoma.

Central Growths

Unfortunately this group, consisting of growths which offer most chance of surgical extirpation, forms a very small proportion of the total cases of malignant disease of the lung. Of 118 cases of primary malignant neoplasm in the lung which I have personally seen, only nine patients could be put in this category. These nine cases were submitted to exploratory thoracotomy, and six were inoperable because of the presence of secondary malignant glands in the mediastinum or widespread adhesion to the chest wall. The other three, comprising two endotheliomata and one squamous-celled carcinoma, were removed, and are worthy of record.

CASE I.—Endothelioma of Lung Involving the Right Lower Lobe and Diaphragm

The patient was a man, aged 60, and the condition necessitated removal of about three-quarters of the right lower lobe and a portion of the diaphragm. The patient died fourteen hours later from haemorrhage from an intercostal artery, which had been penetrated by the needle used in suture of the chest wall.

CASE II.—Endothelioma of Lower Lobe of Right Lung

The patient, a woman aged 48, was operated upon in December, 1928. Pain and discomfort were the only symptoms, apart from expectoration of a small quantity of blood-stained mucus on one occasion. The x rays showed a definite tumour involving the right lower lobe, and after pneumothorax was induced the lung and its contained tumour were seen to be free from pleural adhesions (Fig. 1). Exploration showed the tumour localized in the lung with no involvement of the mediastinal glands. A small cuff of normal lung was turned back at the hilum, and the lobar artery and vein to the right lower lobe ligatured. The bronchus was crushed and ligatured, and the cuff of pulmonary tissue was sewn over the stump after removal of the lobe. Airtight drainage was employed for forty-eight hours. Apart from a subacute dilatation of the stomach, convalescence was uninterrupted. This patient is alive and without symptoms at the present time, one year and eight months after operation (Fig. 2).

CASE III.—Squamous-celled Carcinoma of the Lung Involving the Chest Wall

In July, 1929, the patient, a man aged 55, complained of a slight cough and expectorated a small quantity of blood-stained sputum. X rays showed a mass in the left lung, extending to the chest wall. Another skiagram, fourteen days later, showed a similar appearance, but there was now evidence of spontaneous fracture of the sixth rib. There was an area of dullness and dry pleural cracking in the left axilla. Artificial pneumothorax showed adhesion of the mass, which had extended outward from the lung to the chest wall. In September, 1929, exploration was made, and the mass in the lung and the adherent chest wall was removed. This necessitated the excision of the major part of the left lower lobe and portions of three ribs and the intercostal structures between. Airtight drainage was instituted. The general condition was good after operation. The patient improved for a period, but later there was definite evidence of recurrence, from which death ensued about four months later. Histological section of the tumour showed squamous-celled carcinoma.

Of the remaining six cases in this group one patient had multiple primary growths in the right lower lobe with secondary glands in the mediastinum, and eventually died without further treatment. Three others were treated with radon seeds, and will be mentioned later, as will another who was treated by application of a radium plaque externally. One patient with an inoperable growth died from acute bronchitis involving the opposite lung.

Bronchial Carcinoma

This type will rarely offer any hope for successful removal owing to its position and to its early spread into the mediastinal glands. In one case only have I attempted to remove a growth in this region.

CASE IV.—Carcinoma of Bronchus of Left Upper Lobe

The patient, a woman aged 27, suffering from haemoptysis, had been treated for some months for pulmonary tuberculosis. X-ray examination showed a bilobed shadow in the subapical region of the left upper pulmonary lobe. Its sharp outline suggested a growth, and there was doubt as to whether it was benign or malignant. Pneumothorax, followed by α rays, showed the tumour in the lung tissue (Fig. 3). Exploratory thoracotomy on January 8th, 1930, showed the tumour—about the size of a tangerine—to be situated in the upper lobe. The overlying lung was incised, and the tumour was found to lie within an enormously distended bronchus. To this it was attached by a broad base. This base was excised with the tumour, whereon a sudden haemorrhage from the lobar artery ensued. Seizure of the bleeding area was followed by sudden collapse of the patient. A ligature was applied and the lung tissue rapidly sutured over the opened bronchus. Airtight drainage of the chest through a low intercostal stab wound was instituted, and the chest rapidly closed. A small transfusion of 200 c.cm. of blood was followed by rapid improvement of the patient, although the collapse was probably due to vagal shock. Steady convalescence resulted, and the patient is well and symptomless now, eight months after operation (Fig. 4). The growth proved to be a columnar-celled carcinoma.

Radium Treatment

When radical removal is impossible radium therapy should be considered for all growths of a malignant nature in the lung. Three methods are available: (1) teletherapy, application at a distance of large doses (1 to 4 grams) in a bomb; (2) external application of plaques; (3) insertion of radon seeds.

(1) *Teletherapy*.—This method is very attractive, for it does away with any necessity for operation and, being unaccompanied by pain, appeals to doctor and patient alike. My personal experience with this method is limited, and the possibilities of its variations entail a very extended trial before judgement can be passed. However, in those patients—six in number—which have been treated by this method alone, I have yet to see any, even temporary, improvement, and they have all died.

(2) *External Plaques*.—The application of radium on external plaques of Columbia paste or sorbo rubber having proved valuable in other regions, it has been used for five cases of malignant disease of the lung. It is quite obvious that the intensity of the rays in deep growths is so small that no effective action could be expected on growths in the main bronchi, and therefore it should be reserved for growths adjacent to the chest wall. By this method also I have yet to see improvement in the radiological appearance of the neoplasm, although in three cases where the growth had commenced to involve the chest wall and acute pain was a marked feature, the radium application has resulted in considerable amelioration of this symptom.

(3) *Radon Seeds*.—For growths in the main bronchi there are definite dangers of perforation into the mediastinum by haphazard insertion of seeds into the growth itself, and I have therefore devised a small hollow silver tube which carries eight radon seeds. This tube is inserted into the bronchus through the bronchoscope, and a small thread, attached thereto before insertion, passes up through the larynx, and is fixed to the cheek by strapping. The tube has been retained from five to eight days, and appears to cause little irritation. Withdrawal is easily effected by traction on the thread. This method I have used in four patients, but the records are too recent to decide as to its ultimate value. As these growths are

often of the squamous-celled variety there is a distinct possibility that we may get a curative result in cases where no extension to the mediastinal glands has occurred before irradiation.

For central neoplasms, and for those arising peripheral to the main bronchi, which are inoperable, the direct insertion of radon seeds would appear to be the method of choice. In every case in which the general condition of the patient does not contraindicate, the growth should be explored, its extent determined, and the seeds inserted as regularly as possible under direct vision. Where adhesions are extensive wide exploration is not possible, and, in these, careful study of skiagrams in various directions will aid in the even distribution of the seeds throughout the growth. Originally the seeds were introduced by one of the types of introducers used for growths elsewhere in the body. In the recent cases I have been employing an ingenious modification, which allows of coagulation of the needle-track by the diathermic current before the trocar is withdrawn. This simple but ingenious modification, which has been suggested by Dickson Wright, prevents the escape from the puncture holes of blood containing tumour cells, and the possible transplantation and development of these cells on the pleural surfaces.

The last three cases which I have treated by this method offer considerable encouragement. In one case the tumour shadow became much less dense and diminished in size, the persistent cough disappeared, and the patient, a man of 68, gained 11 lb. in weight. I have recently heard, eleven months after operation, that he has developed a hemiplegia, which is possibly due to a secondary deposit in the brain. A second patient, a woman of 35, was operated upon in February of last year. When she was examined six months later she had completely lost all sputum; pyrexia and cough were absent, and she had gained well over a stone in weight. X rays show almost complete disappearance of the opacity in the lung. The last patient, a man of 64, who also suffered from myocardial changes, is already much improved, although the seeds were only inserted early in June.

Results such as the above are, in my opinion, a distinct encouragement to persevere in the treatment of these cases by radium, and, even if cure cannot be obtained, the relief given and the prolongation of life afforded by these means thoroughly justify the comparatively small risk of the operation.

Deep X-Ray Therapy

This method of treatment has proved very disappointing for primary malignant lung tumours, and I have yet to see a patient who has benefited thereby. The newer and more powerful forms of apparatus may prove of more value, but unless other methods fail I am disinclined to recommend this treatment to my patients at the present time.

Secondary deposits in the lungs are most commonly multiple, and often bilateral. In such cases no question of operative treatment is likely to arise. In rare cases, however, a single secondary deposit may occur in the lung, and in such as the following may justify radical operation.

CASE V.—Secondary Myeloma in the Left Upper Lobe

The patient, a woman aged 57, gave the following history. In 1904 she had a myeloma curetted from the upper end of the right fibula. This operation was repeated in 1907. There was another recurrence in 1921, which was treated by radium. This resulted in considerable pain and swelling, and the right lower limb was amputated three weeks later. On July 22nd, 1927, there was a slight attack of haemoptysis. Inquiry elicited a history of a certain feeling of constriction in the upper chest for the previous six months. On examination there was a small patch of dullness over the left apex

posteriorly. X rays showed a well-defined shadow in the apex of the left upper lobe (Fig. 5). Pneumothorax proved the mass to be intrapulmonary, and, in view of the history, probably a secondary deposit (Fig. 6). The upper two-thirds of the left upper lobe were removed through an intercostal incision. Apart from some temporary collapse of the left lower lobe, due to a mucus plug in the bronchus, which was coughed up later, convalescence was uninterrupted (Fig. 7). This patient is alive and symptomless to-day, three years after operation.

Occasionally the rapidity of development of secondary deposits of limited extent in the lungs is diminished by deep x-ray therapy, and this treatment should be recommended only when its limitations are fully realized.

In the past, when a diagnosis of malignant disease was established, the tendency has been to abandon all hope. As diagnosis is becoming possible at an earlier stage it is surely time that some definite efforts should be made to deal with the problem of malignant disease of the lung in a more active way than by palliation of symptoms alone.

CATARACT OPERATION IN EXTREME OLD AGE

BY

R. H. ELLIOT, M.D., F.R.C.S.

LIEUT.-COLONEL I.M.S. (RET.)

It is commonly urged by relations that a patient is "too old for an operation for the removal of cataract." During the course of my experience in cataract operation, running into many thousands of extractions, I have on four occasions operated on patients 90 years old and over, and on one just before her ninetieth birthday. In addition to this I have operated on a very large number of patients between the ages of 80 and 90. No experienced operator will dispute that a cataract extraction is a shock to the patient, and that the lives of some old people are shortened by procedures of the kind injudiciously undertaken. The fact is worth noting that in all five cases the operation was successful, and also the further fact that in not one single instance does it appear to have shortened life. Before discussing the reason for this a few details of these cases may be of interest.

The first patient was a woman in Madras whose skin hung on her in satiny wrinkled folds, resembling the skin of an animal rather than that of a human being. She claimed to be "very, very old," but, like a number of the natives of India, could not give her exact age. She could say, however, that her eldest son fought in one of the Sikh wars. My assistants were able to calculate from this that, even allowing for the very tender age at which Indian women become mothers, she must have been at least 96 at the time of the operation, and possibly several years older. Not only did she make an uninterrupted and perfect recovery, but she signalized the event by stealing a necklace from under a neighbour's pillow the first night that her eye was released from the bandage. The result was that I had to interview her and the outraged lady in the orderly room next morning.

The next case was that of an old retired civil servant, who, being told that an operation for cataract was necessary, refused to have it done until after his ninetieth birthday, as he had undertaken to give a bride away on that date. He also made an uninterrupted recovery, and appeared to suffer no shock from the operation. A year later he came to see me, asking me to inspect my handiwork; he brought with him cuttings from an illustrated paper with his photograph, and the legend beneath it that he had made a record "drive" for his age. Considering the very small number of people who play golf at 91, the statement might readily be accepted. He was extremely indignant because his drive, as given in the

press, was, he maintained, ten yards short of the length he had really driven. (A true golfer!) I have lost sight of him for a good many years and conclude that he is dead.

The third case was that of an old general who had served many years in the Indian Army. I first operated just after his ninety-fifth birthday for double glaucoma. Six months later I extracted a cataract from one eye. The operations were successful; he stayed only the usual fortnight in the nursing home, and did not appear to be unduly depressed by the shock he had sustained. Whenever I went to see him he had a racy story to tell me. On the occasion of my visit to him in the home on the night before the operation he assured me that he felt quite young, and to satisfy me on this point he insisted on the nurse dancing one of the new steps with him round his room while he vigorously hummed a tune.

In the fourth case the patient was a man very distinguished in public life in his day, and accustomed to having his own way. He had both cataracts extracted towards the close of the ninth decade of his life. An after-ataract formed some time later in one of his eyes. His family were desirous that he should not undergo further operation, and, needless to say, I was very unwilling to undertake any procedure on him after 90 years of age, in view of the fact that the vision was quite reasonably good with the eye in question, while he could read diamond print and had 6/5 vision with the other. I tried to dissuade him, but he turned very determinedly on me and asked, "Do you think I shall see better with that eye if I have it operated on?" I replied, "Yes, but there is an element of risk to your life in this case which I do not feel justified in taking without putting the matter very clearly before you." His reply was, "Well, that is settled. I will come into the home." He was a perfect patient, as he always had been, and once again showed no sign of shock; his recovery was even and uninterrupted, and he regained the sight which he formerly had, equal to that of the other eye.

The fifth patient was operated upon a few months before her ninetieth birthday for a cataract of the Morgagnian variety. The operation was performed in two stages. The patient was stone deaf, and the only possible means of communicating with her was by the hands. We had to establish a code before the operation with signs for "look down," "look up," etc. She was a model patient, and in consideration of her great age she was allowed to sit up in bed the day after the operation. She remained in the home the usual fortnight, making a perfect recovery. Later she was fitted for glasses, though on account of her deafness it was a matter of great difficulty to conduct an accurate test. Her distance vision, under correction, was 6/12, and she read No. 2 (pearl) print easily. She is now over 92, and I recently heard that she was doing well.

Colonel Kirkpatrick has very kindly allowed me to include in this record a case of his own, in an old "John Company" pensioner, 96 years of age, upon whom he operated in Madras. He cannot remember with absolute certainty whether one or both cataracts were extracted, but he is practically sure that the old man came back three or four months later for the second eye to be treated. The point of great interest is that the patient had rhinophyma, causing a marked disfigurement of his nose, and that after the recovery of his sight he insisted on having this operated on at the General Hospital, Madras. He went home with a greatly improved appearance, having regained the strikingly handsome looks of his earlier years—not a bad sequel at about 97.

COMMENTARY

It is probable that very few surgeons have had the opportunity of operating on so many patients at such very advanced ages, and this seems to justify a short note on the cases. What appears to me of much more interest is: (1) that in every one of these cases the operation was successful; and (2) that in none of them was the shock of the operation severe. Remembering how many patients aged 70 or less are in too frail a condition to be submitted

A. TUDOR EDWARDS: MALIGNANT DISEASE OF THE LUNG.

FIG. 1.—Skiagram showing tumour in right lower lobe. Note absence of inflammatory reaction in adjacent pulmonary tissue. (Case 2.)

FIG. 2.—Appearance of chest five weeks after complete removal of right lower lobe of lung. (Case 2.)

A. TUDOR EDWARDS: MALIGNANT DISEASE OF THE LUNG.

FIG. 3.—Skiagram showing tumour in upper left lobe with shadow above, due to area of collapsed lung. (Case 4.)

FIG. 4.—Skiagram taken two months after removal of carcinoma of bronchus. (Case 4.)

A. TUDOR EDWARDS: MALIGNANT DISEASE OF THE LUNG.

FIG. 5.—Shows well-defined tumour at apex of left lung. (Postero-anterior view.) (Case 5.)

FIG. 6.—Appearance in erect posture after induction of pneumothorax. The weight of the contained tumour has displaced the apex of the lung downwards. (Antero-posterior view.) (Case 5.)

A. TUDOR EDWARDS: MALIGNANT DISEASE OF THE LUNG.

FIG. 7.—Appearance four months after partial pneumectomy. Note the displacement of the mediastinal structure to the left to compensate for loss of pulmonary tissue. (Case 5.)

JAMES F. BRAILSFORD: HYDATID DISEASE IN ENGLAND.

FIG. 1.—Photograph of a pig's liver showing innumerable cysts. They were closely approximated throughout the whole organ giving it a honeycomb appearance on cross section.