to that described, save that the specific agglutinating serums are diluted to eliminate coagglutinins and that smears of the mixed pneumococcal exudate and serum are made on slides, dried and stained. Sabin's method therefore sacrifices the opportunity of observing the striking and entirely characteristic change in the appearance of the cocci which follows conjugation with the specific antibody-bearing protein; and, further, his method introduces additional manipulations in the preparation, drying, and staining of the films. Neufeld, twenty years ago, applied the microscopic method to typing pneumococci, but was dissatisfied with the results he obtained with Type III. In my experience, however, no difficulty has arisen so far with this organism or with material from cases of Type III pneumonia. In any event the results with Types I and II are decisive, and are the key to early treatment.

The experiments described here have been carried out with the assistance of the Medical Research Council.

REFERENCES

- ¹ Vide Glasgow and Edinburgh reports, *Lancet*, December 27th, 1930.
² Sabin: *Journ. Infect. Dis.*, 1930, xlv, p. 469.

THE ENIGMA OF THE CHILBLAIN

BY .

RUPERT HALLAM, M.D.

PHYSICIAN, SKIN DEPARTMENT, ROYAL INFIRMARY, SHEFFIELD

The discomforts and disabilities caused by chilblains during the winter are very real, although non-sufferers may regard the complaint as trifling. We have but a fragmentary knowledge of the subject; it is known that cold is the exciting cause, but how one individual escapes and another suffers has never been determined. Consequently it is improbable that efficient means of prevention will be found until more is known of the etiology and pathology of the condition.

The Etiology of Chilblains

Perniosis is essentially an affliction of youth, as the following analysis of 100 consecutive cases shows:

Age	Number of Cases
0-10	54
10-20	29
20-30	4
30-40	5
40-50	4
50-60	3
60-70	1

From the above it is evident that after a period of years an immunity is acquired. Certain persons are free in their early years, but suffer in middle age; strange to say, however, chilblains are seldom met with in the aged, even when serious organic disease is present.

A histological examination in the early stages shows that there is a rapid degeneration of the small vessels, accompanied by perivascular infiltration. There is not a mere transudation of serum, as some writers would have us believe, but a sudden and severe damage affecting the vessels of the papillary layer of the cutis. Attempts to produce a chilblain experimentally by the application of solid carbon dioxide, injection of histamine, etc., have failed. In a susceptible individual a chilblain commences to make its appearance about eighteen hours after exposure to cold of the requisite severity. Should the temperature remain low little alteration in the state of

the chilblain ensues, but a change of wind bringing a warmer atmosphere produces a rapid amelioration of the symptoms; and the chilblain disappears with surprising rapidity. Change of environment may also produce a striking alteration in individual susceptibility. These facts are of importance, for it is not unlikely that they explain the reputation of many of the so-called cures for chilblains.

Chilblains Associated with Other Diseases

In order to investigate some of these suggestions, I endeavoured to ascertain the relative frequency of chilblains in certain diseases of known etiology, and also to determine the incidence in various social states. A more detailed account of these inquiries has been given in a previous communication.¹ The incidence of chilblains among patients attending the casualty department of the Sheffield Royal Infirmary compared favourably with that of others in more affluent circumstances. On the other hand, statistics obtained from the medical out-patient department and from the tuberculosis dispensary in the same city indicate that the incidence rises with a disturbance of the general health, as shown below.

	Percentage Affected with Chilblains
1,000 casualty patients	5.3
1,000 medical patients	9.2
1,000 tuberculosis patients	13.2

The figures in the last group are of particular interest; for some writers, among them Darier,² allege that the chilblain is actually a tuberculide. Although the above figures give some support to this theory, it is probably more accurate to assume that tuberculosis and some other diseases which impair the general health are predisposing causes. An analysis of 1,275 patients attending the medical out-patient department affords some interesting comparisons. Taking first the abnormalities of the circulatory system, 14 of my series had various degrees of arterio-sclerosis, yet in no instance was there a history of concomitant chilblains. On the other hand, of 86 suffering from some form of heart disease there was a record of recent chilblains in 15—that is, 17.4 per cent. From these figures it appears that a gross defect in the arterial wall may exist without materially diminishing the vitality of the capillary. The frequency with which chilblains are met with in disorders of the heart may be accounted for by the lessened propulsion force, and consequent slowing of the blood stream, interfering in some way with the nutrition of the superficial vessels. An examination of patients suffering from some definite imperfection of an endocrine gland is instructive. Among the 1,275 patients referred to 12 were suffering from myxoedema, but none suffered from chilblains; 24 were under treatment for exophthalmic goitre, of whom 3 had had chilblains since the onset of hyperthyroidism; this appears to show conclusively that chilblains are not necessarily associated with excess of thyroid secretion.

"Chilblain Circulation"

Most observers will agree that many chilblain subjects exhibit some peculiarity of the peripheral circulation in winter time. This was confirmed by an investigation of 100 cases of chilblains, for of this number 51 showed an apparently abnormal circulation in the hands and feet. It must be admitted that the records are of little intrinsic value, for it is impossible to define what is meant by normal blood supply. Under varying conditions the subjects might show different responses to heat and cold; furthermore, it is incorrect to assume that the peripheral circulation is necessarily normal because it appears to be

so clinically. Of the 51 cases showing clinical manifestations of variation of the circulation from the normal, the most obvious signs were cold hands and feet, accompanied by cyanosis of the hands and the lower third of the legs—a state of the peripheral circulation which is not infrequently referred to as the “chilblain circulation.” What is the pathology of the abnormal response to cold? A histological examination of the skin from such a cold blue hand, even where there are no chilblains, shows some thickening of the walls of the smaller vessels. It is agreed that cyanosis and coldness indicate a slowing of the blood stream. Krogh³ states that the temperature of the skin is primarily determined by the rate of blood flow, which in turn depends on the state of contraction of the arteries and arterioles. As the shade of the skin is controlled by the state of the subpapillary venous plexus and capillaries, it is obvious that these small vessels must be dilated in perceptible cyanosis. This is confirmed by an examination with the capillary microscope, a method of observation which is unfortunately very limited, for, excepting in the nail folds and lip, it is not possible to see more than a small section of a capillary loop. On examining a pronounced case of this type of cyanosis, the investigator almost instinctively presses the dorsum of the cold blue hand, which fills in slowly from the periphery like a closing diaphragm. In the individual with hands of normal colour, however, the white patch reddens from below as well as from the periphery. The only reasonable explanation of this interesting phenomenon is that first there is a closure of the arterioles, and secondly a return flow of blood from the neighbouring superficial venules. Spalteholz pointed out that there are no valves in these venules, consequently there is no obstruction to the return along them; but the blood cannot reach the venules from the veins in the deeper layers owing to the presence of valves. Again, if the ulnar half of the hand is immersed in hot water it gradually becomes red, whereas the non-immersed portion retains the original cyanotic colour. The line of demarcation is very sharply defined, indicating that the large arteries are not involved. It appears, therefore, that the cyanosis and colour is accounted for by a constriction of the arterioles, together with a dilatation of the capillaries and subpapillary venous plexus. Although this type of circulation is frequently encountered in patients who are subject to chilblains, it is certain that this peculiar pathological change alone will not produce a chilblain, even with an accompanying exposure to cold, for in the syndrome known as acrocyanosis an extreme degree of cyanosis and coldness of the extremities may exist without the slightest evidence of a chilblain. At the same time, it must be admitted that a high percentage of acrocyanotics do suffer from chilblains. Again, in Raynaud's disease, which Lewis⁴ has shown to be a spasmodic arrest of the circulation due to a contraction of the digital arteries, chilblains are not commonly met with. The etiology of the chilblain is therefore still obscure. It appears that there must be an unknown and independent factor producing a change in the wall of the smaller cuticular vessels, in addition to the factor which causes slowing of the blood stream. This change in the vessels is necessary before the exposure to cold is able to damage the tissues, for in normal skin, as Lewis⁵ has pointed out, even after the skin has been sufficiently frozen to form a wheal, the capillaries are afterwards found to be intact.

Biochemical Investigation

The anticipation that biochemical investigation would advance our knowledge on the subject has, unfortunately, not been realized. In view of the fact that calcium salts inhibit the transudation of serum through the capillary

walls, and as a result also of belief in the existence of a diminished coagulability of the blood in chilblain subjects, calcium therapy has been singularly popular for many years. It is significant, however, that Percival and Stewart⁶ find the serum calcium normal as regards both quantity and quality; moreover, these workers also ascertained that the ingestion of calcium salts in quantities usually prescribed does not appreciably raise the amount of serum calcium.

The Blood in Perniosis

My investigations of the blood in both chilblain subjects and acrocyanotics does not support the contention that the coagulation time is delayed. The method of estimation adopted was that of Dale and Laidlaw. The first drop of blood was always selected, and the test was repeated three times. The tubes were immersed in water at 37° C. The following table shows the average time.

Blood Coagulation Time

Ten chilblain patients: 2 min. 16 sec., 2 min. 36 sec., 2 min. 10 sec., 2 min. 26 sec., 2 min. 6 sec., 2 min. 29 sec., 2 min. 9 sec., 2 min. 7 sec., 2 min. 10 sec., 2 min. 3 sec.

Ten acrocyanotic patients: 1 min. 58 sec., 2 min. 18 sec., 2 min. 15 sec., 2 min. 2 sec., 2 min. 52 sec., 2 min. 8 sec., 2 min. 27 sec., 2 min. 38 sec., 2 min. 31 sec., 2 min. 5 sec.

Six normal controls: 2 min. 18 sec., 2 min. 10 sec., 1 min. 55 sec., 2 min. 19 sec., 2 min. 8 sec., 2 min. 11 sec.

Taking into consideration the work of Percival and Stewart, as well as the above, it is doubtful whether there is an adequate reason for the belief in a calcium deficiency.

In consequence of the recently acquired knowledge relating to the influence of vitamin D on the development of bone and teeth in youth, and also of the important function of the accessory food factors in the prevention of disease, it is important to know whether changes in the peripheral vessels may be attributed to a deficiency of one or more constituents of the diet. It has been found, however, that in some groups of school children having an average diet, the incidence of chilblains compares favourably with others having an added amount of vitamins A and D, and it can be stated definitely that some apparently healthy children who have had a generous diet over a long period, containing more than a sufficiency of calcium, with additional amounts of vitamins A and D, may acquire chilblains on exposure to cold. It has also been noted that the incidence of chilblains in a group of London elementary school children (100) exposed to a carbon arc light during the winter was no less than in control groups.

CONCLUSIONS

In conclusion, I submit that the pathology of the chilblain is complex and still imperfectly understood. The actual lesion is not due to a mere transudation through the capillary wall, for the superficial vessels of the susceptible individual are the subject of a pathological change, which enables sudden lowering of the temperature to act with great intensity, destroying the endothelium of the peripheral vessels *en masse*. Among the predisposing causes are hereditary weakness, lowered resistance due to tuberculosis, and other toxæmias.

REFERENCES

- ¹ Hallam, R.: Etiology and Pathology of Chilblains, *Transactions of the Eighth International Congress of Dermatology*, Copenhagen, 1930.
- ² Darier: *Précis de Dermatologie*, 1923.
- ³ Krogh: *The Anatomy and Physiology of the Capillaries*, 1922.
- ⁴ Lewis: *Heart*, xv, No. 1.
- ⁵ Idem: *The Blood Vessels of the Human Skin and their Responses*.
- ⁶ Percival and Stewart: *Brit. Journ. Derm. and Syph.*, 1927.