TREATMENT OF CARBOLIC ACID POISONING

BY

O. S. GIBBS, M.B., CH.B.

FORMERLY PROFESSOR OF PHARMACOLOGY, DALHOUSIE UNIVERSITY,
CANADA; VISITING LECTURER IN PHARMACOLOGY, JOHNS
HOPKINS MEDICAL SCHOOL, U.S.A.

The carbolic acid group constitutes one of the most popular suicidal poisons, and thus possesses a clinical interest to any medical man, especially in regard to immediate and proper treatment. Reference to many of the works on toxicology is often disappointing, since the essential features of treatment are apt to be obscured by a discussion of impracticable, or even useless, methods. Situated in a laboratory one rarely comes in contact with such cases except by consultation, and thus only at a late stage. Two recent cases of this type, however, raised the question of the most practical procedure that could be adopted by the physician in charge.

There is no need to repeat the well-known details of this form of poisoning. Briefly stated, carbolic acid is a quite non-specific poison, causing loss of function and eventual cell death to any tissues with which it may be brought in contact, if it is sufficiently concentrated or acts over a sufficiently long time. Three places are particularly subject to damage. The first is the site of absorption, be it skin, gastro-intestinal tract, or any other area. If the acid is concentrated, coagulation of the tissues will ensue here, leading to mass death of the cells. After absorption the central nervous system is particularly susceptible, as with any fat-soluble depressant, leading to the typical descending type of functional inactivity, with final failure of respiration. Since the kidneys deal with the poison in a relatively more concentrated form, and over a longer period, in their work of elimination, they are therefore very liable to damage, and this constitutes a danger which is very serious. That carbolic acid is a general tissue poison must not be lost sight of, however, for in any case of poisoning, especially if functional weakness or damage previously existed, other organs may be predominantly affected. The liver and heart are of particular interest.

As with most poisons, the problem of treatment divides itself into three parts. The first and most important aim is to limit absorption; secondly, to help the body to withstand the effects of the poison; and thirdly, to assist in its elimination. It is in regard to the first and most urgent problem that I venture to suggest a form of procedure that has the merit of simplicity, and is probably as effective, while it is certainly as safe, as any other recommended method.

The treatment is based on the fact that carbolic acid is much more soluble in oil than in water, and thus readily leaves a watery medium for one of oil, in which it remains pharmacologically inert. In this connexion I may recall the established fact that carbolized catheter oil has no virtue as an antiseptic. It would appear, therefore, to be sound treatment to apply oil freely to the site of absorption; thus, in the case of carbolic acid burns, washing in oil followed by soaked oily dressings is a logical proceeding.

When the acid has been swallowed an immediate administration of a large quantity of oil to the stomach contents would certainly render unabsorbable much of the acid. The most suitable substance for this purpose is ordinary medicinal liquid paraffin, which takes up the acid efficiently, and is entirely non-absorbable even by damaged mucous membrane; nor is it rendered absorbable by digestive processes; in this it possesses a great advantage over vegetable or animal oils, which are occa-

sionally recommended as an adjunctive treatment. The use of this oil in any corrosive poisoning is sound, because not only will it allay irritation by mechanical means, but, as an unabsorbable oil, it readily assists its own elimination along with anything that may be dissolved therein. The effective dose would have to be large—at least half a pint (200 c.cm.), and as much more as one could induce the patient to take. Administration of the oil would not invalidate the use of the wash-out tube, though it would undoubtedly render this somewhat questionable proceeding even more messy. In cases where lavage is carried out, oil might with advantage be readministered prior to withdrawing the tube.

Too much stress cannot be laid on the fact that carbolic acid very rapidly penetrates tissues, especially unprotected tissues, and that even a short delay may be enough to affect the treatment adversely; thus the administration of paraffin has a great practical advantage over the wash-out as a primary method, since the time factor is much lessened. Treatment would then be continued by the use of repeated doses of magnesium sulphate according to the modern method, which is founded on the sound basis of retarding absorption and facilitating elimination from the bowel.

It is fairly well known that if pure carbolic acid is immediately wiped off the skin with pure alcohol a burn may be escaped. The reason for this is that the skin grease may temporarily hold up penetration of the acid, and the alcohol removes both. I had the interesting experience of seeing this idea extended to the use of alcohol dressings, the result of which was that the patient had become completely comatose and remained so for twelve hours. This was not surprising, since it would be difficult to devise a method more favourable to absorption. Alcohol is one of the few substances directly and readily absorbed from the stomach; furthermore, it may actually cause the absorption of drugs not otherwise taken up by this viscus. The use, then, of dilute alcohol as a wash-out medium, recommended in certain books, seems to be based on an imperfect understanding of the facts, and also appears a needlessly dangerous procedure.

In regard to the remaining treatment I have little to add to existing knowledge, though greater emphasis might profitably be placed on the conception that the patient has all his functions seriously impaired, and thus should be treated with the same care that severe surgical shock or anaesthetic poisoning demands. Prolonged journeys, inadequate provision for warmth, or any other cause of functional fatigue, should be rigorously avoided. Large amounts of fluid are indicated in order to facilitate the work of the kidneys, and thus should be administered by other routes than the gastro-intestinal for at least forty-eight hours when the poison has been swallowed, since absorption is not desired in this area. Alkalis seem definitely beneficial, but at least $3\frac{1}{4}$ oz. (20 grams) should be administered with the fluid during the twenty-four-hour period. High fluid intake should be continued until the urine is carbol-negative.

Dangerous symptoms can only be treated according to general principles. Caffeine is the best respiratory stimulant. In view of the fact that I have read in this *Journal* two discussions on morphine and heroin poisoning in which no mention of caffeine was made, it may be pointed out that pharmacological evidence shows that caffeine is the best respiratory stimulant in these conditions, as well as probably in any other. One very good reason for its use is that caffeine has no dangerous secondary effects in large therapeutic doses, providing the simple precaution of slow injection is adopted. The same cannot be said for either strychnine or lobeline. In animals poisoned with morphine I have found lobeline neither so satisfactory nor so reliable as caffeine.

Strychnine is a valuable drug, since it combines a powerful action on the vasomotor system with the property of stimulating the liberation of adrenaline. This latter substance has a direct stimulant action, not only on the vessels, but on the heart. Thus, while strychnine has no direct cardiac action of value, it may act as a stimulant to the heart through this mechanism; which is an interesting sequel to an old fight between clinicians and pharmacologists, in that both are now shown to be justified.

For continued cardiac action one of the digitalis bodies is indicated, strophanthin being probably the most convenient where the oral route is contraindicated. Strophanthin and caffeine are best administered intravenously in such cases, or for urgent need.

Summary

1. The use of liquid paraffin is advocated for carbolic acid burns.
2. The immediate administration of a large dose of liquid paraffin is advocated as the primary treatment of oral carbolic acid poisoning.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

TWELVE-YEAR-OLD ECTOPIC PREGNANCY

The following case of a woman who carried a dead full-time foetus for twelve years in her abdominal cavity, and had three normal deliveries of healthy children before its removal, seems of sufficient interest to publish.

A Chinese woman, aged 34, was admitted to the Wuhu General Hospital (after walking ten miles) complaining of abdominal pain and difficulty in micturition. There was no vomiting, but she felt feverish and was unable to sleep. Her temperature was 101° F.; pulse 100. Bowels opened regularly every day. These symptoms had existed for fifteen days; previously she had been quite healthy.

The patient had had no past illnesses. She had been married for fourteen years and had three children—all normal deliveries—alive and well, aged 9 years, 5 years, and 9 months respectively. No miscarriages had occurred. She had been unable to feed any of the babies as she had no milk. Menstruation, of three days' duration, was regular at thirty-day intervals.

On examination of the abdomen a mass presented, the size, shape, and position of a seven months' gravid uterus. She showed no concern about this, stating that it had been there for twelve years. Evidently it had grown to its present size in a few months, and had never altered since or given any symptoms. During pregnancy it moved upward and to the right, returning to the mid-line after delivery. On palpation the mass was clearly defined, except below, where it was continuous with the pelvic organs. It was hard, immovable, and the skin over it freely movable. It was tender to palpation all over; tenderness also extended into the epigastrium and right hypochondrium. Bimanual vaginal examination showed a low cervix, while continuous with the upper pole of the uterus, and apparently attached to the right appendages, was the hard, tender, immovable tumour. A provisional diagnosis was made of (a) fibromyoma, (b) right ovarian cyst. Urine was passed normally with straining; catheterization produced a few ounces of residual urine.

Abdominal pain gradually grew worse, and a week after admission it was decided to operate. Under general anaesthesia the abdomen was opened by a mid-line incision, and on opening the peritoneum a dark brown mass was seen to occupy the greater part of the abdominal cavity, while surrounding it on all sides were adherent coils of slightly inflamed intestines and omentum. The mass was extremely hard in parts, while in others the finger sank into decomposing tissue with a flow of thick brown fluid. Fears were entertained that

it might be a malignant growth, and the advisability of continuing the operation was considered. However, the separation of adherent coils was started in the easiest part, and on isolating the smooth curve of a foetal spinal cord the nature of the tumour was realized. The foetus was lying in the right occipito-posterior position in an immensely dilated right tube. After dividing adhesions for an hour and a half it was possible to apply a clamp along the right side of the uterus and remove the foetus, with the tube and appendages. The placenta was adherent to the right tube and anterior abdominal wall, and was scooped out with the hand.

The wound was closed in layers, with a drainage tube in the pelvis, and the patient made an uninterrupted recovery. She was discharged home quite well in a month's time.

After operation the following story was obtained from the patient. Twelve years ago, when the lump grew, she thought she was pregnant, and when the time came for delivery pains started, but nothing happened. After three days she was told by a Chinese medicine man that she had not been pregnant, whereupon he needled the lump, and told her she would have it for the rest of her life. The story was verified by her husband.

An x-ray of the removed tumour is reproduced.

X-ray of tumour removed from the abdomen of a Chinese woman, where it had remained for twelve years.

Ectopic pregnancy going to full term is rare, and the omission of the condition in the differential diagnosis of this case is perhaps excusable.

I am indebted to the American Free Church Mission authorities at Wuhu for inviting me to operate at their hospital during three weeks' residence in the port, and to Dr. Watters, the medical superintendent, who gave valuable assistance during the operation in this case.

W. WARD DARLEY, M.R.C.S., L.R.C.P.

H.M.S. Gannet, China Station.

PUERPERAL INVERSION OF UTERUS

Dr. Noel Hypher records a case of acute puerperal inversion, in the *Journal* of August 2nd, 1930 (p. 179). I had a similar case this year, the details of which I now record.

The patient, a fisherman's wife, aged 29, was confined on January 4th with her first baby. As the labour was