angles of tilting will be associated with the corresponding raising of the heel—namely, 15 degrees tilting with $1\frac{1}{2}$ -inch heel ; 30 degrees with $3\frac{1}{2}$ -inch heel ; and 45 degrees tilting with 5-inch heel. But even if the heel of the foot is supported by a raised heel of the boot, the pose of the foot is not very satisfactory, because the calf muscles are thrown out of action to an extent measured by the degree of tilting of the foot, and the patient loses spring and resiliency of step. Probably 15 degrees of tilting is the maximum advisable, representing a gain in height of about an inch, since the other shoe must have a heel of at least three-quarters of an inch. Both instrument makers and surgeons, however, have gone much beyond this limit, and have tried to convert the foot into an extreme equinus position. A special type of boot is widely advertised in which the heel is 5 or 6 inches off the ground, the equinus position being masked by making this elevation of the heel inside the boot. Such an appliance does disguise the deformity, but it does so at the expense of the muscular balance of the leg. In the same way the Mikulicz operation, by which the os calcis and astragalus are removed and the rest of the tarsus and metatarsus brought into a straight line with the tibia, results in a gain of about 5 inches in the length of the leg, but the limb, in consequence, is much less useful for walking than an artificial limb would be. There is an interesting specimen in the Museum of the Royal College of Surgeons representing the leg and foot of a girl, aged 18, on which the Mikulicz operation had been performed by Sir William MacCormac. The patient had infantile paralysis with paralytic equinus, and the leg was amputated by Sir John Bland-Sutton owing to the pain and uselessness of the limb, in spite of the fact that perfect bone union had occurred in good line. It may therefore be laid down as a definite rule in dealing with a weak, short leg, that a plantigrade position of the foot should be preserved, or certainly not departed from by more than 15 degrees of tilting.

SHORTENING OF THE SOUND LEG

For equalizing the length of legs another valuable device, which I think is not resorted to as often as it might be, is the shortening of the longer leg. In the majority of cases we are considering, whether tuberculous hip disease, infantile paralysis, or congenital dislocation of the hip, the long leg is strong and has good sound bones with perfect muscular and vascular supply. In such a leg the femur may be shortened with perfect impunity by 25 per cent. of its total length—that is, in the adult about 4 inches. I have done this operation in various ways, by taking a piece right out and uniting the shaft over a peg, or by a "step-cut" operation, but I now use an impaction method, which is much simpler than either of these (Fig. 1, C and D). If, for example, 4 inches is to be taken from the femur, the bone is exposed by a lateral longitudinal incision in its lower part about 6 inches long. The muscles are divided between the rectus and the vastus externus, and the bone is sawn through about 2 inches above the condyles. The shaft is brought out of the wound and 2 inches are cut off. The remainder is removed by a longitudinal saw-cut 2 inches in length, passing from side to side, the anterior part being cut away, leaving the posterior part of the femur, including the linea aspera, sticking out. This is trimmed by the saw and chisel into a peg about half an inch thick. The lower fragment of the femur is bored by a twist drill of suitable diameter, and the upper fragment is then impacted into it, and fixed in place by a single transverse ivory nail. It is very remarkable how the slack of 4 inches of muscle is taken up within a few months and the full muscular power of the leg regained.

Two examples will suffice to illustrate the possibilities of this method.

CASE I

A woman, aged 25, had a long-standing infantile paralysis with dislocation of the hip on the same side. When she came to me the left leg was $5\frac{1}{2}$ inches off the ground, caused partly by tilting of the pelvis and partly by insufficient growth of the leg, which was encumbered by a heavy apparatus consisting of a calliper splint and high boot. I did three operations. First, the left flail foot was stabilized by a partial tarsectomy (Naughton Dunn's operation), so that she could get firm plantigrade stance without irons ; secondly, an osteotomy was performed below the left trochanter, so as to bring down the pelvis on the left side, lengthening the leg by $1\frac{1}{2}$ inches and allowing her to support her body weight direct on the femur, instead of hanging it on to the capsule of the dislocated joint ; thirdly, six months later, the right femur was shortened by 4 inches, thus bringing her two legs to the same length. She can now walk on limbs of the same length, and for the first time in her life she can wear an ordinary pair of shoes.

CASE II

A man, aged 30, who must have had osteomyelitis of the upper end of the right femur in childhood, came to me wearing the largest "surgical" boot I have ever seen. It was 9 inches high at the heel and weighed 7 lb. He was sent to me for amputation, and it was obvious that an artificial limb would have been much more serviceable to him than the horrid boot. He had, however, an adduction and flexion deformity of the hip-joint, and this would have had to be corrected even if an amputation had been done. He gladly consented to a series of reconstructive operations, by which his own foot and leg have been preserved. An osteotomy was performed below the neck of the left femur, and the leg put up in abduction. This gained 2 inches for the functional use of the leg. The right leg was shortened by 4 inches, and his two legs are now within 3 inches of the same length. He was so pleased with this result that he is now trying a boot raised 3 inches at the heel and $1\frac{1}{2}$ inches in the sole, postponing for the present a possible bone-lengthening operation on the left femur.

As the result of the active campaign that is being carried out to-day for the treatment of crippled children there are fewer of these cases of adult gross deformity than formerly ; but there will always be a sufficient number to make the problems involved of great practical importance.

A CASE OF PERNICIOUS ANAEMIA RESISTANT TO TREATMENT

BY

A. M. KENNEDY, M.D., M.R.C.P.

PROFESSOR OF MEDICINE, UNIVERSITY OF WALES, AND DIRECTOR OF THE MEDICAL UNIT, WELSH NATIONAL SCHOOL OF MEDICINE, CARDIFF

A civil servant, who had always lived a temperate life in this country, at the age of 52 noticed that he was becoming pale and rather easily tired, and a year later, on account of these symptoms having become worse, he spent a month at a convalescent home on the South Coast. Shortness of breath and palpitation on exertion began to trouble him ; sleep did not come easily at night ; and six months after returning home his doctor told him that he was suffering from pernicious anaemia, and prescribed from a quarter to half a pound of liver daily. He did not seem to make any progress, however, either for better or for worse ; and after six months' treatment with liver daily he was admitted to hospital under my care.

He was then 54 years of age, a well-nourished man, 11 st. 10 lb. in weight, with fairly considerable subcutaneous fat. His skin and mucous membranes were pale, the former possessing the slight lemon tint so typical of pernicious anaemia. He had been treated for an appendix abscess at the age of 48, but had had no other illness before the onset of the complaint from which he was now suffering. His appetite had always been good, and he had not suffered from vomiting, diarrhoea, or from troublesome constipation.

A haemic murmur was audible at the base of the heart, but nothing further of note could be made out on examination of his chest. The liver was not enlarged, and the spleen was not palpable below the costal margin, nor was the splenic dullness found to be increased on percussion. He complained of slight tenderness on deep palpation in the left iliac fossa, where, through a resisting abdominal wall, a small though rather indefinite sausage-shaped mass seemed just palpable along the course of the colon. On inquiry he stated that for some years he had passed at varying intervals a little mucus on defaecation, especially if he had allowed his bowels to become constipated. Examination per rectum revealed a few small haemorrhoids only. His teeth were all artificial, his tongue was moist and clean, smooth, but free from any soreness, and he did not admit to having suffered from a sore tongue previously. No enlarged glands were present anywhere, his arteries were not thickened, and the Wassermann reaction of his blood was negative. Blood examination on admission to hospital (April 25th, 1930) revealed the typical picture of pernicious anaemia, as follows:

Red cells, 1,030,000 per c.mm.; white cells, 2,000 per c.mm.; haemoglobin, 28 per cent.; colour index, 1.4; platelets, 61,000 per c.mm. Red cells well stained and filled with haemoglobin, showing marked variation in size and shape; numerous megalocytes and microcytes present, and an occasional megaloblast; reticulocytes found only after prolonged search. Polymorphonuclears, 45 per cent.; eosinophils, 1 per cent.; lymphocytes, 46 per cent.; large mononuclears, 8 per cent.

The patient was then put to bed and given, in addition to the ordinary hospital diet, half a pound of lightly cooked liver and one orange daily for the next two months, and his blood was examined regularly at weekly intervals. His general condition remained much about the same, and any improvement that occurred in his blood was practically negligible. On one occasion only the reticulocytes rose to 2 per cent.; on all other occasions they were much fewer, and found only after prolonged search. At the end of two months' liver treatment examination of the blood showed (June 20th):

Red cells, 2,070,000 per c.mm.; white cells, 4,200 per c.mm.; haemoglobin, 38 per cent.; colour index, 0.91; platelets, 42,000 per c.mm. Red cells well stained, with some variation in size and shape, some larger and others smaller than normal, but no nucleated reds obvious. Reticulocytes found only after prolonged search.

In view of the fact that the patient had had from a quarter to half a pound of liver daily for the six months before coming into hospital, and that, after admission to hospital, two months' treatment with half a pound of liver daily produced an increase in the red cell count of only one million per c.mm. (the highest count being no more than 2,070,000 per c.mm.), with less effect on the other constituents of the blood and with practically no effect on the reticulocytes, it was concluded that the case was resistant to liver therapy. (The changes in the blood occurring at weekly intervals during the two months when the patient was undergoing liver treatment in hospital are shown in the first part of Chart 1.) The liver was thereupon stopped, and the patient given desiccated extract of stomach tissue in the form of ventriculin for the next three and a half months. The contents of three tubes of ventriculin (= 30 grams) were given daily in one dose, and the blood was examined regularly at weekly intervals. A week after the treatment with ventriculin was commenced the examination of the blood showed (June 27th):

Red cells, 1,670,000 per c.mm.; white cells, 2,000 per c.mm.; haemoglobin, 33 per cent.; colour index, 0.99; platelets, 31,000 per c.mm. Red cells well stained, with some variation in size and shape, some larger than normal and some smaller, but no nucleated reds seen; reticulocytes found only after prolonged search.

About two and a half weeks after starting on the ventriculin treatment the patient began to pass a little blood (approximately half an ounce) in the stools, and also some mucus. This continued almost every day for a fortnight. He caught a cold, herpes appeared on his

lips, and his general condition seemed to get much worse. This was confirmed by the blood examination, which showed (August 22nd):

Red cells, 720,000 per c.mm.; white cells, 2,200 per c.mm.; haemoglobin, 21 per cent.; colour index, 1.4; platelets, 13,680 per c.mm. Red cells well stained, with marked variation in size and shape; numerous megalocytes and microcytes; no nucleated reds seen, and reticulocytes found only after prolonged search.

On account of this very serious deterioration in the patient's condition it was decided to resort to blood transfusion, and two days later he was given about ten ounces of blood intravenously. There was some general reaction immediately following the transfusion, but five days later he seemed rather better, his colour had improved, and blood examination revealed (August 29th):

Red cells, 1,220,000 per c.mm.; white cells, 1,600 per c.mm.; haemoglobin, 22 per cent.; colour index, 0.9; platelets, 24,000 per c.mm. Red cells well stained, varying in size and shape; many megalocytes and microcytes present; no nucleated reds seen, and reticulocytes found only after prolonged search.

His cold improved, and the passage of blood in the stools ceased after the transfusion, but a little mucus continued to be passed. He was given a second blood transfusion (11 oz.) a month after the first, and shortly before the end of the course of ventriculin, and no general reaction followed. At the end of the three and a half months' ventriculin treatment blood examination revealed (October 3rd):

Red cells, 1,420,000 per c.mm.; white cells, 3,600 per c.mm.; haemoglobin, 34 per cent.; colour index, 1.1; platelets, 29,820 per c.mm. Red cells well stained, with marked variation in size and shape; megalocytes and microcytes present; no nucleated reds and no reticulocytes found after prolonged search.

The weekly blood examinations failed to show that the ventriculin had at any time produced even the slightest improvement in the blood, and reticulocytes were always difficult to find. The changes in the blood, as they occurred at weekly intervals during the three and a half months when the patient was undergoing ventriculin treatment, are shown in the second part of Chart 1.

It was thus obvious that the ventriculin treatment was useless, even though it had been reinforced by two blood transfusions; and as previous liver treatment had also failed, the case presented a difficult problem. The presence of tenderness and of an indefinite small sausage-shaped mass palpable with difficulty in the left iliac fossa, associated as it was with the passage of mucus and some blood in the stools (even though small haemorrhoids were present), added to one's anxiety lest a carcinoma be present. Radiological examinations after a barium meal and a barium enema showed on one occasion some indefinite narrowing at the site, which a subsequent investigation did not confirm.

After consideration it was thought, on the whole, probably wiser to clear up the question by exploratory laparotomy. This was done (October 9th) by my colleague Mr. Lambert Rogers. Some adhesions were found round the caecum (the appendix had been removed six years previously), but no tumour was present in the colon or elsewhere, and none of the other viscera showed any abnormality. The abdominal wound healed quickly, except for a stitch abscess, which continued to discharge for about three weeks. Twelve days after the laparotomy, and twenty-one days after a blood transfusion, examination of the patient's blood showed (October 21st):

Red cells, 2,200,000 per c.mm.; white cells, 3,900 per c.mm.; haemoglobin, 41 per cent.; colour index, 0.9; platelets, 37,000 per c.mm. Red cells well stained, marked variation in size and shape; megalocytes and microcytes; no nucleated reds, and reticulocytes found only after prolonged search.

Ten days after this the blood count was (October 31st):

Red cells, 1,140,000 per c.mm.; white cells, 4,200 per c.mm.; haemoglobin, 30 per cent.; colour index, 1.3; platelets, 31,200 per c.mm. Films: no change from preceding examination.

Koessler, Maurer, and Loughlin,¹ as the result of observation and experiments on rats, suggested that chronic vitamin deficiency (A, B, and C) might play a part in the production of pernicious and other severe anaemias, and that vitamin A in the diet was essential to blood regeneration. It was therefore decided to try multiple vitamin therapy in this case.

The patient was given vitamins A and D in the form of radiostoleum, half a drachm twice a day, vitamin B (also A and E) as bemax, half an ounce once a day, and

a lower level had been reached than at any previous stage during the administration of liver, ventriculin, or of vitamins. The blood examination was as follows (March 6th):

Red cells, 1,150,000 per c.mm.; white cells, 3,000 per c.mm.; haemoglobin, 27 per cent.; colour index, 1.1; platelets, 21,850 per c.mm.; reticulocytes, 0.8 per cent. Red cells well stained, marked variation in size and shape; megalocytes and microcytes; no nucleated reds.

The patient was now given liver extract (= 1/2 lb. liver) and 30 grams (three tubes in one dose) of ventriculin daily; but he got worse, gradually sank, and, in spite of a blood transfusion (14 oz.), succumbed fourteen days later with a terminal broncho-pneumonia. On the day before death blood examination showed (March 20th):

Red cells, 660,000 per c.mm.; white cells, 1,100 per c.mm.; haemoglobin, 20 per cent.; colour index, 1.5; platelets, 7,800 per c.mm.; reticulocytes, 0.2 per cent. Red cells well stained, varying in size and shape; many megalocytes and a few microcytes present; no nucleated reds seen.

The changes in the blood as they occurred at weekly intervals during the three and a half months when the patient was undergoing multiple vitamin treatment, and up to the date of death, are shown in Chart 2.

COMMENTARY

This case presented the general appearances usually seen in Addisonian pernicious anaemia. It was that of a well-nourished man having a smooth skin with lemon-yellow tint, languid and rather indifferent to the progress of his chronic disease. The blood picture—with megalocytes, microcytes, and poikilocytes, high colour index, leucopenia, and no abnormal white cells—was also characteristic. Erythroblasts were found only once, but the case was not one of aplastic anaemia. The post-mortem examination revealed findings typical of pernicious anaemia;

CHART 1.—Showing the changes in the blood occurring while the patient was undergoing treatment with (a) liver and (b) ventriculin.

vitamin C in the form of one orange—later increased to two—daily. In addition, of course, he was receiving ordinary diet. This was kept up for over three and a half months (November, 1930, to February, 1931).

The day before the multiple vitamin therapy was begun the blood count was:

Red cells, 1,730,000 per c.mm.; white cells, 3,400 per c.mm.; haemoglobin, 30 per cent.; colour index, 0.9; platelets, 25,000 per c.mm. Red cells well stained, marked variation in size and shape; megalocytes and microcytes; no nucleated reds present, and reticulocytes found only after prolonged search.

Weekly examinations of the blood thereafter showed that from the beginning of the multiple vitamin treatment the red cells, haemoglobin, leucocytes, and platelets steadily increased for the next two to two and a half months. The haemoglobin, leucocytes, and platelets reached their highest limits exactly two months after the treatment had been commenced, and the red cells two weeks later. Blood examination at the end of two months' vitamin treatment showed (January 9th, 1931):

Red cells, 3,500,000 per c.mm.; white cells, 5,400 per c.mm.; haemoglobin, 73 per cent.; colour index, 1.05; platelets, 174,000 per c.mm.; reticulocytes, 0.8 per cent. Red cells well stained and showing variation in size and shape, a few tending to be larger than normal, and others definitely microcytes.

At the end of two and a half months' vitamin treatment the blood showed (January 23rd):

Red cells, 3,840,000 per c.mm.; white cells, 3,200 per c.mm.; haemoglobin, 69 per cent.; colour index, 0.9; platelets, 92,000 per c.mm.; reticulocytes, 0.7 per cent. Red cells well stained, but still marked variation in size; megalocytes and microcytes present.

The progress of the patient was thus far exceedingly satisfactory. At no time since the onset of his illness had he felt so well. The red cells, haemoglobin, and platelets had increased more than ever before, and on only one previous occasion had the leucocytes reached the same level. The reticulocytes, however, were always scanty; once only did they reach 1.8 per cent. Further satisfactory progress, however, was not maintained. The road to recovery had apparently reached its peak, and now began to decline steadily and rapidly. The multiple vitamin therapy was persevered with for another month, but the number of red cells and platelets, and the amount of haemoglobin, rapidly diminished; the blood examination made a few days after stopping the treatment showed that

CHART 2.—Showing the changes in the blood occurring while the patient was undergoing treatment with multiple vitamins and, at the end, ventriculin and liver extract.

there was a marked hyperplastic, and not aplastic (atrophic), condition of the marrow of the long bones, and a well-marked haemosiderin reaction was obtained in the liver, spleen, and kidneys.

An unusual feature of the case, however, was the presence of hydrochloric acid in the stomach contents. Fractional test meals were examined about a month after the course of liver treatment was begun in hospital, and also a few days after the beginning of ventriculin treatment. Both showed similar findings. The free HCl

and total acidity followed more or less normal curves. Free HCl went up to about 30 (N/10 NaOH) one and a quarter to one and a half hours after partaking of the meal, and the total acidity to 55. There was no odour in any of the samples, and no blood, bile, or lactic acid; starch was present up to one and a half hours, and resting juice amounted to 10 c.cm.

Throughout the long courses of treatment with liver, ventriculin, and multiple vitamins, reticulocytes never exceeded 2 per cent., and usually they could be found only after long searching. Under treatment with multiple vitamins the patient, for the first time, showed steady and marked improvement, which reached its maximum degree (3,840,000 red cells per c.mm.) at the end of about two and a half months. At this point, however, in spite of continuation of the treatment, he quickly relapsed, and died a month later with a red cell count of 660,000 per c.mm.

The case, therefore, was resistant to liver therapy and to treatment with desiccated extract of stomach tissue. Although a marked improvement took place under treatment with vitamins A, B, C, D, and E, it was not maintained; and knowing the tendency of the disease to show periods of natural remission even without treatment, one cannot avoid the conclusion that this may be the explanation, and that vitamin deficiency *per se* is not an essential factor in the causation of the disease. For two weeks before he died he was given both liver extract and ventriculin, but without effect. On three occasions also the other forms of treatment were reinforced by blood transfusion, but the progress of the disease was not influenced thereby.

SUMMARY

- (1) A case of Addisonian pernicious anaemia, which resisted treatment by (a) liver and (b) desiccated extract of stomach tissue, is recorded.
- (2) Treatment with multiple vitamins (A, B, C, D, and E) was tried, and improvement took place for two and a half months, but was not maintained, and relapse ensued.
- (3) Liver extract and desiccated extract of stomach tissue were then given, but the relapse progressed and ended in death.
- (4) Blood transfusions also did not influence the condition.
- (5) The improvement which took place under multiple vitamin therapy may have been but a natural remission of the disease, and so vitamin deficiency *per se* may not be an essential factor in its causation.

REFERENCE

- ¹ *Journ. Amer. Med. Assoc.*, 1926, lxxxvii, 476.

SUBACUTE BACTERIAL ENDOCARDITIS FOLLOWING REMOVAL OF SEPTIC FOCI

BY

LEONARD ABRAHAMSON, M.D., F.R.C.P.I.

PHYSICIAN TO THE RICHMOND, WHITWORTH, AND HARDWICKE
HOSPITALS; PROFESSOR OF PHARMACOLOGY, ROYAL
COLLEGE OF SURGEONS IN IRELAND

The eradication of septic foci, particularly in connexion with teeth or tonsils, is advocated in all textbooks of medicine. Such advocacy belongs to the realm of elementary teaching, for it is realized that a large number of cardiac patients owe their death to the supervention of subacute bacterial endocarditis, and it is presumed on reasonable grounds that infection in such cases derives from teeth or tonsils. Yet the removal of septic teeth or tonsils in cases of chronic heart disease does not constitute quite so simple a problem as appears on the surface. This is amply demonstrated by the following cases in

which subacute endocarditis supervened upon removal of septic foci—teeth (two cases) and tonsils (one case)—in circumstances which leave no doubt that the fatal issue was the result of removal of these foci.

CASE I

A man, aged 39, was seen by me on June 24th, 1925. He had been informed twelve years previously that he had well-compensated mitral regurgitation. From this he had experienced no symptoms whatever. In the middle of January, 1925, he consulted a dentist, who extracted five teeth under local anaesthesia. A week later six more teeth were extracted. Shortly after the extraction he noticed small red spots on his fingers. These were attributed by him to the effects of the cocaine used in the extractions. Soon afterwards he noticed stiffness of his calves and thighs. On April 25th he developed sudden loss of power of his right arm and leg, without loss of consciousness. When I saw him strength had returned to his limbs, but a number of petechiae were noted on his fingers. These were, he stated, similar to the earlier spots. His temperature showed an evening rise varying from 99.4° to 101°, while his pulse rate varied from 90 to 120. Examination showed the presence of mitral regurgitation, without enlargement of the heart. Blood culture yielded a growth of *Streptococcus viridans*.

The next event in the case was the appearance of tender nodules on a finger, on the right arm, and on the right buttock. Crops of petechiae appeared from time to time in various parts. On July 25th sudden pain was felt in the right shoulder. This lasted for two days, and was followed by a shower of petechiae in the region of the right clavicle and scapula. More tender nodules were observed on the left wrist, left leg, and buttock. On September 20th, 1929, intense pain was experienced in the back of the head. This was followed by unconsciousness and, in a few hours, by death.

CASE II

A male, aged 21, had a history of rheumatic fever in childhood. A year after this infection he was able to indulge in vigorous exercise without discomfort. After a time he hunted, and rode in point-to-point races with considerable success. In July, 1929, he obtained his degree, but failed to pass the medical examination for the British Army. Until this time he was quite unconscious of the presence of any defect in his heart. The result of his medical examination caused him considerable surprise, and he thereupon consulted a London cardiologist, who found a valvular lesion, and advised removal of tonsils, which were, he found, in a septic condition. The tonsillectomy was followed by pyrexia, and ten days later the patient returned to Ireland. On the evening after his return he had a violent pain in his right hip, and was then seen by Dr. Drennan, who has kindly permitted me to publish this case. Examination revealed a good deal of sepsis in the throat, swelling and tenderness of the right hip, a temperature of 103°, and a pulse rate of 120. An aortic diastolic murmur was heard.

After some days the throat condition improved and the swelling of the right hip subsided. The temperature came down, but failed to reach the normal, and for some weeks there was an evening rise towards 100°. Pain and swelling attacked a number of joints, and with this a progressive anaemia became apparent. Six weeks after his return to Ireland I was called to see the patient. The throat was still septic, the temperature was 102°, and the pulse rate 120. The patient looked extremely pale, almost ashen; a few petechial spots were seen on the legs, and there was considerable sweating. The heart showed moderate enlargement, with aortic and mitral regurgitation. Blood culture yielded a pure growth of *Streptococcus viridans*. The course of the infection after this date followed the lines previously exhibited. Remissions occurred in which the general condition improved and the temperature fell, though never reaching normal. After each remission an exacerbation was ushered in by joint-pains, weakness, increased pyrexia, and sweating. Cardiac enlargement became more marked, and anaemia more profound. After seven months death ensued from cerebral embolism.