A NEW METHOD OF TREATMENT OF THE FRACTURED CLAVICLE

BY

JAMES KEMBLE, CH.M., F.R.C.S.

CHIEF CLINICAL ASSISTANT, WEST LONDON HOSPITAL

The multiplicity of the methods employed for the treatment of the fractured clavicle bears witness to the want of satisfaction which is so often felt in dealing with this common lesion. Owing to its peculiar location and to the unique mechanical purpose which it serves, a clavicle which shows any residual shortening after fracture occasions a very appreciable impairment of the strength and functional utility of the upper limb for a considerable time after the accident. Moreover, since in women décolletage is an established fashion, any mal-alignment will jeopardize the prospects of cosmetic perfection. In introducing this new mode of treatment I am convinced, after reviewing my series of cases, of the simplicity of its application, of the comfort afforded the patient, of the more perfect alignment obtainable, and of the definite curtailment of the period of incapacity occasioned by the injury.

CLINICAL FEATURES

The cause of these fractures is almost invariably indirect violence due to falls on the hand, the elbow, or the point of the shoulder. It is applied, as a rule, to the under surface of the outer end of the clavicle in an upward and inward direction, the relative magnitude of the upward and inward components varying in different cases.

The actual fracture occurs most frequently at or near the junction of the inner two-thirds and the outer third. This is the point of meeting of the two curvatures of the bone. Moreover, the inner two-thirds of the bone, constituted by the anterior convex curve, is cylindrical in form, and consists of a compact layer of considerable thickness. It therefore resists a fracturing force more effectually. The fracture may be of the "greenstick" type, this being common in young children. The angulation is directed upwards and forwards. Of the complete varieties the line of fracture may be transverse or oblique, the latter being the commoner. In the majority of cases, the obliquity takes a direction from above downwards, inwards, and backwards. The position taken up by the bone ends relative to one another is due partly to the direction of the violence and partly to the weight of the limb. The inner end of the outer fragment comes to lie below and in front of the outer end of the inner fragment. The whole outer fragment also moves inwards, so that the bone ends overlap. Other deformities occur, but are less common. In the most common type of lesion—the oblique fracture with the deformity described above—the part played by the violence in determining the direction of the fracture and the displacement of the ends is probably the same as in the case of the abduction-eversion fractures at the ankle. In the case of the clavicle, the coraco-clavicular ligaments act as a fulcrum, and the force applied to the under surface of the outer end of the bone produces an oblique fracture to the inner side of the ligaments, with the characteristic displacement. In the case of the ankle the abduction-eversion force driving the astragalus outwards against the external malleolus causes an oblique fracture of the fibula some two inches above the inferior tibio-fibular ligaments, which act as the fulcrum. The other factor producing and maintaining this position of the bone ends is the weight of the limb. The whole outer fragment drops. Moreover, as the main function of the clavicle is to buttress the shoulder outwards and backwards, when a fracture occurs, the outer fragment of necessity moves inwards and forwards.

TREATMENT

A satisfactory method of treatment must take cognizance of every one of these facts regarding the fracture. In dealing with all fractures the three principles of treatment are: first, the reduction; secondly, the retention of the corrected position; and thirdly, the maintenance of nutrition and early restoration of functional utility.

In the case of the clavicle, reduction is always easily accomplished. It is the retention which has always presented the difficulty.

To retain the position after reduction, some means must be utilized which will prevent the outer fragment from moving inwards, forwards, and downwards, and will at the same time act as a splint between the two portions of the divided bone. Moreover, the upper limb must be left free. In many of the methods which have been used, especially that known as Sayre's, the whole upper limb, including the hand, was fixed against the side and front of the chest by firm strapping or bandage. The patients always complained of great discomfort, the skin became sodden from perspiration, or blistered from strapping, while the greatest objection of all was the stiffness of the whole arm and hand, which was the outcome of the immobilization and disuse. This prolonged the period of incapacity very appreciably. Other methods, such as those depending merely on loops passed round the shoulders and tied across the back, were ineffectual. A secure and effectual retention of the reduced fracture must be ensured, not only on account of the great likelihood of recurrence of the deformity in this situation owing to the mechanics of the part, but also owing to the saltatory and restless disposition of the young patients in whom fractured clavicles are most common. The method must be essentially simple in application, owing to the ubiquitous incidence of this type of fracture and the everyday circumstances under which it occurs. It must be available to the general practitioner as well as to the surgeon. For this reason any elaborate appliance which involves the use of steel frames, fitted splints, or extensive plaster jackets is impracticable. Recumbency is undesirable. Open reduction with the introduction of a plate or intramedullary peg, though it may give ideal alignment, is as unnecessarily drastic as it is wantonly disfiguring.

I have therefore attempted to meet these exigencies by the utilization of a moulded plaster-of-Paris slab. The technique is as follows:

Make a fairly thick elongated slab by sandwiching several layers of plaster-of-Paris bandage between two layers of lint. This slab is roughly rectangular in shape,

FIG. 1.—Diagram demonstrating the moulded plaster slab in position.

but is somewhat wider at the ends than at the centre. It is made to an exact size for each case. Its length is one inch greater than the distance between the deltopectoral folds in front of each shoulder. It is 4 inches wide in an adult and proportionately narrower in infants. When the precise position of the fractured ends in the radiogram have been recognized the patient is seated on a backless chair. An assistant stands behind the patient, puts his left knee between the shoulder-blades, and with the fingers of either hand in front of the shoulders, draws them firmly backwards, outwards, and upwards. The

surgeon stands in front and manipulates the fragments till reduction is perfect. The assistant retains this by maintaining the traction.

The plaster slab is then applied by the surgeon in front. Each extremity is moulded so as to be crutch-ended and to fit into the delto-pectoral hollow in front of each shoulder. It is made to come well down into the axilla beneath the shoulder so as to obtain sufficient upward lift. Owing to the prearranged length of the slab, the central part, which is in front of the sternum and inner portions of the clavicles, is bowed out from the skin surface slightly. Over the outer half to two-thirds of the clavicle the slab is moulded exactly to the contour of the bone in its corrected alignment. It thus embraces the fragments on both sides of the fracture. The plaster is held till set. A nurse then applies bandages in the following fashion. First, a double "figure of eight" is applied round both shoulders, with the intersection in front over the centre of the slab. The assistant then relinquishes his hold. The effect of this bandage, very firmly applied, is to exert a force backwards, outwards, and upwards on the centre of the bow-shaped slab. The plaster here stands off a short distance from the sternum, so that this pressure is transmitted through the crutch-ends of the slab and *pushes* the shoulders and outer ends

FIG. 2.—Photograph of a patient with the moulded plaster slab in position.

of the clavicles backwards, outwards, and upwards. The slab acts, indeed, exactly as an adventitious clavicle, replacing the bone both in position and function as closely as is possible. It constitutes an efficient substitute till the continuity of the prototype is restored. Properly applied, this bandage is sufficient, for the slab maintains the shoulders in the braced back and supported position, when the assistant relaxes his hold. It is best, however, in order to ensure immobility of the plaster, to apply another "figure of eight" bandage round the shoulders, with the intersection behind. The portion of the slab which has been moulded so as to lie in intimate coaptation with the clavicle over the site of fracture acts as a perfect splint, preventing any possible movement at this point. The arm is supported in a sling at first, though this is not always necessary, and the patient goes home. The movements of the shoulder are partly restricted, but the rest of the arm and hand are perfectly free for use. The plaster is retained for from ten to fourteen days. Throughout, the patient is perfectly comfortable, and on removal of the slab finds he has full movements of the whole limb.

In the series of cases treated by this method a radiogram has been taken in every patient after reduction and application of the plaster. This has confirmed a satisfactory reduction.

SUMMARY

The advantages and results can therefore be summarized thus:

The method is simple.

It is effective in securely retaining perfect alignment after reduction, so that a gratifying cosmetic result is assured, and there is no risk of disability due to residual shortening of the bone.

The patient is completely comfortable throughout the treatment, and has free use of his hand and arm.

As a result of this last circumstance the period of incapacity due to the accident is considerably curtailed by the employment of this technique.

For the opportunity of carrying out this investigation I wish to express my gratitude to Mr. Sangster Simmonds, whose clinic at the West London Hospital these patients have attended.

RENAL COLIC AND HAEMATURIA FOLLOWING RECUMBENCY

BY

W. ETHERINGTON WILSON, F.R.C.S.

TORQUAY

During the latter period of the great war many of the hospitals were made into centres where certain types of gunshot wounds were concentrated for reasons of efficiency and team work. The 1st London General at Camberwell became one of the centres for fractured femurs, and it was among these cases that I came across a remarkable series of cases of renal colic and haematuria. Of some hundred and fifty men who passed through the special femur wards I have notes of twenty-three who exhibited this renal complication. I publish them now because I have often been asked about them, and hope that they may be of some assistance to research workers on kidney disease and function.

I have to admit from the outset that many investigations were incomplete and left undone at that time in trying to find a cause for the colic and bleeding, but I am convinced that, had *x* rays been used, a larger number of urines examined carefully, or cystoscopies undertaken, no assistance would have been forthcoming in elucidating these phenomena.

It is necessary to draw attention to the following facts.

(1) These were all femur cases, and were treated by the Sinclair method. (2) All the patients remained recumbent on their backs for periods varying from three to twelve months or more. (3) The foot of the bed was raised at least nine inches during the earlier weeks of their period of treatment—that is, the stage of extension. (4) Most of the patients concerned were kept recumbent for a long time on account of severe wounds, prolonged sepsis, delayed or non-union, or other wounds. Only one had a simple fracture with delayed union, and he was treated on a net bed, with the usual sloping feet-upward position. (5) When union was considered firm (normally two and a half to three and a half months) a walking calliper was fitted, and the men allowed up on crutches fifteen minutes or so on the first day. On the second day they were allowed to be up longer, and it was remarkable how much they did get about on their feet. I have known one of them to be seen walking in the corridor as if he had been up for a fortnight. Eagerness, competition, and the cheers and envy of the less lucky ones in the ward were, of course, responsible for these feats. It was on this day that several cases of renal colic with haematuria occurred. (6) With regard to the amount of sepsis present, small sinuses in the thigh, sequestra, and flaring-up of healed wounds were common occurrences. The more serious acute septic conditions—that is, secondary haemorrhages, blood infections, joint infections, etc.—did not occur in any of the present twenty-three cases under consideration. (7) Concerning diet, there was an abundance of greenstuffs eaten by the men at dinner, and it is perhaps wise to remember that during the period 1918-19 our vitamin stocks were probably somewhat lacking. (8) The total number of cases of renal colic and haematuria of which I have some record and