of opinion as to priority and the niceties of terminology, and to unite in a combined study of the problems before us, using a common classification for a starting point.

BIBLIOGRAPHY

- Bailey, P., and Cushing, H.: *A Classification of the Tumours of the Glioma Group*, Philadelphia, 1926.
 Cushing, H., and Bailey, P.: Haemangiomas of Cerebellum and Retina, *Arch. of Ophthalmol.*, 1928, lvii, 447.
 Cutler, E., Moritz, A., and Zollinger, R.: A Histological Study of Tumours of the Central Nervous System, *Ohio State Med. Journ.*, April, 1929.
 Globus, J. H., and Strauss, I.: Spongioblastoma Multiforme, *Arch. Neurol. and Psychiat.*, 1925, xiv, 139.
 Lindau, A.: Studien über Kleinhirncysten, *Acta path. et microbiol. Scandinav.*, 1926, Supplement i, 128.
 Penfield, W.: Principles of the Pathology of Neurosurgery, *Nelson's Loose Leaf Surgery*, 1927.
 Idem: The Encapsulated Tumours of the Nervous System, *Surg., Gynecol. and Obstet.*, 1927, xlv, 178.
 Idem: Chapter on Neuroglia, *The Cytology and Cellular Pathology of the Nervous System*, Hoeber, New York, 1930.
 Ribbert, H.: Über das Spongioblastom und das Gliom, *Virchow's Arch. f. path. Anat.*, 1918, ccxxv, 195.
 Roussy, G., Lhermitte, Jr., and Cornil, L.: Essai de classification des tumeurs cérébrales, *Ann. d'anat. path. med.-chir.*, 1924, i, 333.
 Sargent, P., and Greenfield, J.: Haemangiomas of the Cerebellum, *Brit. Journ. of Surg.*, 1929, xvii, 84.
 Strauss, I., and Globus, J. H.: Spongioblastoma with Unusually Rapid Growth following Decompression, *Neurol. Bull.*, 1918, i, 273.
 Tooth, H. H.: Some Observations on the Growth and Survival Period of Intracranial Tumours based on the Records of 500 Cases, with Special Reference to the Pathology of Gliomata, *Brain*, 1912-13, xxxv, 61.
 Weed, L.: Chapter on Meninges, *The Cytology and Cellular Pathology of the Nervous System*, Hoeber, New York, 1930.

SOME ASPECTS OF PREVENTION OF TUBERCULOSIS IN CHILDREN*

BY

F. MAURICE MCPHEDRAN, M.D.

(From the Henry Phipps Institute, University of Pennsylvania)

(With Special Plate)

In treating the important topic of tuberculosis in children, my point of view is necessarily much influenced by an interest, dominant during the past seven years, in the pathogenesis of the disease and in the course of its evolution. At the Henry Phipps Institute the incidence and progress of infections have been studied in family groups. No hospital beds are in the dispensary, nor have we access to a sufficient number of beds elsewhere to afford prompt hospitalization and sufficiently long treatment to many of our cases. This advantage, common to many communities, is peculiarly inimical to attempts to stamp out the beginning infiltrations of children. It is difficult at best to persuade an adolescent or the parents of a younger child, when symptoms are trivial or absent, that rest in bed is necessary to avert a serious illness. Delayed admission to an institution known to be crowded does not strengthen the appeal of an enforced rest. However, serial x-ray observations and correlated attempts to regulate the lives of children who defer or refuse institutional care are sometimes followed by retrogression of their lesions, and always place one in a stronger position to urge absolute rest in bed at the onset of definite symptoms. Furthermore, one is in a position to record the fluctuations of a lesion and to learn how extensive may be the damage capable of retrogression and healing without excessive surgery, essential and life-saving though surgery is in very many cases. Although one's observations are restricted through concentration on certain problems, it is clearly impossible

within a brief space to do more than outline a few aspects of tuberculosis in children, and to indicate means of detecting, preventing, and treating its protean appearances in the diverse phases of infancy, school age, and adolescence. It is only by discussion from many angles and by free exchange of the fruits of experience and controlled observation, that we may hope for "new methods of approach to a subject which still defies exact comprehension and as to which there is still so little unanimity in the minds of investigators."¹ The interaction of seed and soil,² of infecting micro-organisms and host, produces a baffling and complex disease. In its development and manifestations, the type of tubercle bacillus, the size, number, and spacing of the "doses" on the one hand, and on the other the portal of entry and the race,^{3 4} age,⁵ sex, and environment of the infected person, are leading factors. The bovine sources of contagion will not be referred to in this brief consideration. For those interested in pulmonary tuberculosis, in its pathogenesis, epidemiology and clinical manifestations, the duration and intensity of infection or contagion,⁶ and race and age, are perhaps the dominant considerations.

"The available amount of energy and money which can be applied to the extermination of tuberculosis is not unlimited, and so much of it as is spent on measures which are less likely to yield results than others must be considered to some extent to be wasted."⁷ News-holme's warning is not less significant to-day in view of the enormous economic waste of current methods. When the patient is ill enough to seek the physician the disease is usually so far advanced that months, even years, of treatment are a mere palliative. If the patient is retained in hospital, expensive facilities and care are required to secure a clinical recovery, which must be followed by years of careful rehabilitation; if he has a shorter inadequate hospital stay, he goes home a well of contagion, to infect those living in his environment. There is a tendency among those faced with these difficulties to find in the declining death rate a reassurance that only too often results in paralysis of effort. But the fall in mortality, the nature of which has been well defined by Cobbett,² should not be so interpreted that activities languish and fail to be progressively directed against the most destructive phases of tuberculosis. Moreover, amazing though the fall in the death rate is, tuberculosis mortality in England and Wales in infants under 1 year was 109 per 100,000 births in 1927,⁸ and in the age periods from 5 to 10 and from 10 to 15 years, although the death rate from all causes was low, tuberculosis accounted respectively for 14.43 per cent. and 22.53 per cent. of the deaths. But it is in adolescence and in early adult life that we find the peak of tuberculosis mortality, and here, too, we find in tuberculosis economically the most destructive and wasteful of diseases. It cripples more or less permanently or destroys at the beginning of usefulness the carefully trained economic unit representing years of education and the summation of family hopes. It strikes the patient down sometimes rapidly, more often in a series of relapsing illnesses. Further, during the quiescent intervals, the patient may pass on the infection to the next generation. Moreover, there is evidence that, at least in some large cities, there is not a diminution in the percentage of the population infected corresponding to the diminished total tuberculosis mortality.⁹ The tuberculin reaction shows that infection with the tubercle bacillus is still as nearly universal in city dwellers of European stock as it was twenty-one years ago.¹⁰ It is impossible to ascertain how many of these infections would flare up under conditions of unusual strain or privation, as is suggested by the steep ascent of the tuberculosis mortality curves^{11 12 13} in civil popula-

* Read in opening a discussion in the Section of Tuberculosis at the Annual Meeting of the British Medical Association, Winnipeg, 1930.

tions during war. In normal circumstances, many latent infections escape attention because good living conditions constitute an undesigned but adequate prophylaxis for this group. That other large group of cases which even under normal conditions develop into demonstrable latent infiltrations is the chief object of the search and attack in tuberculosis control.

HOUSEHOLD CONTAGION FOCUS OF STUDY, PREVENTION, AND TREATMENT IN TUBERCULOSIS

In seeking the most practical attack on infantile and adolescent mortality (and knowing that the combating of the latter is often grounded in the lesion, latent or clinical, of adolescence), a most rewarding procedure is to be found in the serial examination of those living in household contact with sputum-positive tuberculosis. We have used the term "latent" somewhat arbitrarily, but with a meaning that is appropriate to case-finding methods, to designate the pre-clinical stage of a tuberculous pulmonary infiltration, when it is demonstrable by *x*-ray examination but does not yet manifest itself clinically by symptoms or signs perceptible to the patient or by the usual methods of classical physical examination. In a report⁶ based on observations extending over slightly more than two years, it was shown that among a group of children of whom either the father or mother had positive sputum, about 9 per cent. showed clinical tuberculosis, about 2 per cent. well-defined latent pulmonary infiltrations, and about 20 per cent. calcifications of the tracheo-bronchial lymph nodes. In these children the number of pulmonary lesions has increased during four years of subsequent observation. Among a control group of children living in households no member of which had a history of pulmonary tuberculosis, there was no instance of clinical tuberculosis, and only about 4 per cent. showed calcifications of the tracheo-bronchial nodes. Among the first 102 families reported, 34 had one or the other parent sputum-positive, 33 had no member with tuberculosis, and in 7, one of the children (the eldest in five instances) had open tuberculosis. Although it is possible that in these seven cases the source of infection of the patient was extrafamilial, we were not able to examine the parents for pulmonary infiltration or to obtain adequate evidence of the cause of death of parents not living.

The first examination of contacts of a freshly reported case of open tuberculosis often discovers remarkably little evidence¹⁴ of severe infections which reward supervision. Sometimes, indeed, even the tuberculin reactions may be slight or absent in certain children. If the tuberculin test is made intracutaneously with measured quantities of a standardized tuberculin,¹⁵ and reactions occur only with the larger quantities (0.1 mg. and 1.0 mg. O.T.), strong evidence has been obtained that the exposure to positive sputum is both recent and short, and, in substantiation of this, the *x*-ray picture may show in the source of contagion a lesion that is evidently wholly soft and recent, or may present evidence of an old lesion with recent exacerbation. If, in addition, roentgenograms of the children show no evidences of long-standing exposure—such as tracheo-bronchial calcifications, etc.—the bacillary content of some twenty specimens of sputum as measured roughly by the Gaffky count, and the frequency of expectoration, will give the nearest approximation to a measure of the intensity of exposure in such cases. If the sputum shows few bacilli and the cough is infrequent, it is probable that the sum of infection has not been severe. Nevertheless, because of the uncertainty of all these measures of contagion, the tuberculin test should be repeated in some four to six months, and neither satisfactory hygienic living conditions nor serial *x*-ray

supervision should be neglected. A family history is appended:

Father, aged 31, ill five weeks with cough, sputum, and haemorrhage; onset acute, of the type resembling gripe. *X* rays show a few old strands in the extreme apex, right and left, and in the right upper lobe flocculent spots becoming confluent to consolidate for some four centimetres beside the interlobar fissure in the axilla only. Twelve specimens of sputum negative, one positive, Gaffky IV. Patient hospitalized fourteen weeks after the onset of symptoms; died five months later of acute tuberculosis. Mother, aged 27, showed a small tracheo-bronchial calcification, but no pulmonary infiltration. Tuberculin positive to 0.1 mg. O.T. Santo, aged 8 years 7 months, showed a large pulmonary calcified nodule and large tracheo-bronchial calcifications. Tuberculin positive to 0.1 mg. O.T. John, aged 7 years 2 months, had a negative *x*-ray, and was negative to 1.0 mg. O.T. Carmella, aged 5 years 4 months, had a negative *x*-ray, and was negative to 1.0 mg. O.T.

Here the acute onset and the soft mottling and consolidation in the father and the negative tuberculin reactions of the younger children, one of pre-school age, suggest exposure of short duration. The relative infrequency of positive sputum indicates no great intensity of exposure. Possibly the tracheo-bronchial calcification and weak tuberculin reaction of the oldest child and the few old strands seen in the father's apices may, taken together, be interpreted as evidence that during the infancy of the eldest child the father had a slight lesion, which became arrested before the second child was born. Careful inquiry failed to discover that anyone else had lived with the family. That this family was not heavily infected during fourteen weeks' contact with the father is shown by a tuberculin reaction in the youngest child persistently negative annually for three subsequent years and by the occurrence in the second child of merely weak reactions to the highest doses (0.1 and 1 mg. O.T.) during the same period.

The immediate problem of supervision is, of course, the severity and length of infection to which any infants in the household have been exposed. In an infant, even a weak tuberculin reaction may signalize a dangerous infection. There is evidence⁵ that even young infants may recover from lesions resulting from transitory severe exposure or slight exposure lasting several weeks. But in all cases there is grave danger that even a small lesion, whether of the lung or of one or several tracheo-bronchial lymph nodes, may progress or that metastasis to other areas, in particular the meninges, may occur. Therefore every precaution should be taken to prevent reinfection and to maintain the infant in the best possible living conditions, preferably in its mother's care unless she is the source of infection, and to ensure that the mother has time and means to care for it properly. The importance of preventing reinfection or superinfection is to-day probably generally recognized. But perhaps it is not so generally understood how the attachment to the infant of the source of infection, whether parent, grandparent, aunt, or uncle, will often result in an interruption of the patient's isolation from the baby and all that concerns it. Aseptic technique and clear appreciation of possible modes of transmission of infectious material come hard even to some physicians. In the absence of exact knowledge of the chances of infection—for example, by dry dust and by contamination of food and utensils and toys—merely to require that the patient with the open lesion shall not handle the baby is to take an indefensible risk with the infant's life. Although a positive tuberculin reaction in an infant is not conclusive evidence of immediate contact, it is sufficient to warrant the institution of every measure to ensure the prevention of further infection, including *x*-ray examination of all persons who have been in its environment.

At the other extreme is the result of the first examination of the contacts, typically father or mother, of a patient. One may have coughed and expectorated for years, with sputum presenting many bacilli in most specimens, and an x -ray film showing an old lesion, partly strand-like, with one or many spherical or multilocular smooth-walled cavities. There may be a history of a baby recently dead of meningitis. Of the younger children, half may show calcified tracheo-bronchial nodes, and of those past or at puberty usually one or more will show an infiltration of the upper third of the lung field, with or without symptoms and classical physical signs. The tuberculin reactions vary in intensity, but all, or all but one or two, react to 0.01 mg. O.T. The following cases are illustrative.

CASE I

Father, aged 39, sputum-positive, with symptoms extending back some seven years and an old strand-like lesion, largely excavated. Mother, aged 41, had no pulmonary lesion. Her tuberculin reaction was positive to 0.01 mg. O.T. At the first visit all eight children had positive reactions to 0.01 mg. O.T. A child 2½ years old had recently died of tuberculous meningitis. X rays showed that the three oldest children, aged 17, 16, and 15 years, had latent apical infiltrations, the third with associated tracheo-bronchial calcification suggesting, in conformity with observations in our studies of contagion, that her first infection was of intrafamilial origin. (The significance of calcification of these lymph nodes has been fully discussed elsewhere.⁵) Within three years from the first examination the oldest child, refusing rest in bed, became far advanced. The apical lesion of the next child at first advanced, but was improved by sanatorium care, unfortunately ambulant. The apical lesion of the third child (Figs. 1 and 2, Special Plate) progressed, despite a gain in weight, during ambulant sanatorium care. Another child in whom a fresh apical infiltration developed, with rapid spread, has improved remarkably with complete rest in bed (Figs. 27, 28, and 29 of Reference 5).

CASE II

The father of another family refused examination. The mother died of tuberculosis in 1918. The oldest child, a girl, died, also of tuberculosis, in 1921. The next, a girl aged 20, our first patient from this family, had a far advanced lesion with tracheo-bronchial calcification. The next child, a sister aged 17, had a latent apical lesion. The remaining three children, aged 15, 13, and 12, all showed tracheo-bronchial calcifications. The tuberculin reactions were all positive to 0.01 mg. O.T. Within one year from the beginning of our supervision the youngest child developed a severe infiltration, which has yielded to complete bed rest. (Figs. 3 and 4, Special Plate, and Figs. 16, 17, and 17A of Reference 5.)

These few diverse instances indicate at once how various and how logical may be the results of the first examination of contacts. The children of each household present evidences of infection that are, broadly speaking, reflections of the history and of the anatomical extent and condition of the lesion of the source of contagion. The correlation is clearer when a parent is the patient and the household has not at one time or another contained brothers, sisters, father, or mother of the patient—that is, when other and earlier sources of infection may be ruled out. In nearly all households, whatever the source of infection, the danger is immediate for children under 2 years, somewhat less urgent but more protracted for adolescents, and for children between these ages real, but for the most part deferred until puberty. In short, in children, as Newsholme⁷ showed to be the case in adults, the latent period between the cessation of exposure and the onset of a clinically manifest lesion is often a matter of years.

An eldest daughter, all members of whose household had been serially examined (Fig. 11 of Reference 6), first showed an apical infiltration four and a half years after her father's death. Again, a boy, whose sputum-positive mother died

in his sixth year, first showed symptoms in his fourteenth year. In this somewhat unusual case there is no evidence of other intrafamilial contact. Strong evidence that there was none is found in the negative tuberculin reactions of three younger brothers, who were aged 9, 7, and 5, within less than six months before the appearance of a positive sputum in this patient. No extrafamilial contact, always a possibility to be considered, could be traced.

It is not uncommon for children past puberty first to present infiltrations from the second to the fourth year after the cessation of contact. Jessel¹⁶ found three and a half years the average interval. I have not found that annual x -ray examinations after contact has ceased could be dispensed with during a period covered by seven years' supervision of contact children. Supervision should be closer and more prolonged the nearer the period of contact to the years of adolescence. Known contagion, centring about a diagnosed case of sputum-positive tuberculosis, is the most economic starting point for discovering pulmonary tuberculosis in its most tractable phase. X -ray examination of school children^{9 17} selected by the tuberculin reaction and the routine examination of young adults seeking employment¹⁸ stand perhaps next in order of demonstrable value. X -ray examination of all school children, though more laborious and expensive, has the added advantage that it will reveal many chronic non-tuberculous pulmonary infiltrations, which, undetected and untreated, are productive of serious invalidism. Furthermore, adequate films will pick up indications, in heart size and shape and in pulmonary markings, of many cardiac lesions needing supervision that would nevertheless be overlooked by routine physical examination.

X RAYS IN DIAGNOSIS, SUPERVISION, AND CLINICAL INVESTIGATION

In a general way the importance of the x rays in the examination of the lungs is recognized. It is less generally known or admitted that they can afford basic information about the state and possibilities of a lesion, information that is subject to modification by evidence such as elevated temperature and various abnormal blood pictures, but is nevertheless in itself sufficient warrant for the institution and continuance of treatment in the absence of corroborative accessory findings. Ignored, the x -ray evidence will sooner or later receive disastrous amplification in signs, symptoms, and laboratory tests.

In earlier papers^{19 20 5} histories are cited and lesions recorded, particularly in Figures 5, 14, 16, and 17, and 24 to 35 of Reference 5, illustrating how rapidly, how extensively, and for how prolonged a time tuberculous infiltrations can invade the lung substance and even destroy it while remaining latent in the sense defined above (Figs. 1 and 2, Special Plate). More recently, comparative studies, as yet unpublished, have shown that, although the blood picture^{21 22 23 24} or the sedimentation test, or both, when repeated at relatively short intervals, rarely fail to reflect definitely progressive infiltration, both may be normal at a time when the x ray records a lesion the softness of which indicates an instability subsequently verified by spread. We should be most unwilling to dispense with the aid of either of these valuable indications of activity, properly defined. We hope that through extended comparative studies, subjected to continued analysis, they may throw further light on the course, particularly, of latent lesions, and furnish more certain criteria on which to base their care. But we do not yet know in what relation to place the evidence afforded by each of them in its various degrees of positiveness, and it is essential to evaluate all evidence carefully in dealing with a disease at once so insidious and so volcanic as tuberculosis. Having found that the

x-ray evidence may be alone decisive, we feel that in all cases, at least for the present, the anatomical extent, site, and condition, or softness, of the lesion as revealed by the *x* ray should be the major consideration, emphasized by these important but accessory tests when positive. This distinction should, above all, be insisted upon during the retrogression of infiltration, when only the *x* ray may reveal the extent of the lesion and lack of delimitation, denoting capacity for violent resurgence.

Economically, the greatest waste in the recognition and supervision of tuberculosis arises probably from the failure to institute promptly, and to continue, treatment for the lesions of children, particularly adolescents, until they show, by serial *x*-ray examinations, clear evidence of strand-like anatomical stability. To discharge adolescent children, symptom-free for a few weeks, without such results, is a cat-and-mouse form of treatment, building up hope over months, to destroy it over weeks or months or years. Nothing is more futile and visionary than to lavish attention and costly infirmary beds only upon the clinically manifest activity of so insidious a disease. Inexpensive accommodation can provide adequately for much of the stabilizing period of anatomical arrest if medical supervision is equipped with good *x*-ray and laboratory facilities, and informed by experience with tuberculosis and the tuberculous. Unquestionably, if there is evidence by *x* ray, sedimentation test, or differential count that a lesion is increasing, the patient should be returned to an infirmary bed. Infirmary beds should likewise be sufficient in number to permit alternating periods of complete rest in bed and of varying degrees of freedom for those whose lesions are only moderately well delimited and sluggish, showing little progress to stable strand-formation, despite, perhaps, negative blood tests. No doubt there will still be found those who, objecting that to give such a place to the skiagram is to allow it to define activity and so to determine treatment, will boggle at such an idea, largely because of the association of activity with venerated definitions. But definitions that are time-honoured may come to be time-worn. The value of any conception or definition of activity must depend on the accuracy with which it points to the course and potentialities of the morbid process. Before the development of dependable *x*-ray methods and the correlation of skiagraphic appearances in the living with those of excised material subsequently sectioned, certain physical signs, in particular rales, formed for many physicians the most dependable evidence of active tuberculosis. Physical signs had to be discarded as the index of activity because, on the one hand, they usually lag far behind the initial invasion and the development of fresh infiltrations, and, on the other hand, they may persist for years over anatomically stable and functionally cured lesions. Elevation of temperature and of pulse, the signs currently accepted as evidence of activity, are significant only in that they usually indicate an acceleration, intensification, or extension of the existing lesion. Like rales and symptoms, they neither coincide with the onset nor, by their disappearance, signalize the cessation of progressive infiltration or excavation. Alike in the latent or pre-clinical phase of pulmonary tuberculosis and during clinical disease, widespread extension and even excavations are commonly recorded, particularly in children, without changes in temperature and pulse. These are usually late and essentially secondary effects of the morbid process. When present they occur with no definite, much less proportionate, relation to the intensity and extent of the underlying infiltration, the means by which tuberculosis destroys lung and eventually life. Attempts to link these signs directly with phases of allergy, immunity, or lack of immunity, or with the so-called exudative stage of

tuberculosis in children will receive scant support if closely spaced serial films are made of symptomless contacts (Figs. 1, 2, 3, and 4, Special Plate). (See also Figs. 5, 14, 16, 17, 17A, 20, 24, 27, 28, 30, and 31 of Reference 5.)

A pulmonary infiltration sufficient in extent to cause fever, or other signs and symptoms, rarely, if ever, escapes ready recognition by an adequate roentgenogram interpreted with regard to the anatomy of the chest, the pathology of tuberculosis, and the properties of the *x* rays.^{5 25} In children and adolescents particularly, tuberculosis characteristically causes extensive damage with no febrile disturbance, or with such transient and trivial disturbances as commonly occur in healthy children. A conception of activity in which emphasis is transferred from the direct anatomical evidence presented in the *x*-ray film to such secondary haphazard effects, has lost the essential protection it can give the patient, and will allow negative evidence to camouflage destruction. A patient in whom a tuberculous pulmonary infiltration appears in the *x*-ray film as soft ill-delimited spots, especially if these tend to confluence, should be treated by rest in bed until the insufficiency of such care is proved by adequate observation. As long as such a lesion, roentgenographically recorded at three months' intervals, continues to show a material retrogression from one quarterly record to the next, the patient must be regarded as one who still needs complete rest in bed. Such a lesion is clearly very unstable. An ambulant patient should be one in whom the lesion is not obviously ill delimited and, in quarterly roentgenographic records, either continues to show retrogression or maintains anatomical stability. Even without the warning of other signs of activity, such as may be found in the blood test, the sedimentation test, temperature, or by physical examination, a patient is not suitable for ambulant care who shows a soft ill-delimited consolidation or mottling, despite evidence of retrogression in quarterly roentgenograms.

Eligibility for discharge should be based on the degree to which the infiltration is reduced to sharp strands and calcium. The younger the patient, the less are the therapeutic and economic obstacles to adequate treatment. Particularly in adolescents and young adults the volcanic character of this disease, its potentiality in them for swift relapse on the one hand and relatively rapid retrogression on the other, should be recognized by the rigid application of a roentgenographic standard of anatomic stability. The younger the patient the more he has to lose or to gain by these alternatives. By so much the more, therefore, should we provide that he is not discharged from close supervision before his lesion is in a stage to support the stresses that heedlessness and economic necessities will bring upon it. Serial roentgenograms are the justification and guide for treatment, and the patient's best protection. Boards of trustees and managers have been known to object to the adequate use of the *x* rays because of cost, which is equivalent to the charges for about three days' care, or, for quarterly films, to twelve days' sanatorium residence in the year. The implication is that merely being in a sanatorium for 365 days in a year is more important than having the medical care accurately adjusted to the needs of the disease every three months and keeping the patient in sanatorium only 353 days of the year. Except for the moribund, is there any group of tuberculous patients who cannot better afford to dispense with a small fraction of their sanatorium stay than to suffer the hazards of uninformed treatment? It is implicit in the foregoing that one must not expect some two or three months' rest in bed to nullify an extensive soft lesion merely because

it has not caused signs or symptoms. There is evidence that of two children with lesions of equal extent the one without symptoms or signs clears more rapidly and completely than the one with clinically manifest disease, but even with the former, time is needed to secure a stable lesion. When unsupported by the warning of symptoms the physician requires, to put the patient in bed or to keep him in bed, the ability to attain, at more appropriate seasons, the Fat Boy's desire, "I wants to make your flesh creep." It is only by the sufficiently early institution of treatment long enough continued that he can succeed in preventing and checking tuberculosis in children and adolescents, and so in forestalling the greater part of the tuberculosis of early adult life. To achieve and establish the benefits of preventive medicine in this difficult but most rewarding field, where the prevention of clinical disease is in turn a means of preventing the spread of infection, even into the next generation, he needs all his resourcefulness, tact, optimism, and the sharpest discrimination in evaluating evidence.

Note.—The use of B.C.G. involves so many considerations that it has not seemed desirable to discuss it in this paper. The figures of Cox²⁶ and his co-workers give a much lower mortality among children intimately exposed to sputum-positive tuberculosis than are given by French workers. We have reserved B.C.G. for infants born into households containing a person with positive sputum, as a measure whose exact value has not yet been deter-

mined. The number of such infants is not large, but, especially if the intelligence of the family is low, the danger is grave.

REFERENCES

- ¹ Cummins, S. L.: *Proc. Roy. Soc. Med.*, 1930, xxiii, 47.
- ² Cobbett, L.: *The Causes of Tuberculosis*, Cambridge, 1917.
- ³ Borrell, A.: *Ann. de l'Inst. Pasteur*, 1920, xxxiv, 105.
- ⁴ Opie, E. L., Landis, H. R. M., McPhedran, F. M., and Hetherington, H. W.: *Amer. Rev. Tuberc.*, 1929, xx, 413.
- ⁵ McPhedran, F. M.: *Ibid.*, 1929, xx, 532.
- ⁶ Opie, E. L., and McPhedran, F. M.: *Ibid.*, 1926, xiv, 347.
- ⁷ Newsholme, A.: *The Prevention of Tuberculosis*, E. P. Dutton and Co., New York, 1908.
- ⁸ Dow, D. J., and Lloyd, W. E.: *Tuberculosis Mortality in Children*, Richard Clay and Sons, Ltd., 1930.
- ⁹ Hetherington, H. W., McPhedran, F. M., Landis, H. R. M., and Opie, E. L.: *Amer. Rev. Tuberc.*, 1929, xx, 421.
- ¹⁰ Hamburger, F., and Monti, R.: *Münch. med. Woch.*, 1909, lvi, 449.
- ¹¹ Kirchner, M.: *Zeit. f. Tuberk.*, 1921, xxxiv, 228.
- ¹² Glover, E. G.: *Lancet*, 1921, cc, 715.
- ¹³ Drolet, G. J.: *Bull. New York Tuberc. Assoc.*, 1923, iv, No. 5, 13; *Amer. Rev. Tuberc.*, 1924, ix, Abstracts Section, 41.
- ¹⁴ Wood, W. B.: *Tubercle*, 1930, xi, 489.
- ¹⁵ Aronson, J. D.: *Amer. Rev. Tuberc.*, 1926, xiii, 263.
- ¹⁶ Jessel, G.: *Tubercle*, 1930, xi, 493.
- ¹⁷ Hetherington, H. W.: *Amer. Rev. Tuberc.*, 1929, xx, 511.
- ¹⁸ Reid, A. C.: *Ibid.*, 1929, xx, 46.
- ¹⁹ McPhedran, F. M.: *Ibid.*, 1927, xvi, 420 and 479.
- ²⁰ *Idem*: *Tubercle*, 1928-29, x, 14.
- ²¹ Medlar, E. M.: *Ibid.*, 1927-28, ix, 218.
- ²² Blackfan, K. D., and Diamond, L. K.: *Amer. Journ. Dis. Child.*, 1929, xxxvii, 233.
- ²³ Sabin, F. R., Cunningham, R. S., Doan, C. A., and Kindwall, J. A.: *Bull. Johns Hopkins Hosp.*, 1925, xxxvii, 14.
- ²⁴ Cunningham, R. S., Sabin, F. R., Sugiyama, S., and Kindwall, J. A.: *Ibid.*, 1925, xxxvii, 231.
- ²⁵ McPhedran, F. M., and Weyl, C. N.: *Amer. Journ. Med. Sci.*, 1927, clxxiii, 313.
- ²⁶ Cox, G. L.: *Tubercle*, 1929, x, 497.

DIVERTICULA OF THE DUODENUM

BY

S. GILBERT SCOTT, M.R.C.S., L.R.C.P.

D.M.R. AND E.CAMB.

CONSULTANT RADIOLOGIST TO THE LONDON HOSPITAL

(With Special Plate)

THE clinical diagnosis of diverticula of the intestinal tract, excluding possibly those of the proximal and distal sections, the oesophagus, and colon, may be said to be practically impossible without the help of an opaque meal investigation.

By this method these pouches or local bulgings of the intestinal wall have been detected on almost all parts of the intestinal tract; as curiosities, I am reproducing radiographs of minute diverticula of the ileum and appendix (Figs. 3 and 4). The latter brings to mind the old rhyme, "Great fleas have little fleas upon their backs to bite 'em, and little fleas have lesser fleas, and so *ad infinitum*," for the appendix itself may be considered a diverticulum. I have been unable to find any published record of diverticula in these two sections of the intestinal tract.

A definite case of diverticulum of the duodenum, containing stones found by chance at necropsy, was described by Chomel as early as 1710—that is, in the reign of Queen Anne. About one hundred cases had been described up to 1910, none of which were diagnosed clinically. Since the opaque meal examination became possible, the literature on the subject has become considerable.

While granting that diverticula may be present in large numbers in the colon without giving rise to symptoms, my experience leads me to think that those arising from any section of the duodenum eventually produce some form of abdominal disturbance or distress, whether from direct or reflex causes, and in some cases this may even be of an acute nature, although more frequently it is met with as a chronic condition (diverticulitis duodenale). In fact, I now include a duodenal diverticulum in the

list of undiagnosed causes of the condition known as "chronic abdomen," which is, in my experience, more often than not, due to one of the following, placed in order of their frequency: (1) duodenal ileus, (2) splenic drag (incomplete enteroptosis), and (3) duodenal diverticulum.

The early diagnosis of any of these is important, for it is in the early stages of development that measures must be taken if the patient is to be prevented from joining the large and pathetic tribe of "chronic abdomens." The majority of these unfortunate cases are but undiagnosed or undiagnosable abdominal lesions.

It will be realized, therefore, that diverticula, especially those arising in connexion with the duodenum, have ceased to be only of academic interest, and call for much more careful consideration and investigation than has hitherto been given them.

It is usual to describe two types of diverticula—the congenital and the acquired. The acquired—namely, those resulting from traction of adhesions, ulcers, etc.—should not, strictly speaking, be placed in the same category as the true or congenital type of sacculation.

All the normal structures and layers pertaining to the duodenum should be found in the true diverticulum, while the individual thickness of the wall is dependent on the degree of distension of the pouch.

As already mentioned, the clinical diagnosis is practically impossible, as the abdominal symptoms are so varied and at times resemble very closely other pathological lesions. In fact, most of the cases I have had to investigate have been sent with the provisional diagnosis of gastric or duodenal ulcer, cholecystitis, or appendicitis—practically the identical list of lesions that the symptoms in duodenal ileus suggest. The causes of symptoms are direct and reflex. As with an appendix, an acute or chronic infection or inflammation may occur. As a rule, however, these changes are only catarrhal, but in some cases suppuration, ulceration, and even perforation are met with.

It is a mistake to conclude that because a diverticulum, found by chance at a post-mortem examination, shows no

F. MAURICE McPHEDRAN: PREVENTION OF TUBERCULOSIS IN CHILDREN

FIG. 1.—Female, 16½, October, 1928. Right: calcified lymph node above 6th post. rib; faint inverted salients below neck of 2nd post. rib. Left: irregular soft clouding in post. 2nd, 3rd, and 4th I.S. FIG. 2.—May, 1930: shows increased lesion. History: December, 1926; delicate salients projecting below the neck of 2nd rib (r. and l.)—the only parenchymal lesion. February, 1928; lesion on left reached 3rd and 4th post. I.S.; no signs or symptoms. May, 1928: a few rales only. May-October, 1928, patient ambulant in sanatorium; gained 5½ lb., averaged 94 per cent. normal wgt.; infiltration steadily increasing. January, 1929; slight retrogression. May, 1930; definite rales; pulse and temperature normal. No symptoms.

Male, November, 1927, aged 11. X-ray showed calcified nodules and nodes; same picture March, 1928. September, 1928, TBN was + to 0.01 mg.; temperature 98.6°; wgt. 90 per cent.; no symptoms. October 1, cough; sputum ¼ oz.; five specimens negative. FIG. 3.—13/10/28. Right; 2 calcified nodules on 8th and 9th ribs near tip of 5th rib; with four associated nodal calcifications in hilum in 7th and 8th I.S. and on 8th rib; fresh, soft, unequal density in 5th and 6th post. I.S. Oblique films showed this as situated on post. chest wall. The small density in 5th post. I.S. close to spine, is due to a vessel. Left: granular nodal calcification in hilum at 7th post. I.S. Symptoms: morning cough only; patient in bed at home except for dispensary visits. FIG. 4.—Increased lesion with central excavation despite gain in wgt., absence of fatigue and fever, and decrease in cough; sputum negative; wgt. 97 per cent.; rest in bed continued. January, 1929, retrogression definitely begun. June, 1929, still improving.