The first principle of treatment, therefore, should be to avoid an open pneumothorax. This may be done by using a drainage tube with a valve, which allows the pus to leave but prevents air from entering the pleural cavity. It is usually better, however, to aspirate the fluid at first until adhesions have formed, when incision and open drainage can be performed if necessary, although a certain proportion of cases get well with aspiration alone. In practice it is found that pneumococcal empyema in lobar pneumonia occurs after the illness. The pus is thick, and there are adhesions, so that it is safe to resect a rib and drain the cavity. Moreover, by the time the empyema forms, the patient has, as a rule, recovered from the toxæmia of the pneumonia. For these reasons treatment by incision and drainage is usually successful in pneumococcal cases, and when once the pus is found to be thick, it is better to drain the empyema.

In streptococcal cases, however, the effusion forms early, whilst the patient is still suffering from the toxæmia of the disease. The fluid is at first cloudy and thin, and there are no adhesions, consequently incision and drainage is likely to be fraught with danger. Dr. W. J. Stone published the results of three series of cases.¹ In the first series 85 cases were treated by early operation and drainage, and the mortality was 61.2 per cent. In the second series 96 cases were treated by early aspiration and late operation, and the mortality was 15.6 per cent. In the third series 94 were treated by early and late operation, and had a mortality of 9.5 per cent. In some cases aspiration alone is sufficient, but in the majority subsequent drainage becomes necessary, and when once the fluid has turned into frank pus it is usually safe to operate. After operation, in every case of empyema, it is essential to wash out the pleural cavity frequently with Dakin's or some similar solution in order to encourage re-expansion of the lung, and to disinfect the pleural cavity. The principles, therefore, in the treatment of pyogenic empyema are:

1. Remove the pus, but always avoid an open pneumothorax.
2. Encourage the lung to re-expand and obliterate the empyema cavity. This should be done by washing out the cavity, and any drainage should be arranged so that air cannot enter from outside.
3. Sterilize the cavity by frequent irrigations.
4. Attend to the general condition and nourishment of the patient.

REFERENCE

- ¹ *Amer. Journ. Med. Sci.*, 1916, clviii, 1.

BATH TREATMENT FOR DEAFNESS

BY

JAMES ADAM, M.A., M.D.

SURGEON FOR DISEASES OF EAR, NOSE, AND THROAT, GLASGOW
CITY HOSPITALS

Any measure which will help, even in a minority of cases, what is commonly called chronic progressive deafness is worth consideration, especially if it sometimes succeeds where the usually approved methods fail, and if it is a simple procedure capable of supervision by the general practitioner. The following case illustrates the point.

CASE I

A man, aged 42, consulted me in December, 1927, about deafness in the left ear, which had increased during the previous two years. There was no obvious retraction or thickening of the tympanic membranes, but he had difficulty in self-inflation. With the right ear he could hear a whisper across the room (say 15 feet away); with the left ear, 5 feet away. After treatment by catheter, bougie, and inflation for a year, the condition was worse. A whisper was then heard by the right ear at a distance of 5 feet; by the left ear, 2 feet. He was ordered hot baths, with a pound of commercial Epsom

salts added to each bath; immersion for ten to fifteen minutes was advised at bedtime every second night for a week, once during the following week, and then once a fortnight. Within a week, the hearing distances for whisper were 15 and 9 feet (instead of 5 and 2 feet) for right and left ears respectively. By May, 1930, the distance for each ear had increased to 30 feet. Improvement is still maintained, and is valuable not only to himself but to his colleagues. The baths are still necessary, but they are taken at longer intervals.

As often happens with new treatment, this first case has perhaps been the most striking. The success induced me to try the method in other cases, and sometimes gratifying results were obtained.

CASE II

For many years the manager of a large business, aged 64, had increasing deafness for which I tried in vain the usual measures. Some years elapsed, during which he did not consult me, but after the experience mentioned I asked him to see me. On examination, the right membrane was dull and indrawn, the left was ruptured and atrophied, and the ossicles were bound in cicatrix. Conversation voice was heard 4 feet away on the right side, 4 inches away on the left side. The latter figure is naturally open to question, because of possible hearing by the right ear even when closed. After treatment by baths for two months, the distances were 10 feet and 2 feet respectively, to his great delight.

CASE III

A hospital matron, aged 33, had tinnitus in the left ear, and slight deafness. The Eustachian tubes were narrow; systolic blood pressure 100 mm. Hg. Resection of nasal septum, removal of tonsils, and passage of Eustachian catheter and bougie were all tried, with no obvious benefit. When the patient reached the age of 38, the tinnitus was worse and was in both ears. When, two years later, in May, 1930, a whisper was heard by both ears at 8 feet, I asked her to try baths. A fortnight later the figures were 21 and 25 feet respectively, tinnitus not having improved.

CASE IV

A lady, aged 60, complaining of deafness beginning in the left ear ten years earlier, came to see me in October, 1930. Both membranes were dull, and inflation by catheter was difficult and imperfect. She heard the conversational voice 15 inches away on the right side, and 5 inches away on the left; Weber referred to left. As there was no improvement on inflation, she was asked to try baths. At the middle of December, conversation was heard on the right side at 16 feet distance (instead of 15 inches), on the left at 10 inches.

CASE V

A man, aged 58, had noticed deafness and tinnitus in the right ear for five months. Both membranes were indrawn. A whisper on the right side was heard 2 inches away, but after three weeks of bath treatment the distance had increased to 7 feet; tinnitus was still present, but less marked.

The less chronic cases respond more readily. Those described above were probably "catarrhal" cases, with narrowing of the Eustachian tube—the type in which sweating might be expected to help. Encouraged, however, by the improvement in the cicatricial ear of the patient (Case II) mentioned above, I resolved to try baths in the following case:

CASE VI

A man of 39, who complained of deafness in the right ear after an explosion in the trenches in 1916, came to me in June, 1930; there was also tinnitus in the right ear. The left ear was normal; the right drum was "gone," ossicles adherent, no discharge. He was asked to try baths. The hearing for whisper before treatment was at a distance of 3 feet; after two months' treatment, at 60 feet.

One might expect benefit from this treatment in patients suffering from Ménière's complex, if it is tried early. I have had the opportunity in only two such cases, both late ones. In the first there was complete failure; in the second there seems to have been some success, as shown by the history.

CASE VII

A widow, multipara, aged 47, weight 12 st. 10 lb., height 5 ft. 3 in. was first seen in January, 1930. She had a typical Ménière's syndrome for three years, beginning in the left ear, in which she was now stone deaf. The right ear had been affected for two years. She had well-marked tinnitus and frequent attacks of giddiness on turning quickly; on going upstairs, she had to cling to the rail and sometimes fell. Tinnitus, giddiness, and deafness were worse at menstruation. She could just hear a loud noise with the right ear at 3 inches distance. The Wassermann reaction was negative. Baths did no good. They did not make her sweat till she added 2 lb. of magnesium sulphate to the bath. With this treatment, however, she persevered. She now hears loud speech at 5 feet, has no giddiness, and can walk up and down stairs with ease. The baths may have nothing to do with this improvement. The course of Ménière's complex is sometimes capricious, but anyone who has read Dederding's monograph¹ would consider treatment by sweating not unlikely to help; to be a real help it would need to be used regularly and before irretrievable damage is done to the inner ear.

This treatment is as yet in the tentative stage, but these notes may suggest to others that the method is worthy of trial. It is likely to succeed only in a minority of chronic cases, though it has had good results in cases in which the usual methods had proved an utter failure and in which treatment had been abandoned. Success is more probable in the less chronic cases and in acute cases. It fails when it fails to sweat the patient. Whether magnesium sulphate is essential I do not know; it is hygroscopic and has been used in packs for chronic rheumatism. Doubling the quantity of salt to the bath seemed effectual in causing the last-mentioned patient to sweat. Treatment by the emunctories is an old-established method, but far too little attention has been paid to the skin and to the effect of sweating on the mucous membranes; probably civilized peoples sweat themselves too little and pay a price in catarrhal affections. Thirty years ago treatment by pilocarpine was tried, but had only a short vogue as "the remedy was worse than the disease." Caution must be used with baths, too. Hot baths are exhausting and too risky for elderly, weakly, and cardiac patients; ten minutes immersion may be long enough, and should be followed by a brisk towelling and a warm bed.

REFERENCE

- ¹ Dida Dederding: *Morbus Menieri, Acta Oto-laryngologica*, 1929, Supplement x.

CHYLOUS ASCITES COMPLICATING ABDOMINAL LYMPHADENOMA

BY

THOS. A. PRATT, AND H. W. O. FREW,
M.D. M.D., D.P.H.

ASSISTANT PHYSICIANS, CITY OF GLASGOW FEVER HOSPITAL,
KNIGHTSWOOD

The presence of serous fluid in the peritoneal cavity may be due to a variety of causes, but it is very seldom that cases are reported in which the fluid is chylous in nature. One such case came under our observation, and was considered, on account of its rarity, worthy of record.

Case History

The history of the case dates from September, 1928, when the patient, who was a young man of 20 years, first complained of weakness, tiredness, and loss of flesh. Though able for a time to carry on his work, he had ultimately to give it up on account of the weakness. In January of the following year he developed an attack of influenza, and during convalescence from this illness noticed for the first time, owing to slight pain, a fullness of his abdomen. It was not, however, till coughing and spitting became troublesome that he

consulted his doctor. From the history given, the presence of tuberculosis was suspected, but examination of the chest revealed only a few moist sounds at the base of both lungs. X-ray examination confirmed the absence of any evidence of tuberculosis of the lungs. The sputum was examined, and was found to be negative for tubercle bacilli. From this time till his admission to hospital his condition rapidly became worse. The general weakness increased, the abdomen became much more distended, and he was in consequence confined to bed.

When admitted to hospital, in the middle of May, the patient's condition was obviously serious. He appeared thin, wasted, and cyanosed, and complained of pain in his left thigh and leg. The distension of the abdomen was marked, and gave rise to considerable respiratory embarrassment. In the feet and legs there was a moderate amount of oedema. Examination of the chest was negative, save for signs of congestion at both bases, and the sputum was negative for tubercle bacilli. The heart showed no evidence of organic disease. The urine, which was diminished in amount, contained albumin. During the succeeding days the oedema present in the legs spread upwards till even the chest wall was involved, and with the increase in the abdominal swelling the breathing became more distressed.

On May 24th, 1929, paracentesis was performed, and 26 ounces of milky fluid were removed from the abdomen. The procedure afforded relief, which, of course, was only temporary, and the operation had again to be performed. Altogether the abdomen was aspirated six times, and 38 pints of fluid were withdrawn. When the abdomen was palpated just after paracentesis a hard fixed mass was felt in the epigastric region. In addition to the large mass, smaller masses were felt throughout the abdomen. In spite of these aspirations his condition gradually became worse, and he died on June 18th, 1929.

The fluid which was removed from the abdomen was subjected to chemical and bacteriological examination. To the naked eye it had the appearance of milk. Under the microscope it appeared as a fine fat emulsion, in which were observed a few red corpuscles and a few leucocytes containing fat globules. A differential cell count was performed, and the mononuclear leucocytes were found to be slightly in excess of the polymorphonuclear leucocytes. Stained smears failed to reveal acid-fast bacilli, and no growth resulted from inoculation of culture media. A guinea-pig was inoculated with the chylous fluid, and when killed at a later date was found to be normal.

The result of the chemical analysis was as follows:

Fat	...	...	...	...	...	11.42	per cent.
Proteins	...	...	...	...	...	11.57	"
Other organic matters	...	...	...	...	...	1.10	"
Mineral salts	...	...	...	...	...	7.89	"
Total solids	...	...	...	...	...	31.98	"

Of the mineral salts 58.22 per cent. consisted of sodium chloride.

Necropsy

The body appeared very emaciated. Post-mortem mottling was most marked over the right side of the chest. The abdomen was greatly distended, the linea alba being very marked, especially on the right side. When the body cavities were opened they were found to contain large quantities of milky fluid; 140 ounces were recovered from the abdominal cavity, 50 ounces from the left pleural cavity, and 36 ounces from the right pleural cavity. The fluid from the abdomen seemed to be of a more yellowish colour than that collected from the pleural cavities.

Heart.—The pericardium was found to contain about 6 c.cm. of straw-coloured fluid. The heart was small and in a state of contraction. The surface of the organ presented a whitish appearance, and the blood vessels were outlined by a white marginal line on each side. The heart muscle in its cut surface presented a finely streaked appearance due to the presence of delicate white lines radiating unevenly towards the surface. The muscle itself did not assume the normal reddish appearance, being dull and glistening and slightly brownish in colour.

Lungs.—Beyond a slight degree of hyperaemia the lungs appeared to be normal. The left lung was slightly adherent at its base, but no evidence of involvement of the bronchial glands was discovered.