directly may be concerned with cancer cases. This nucleus in one locality may be most conveniently based on the Public Health Department, but in another may best be executed through some hospital or voluntary association with which the Public Health Department co-operates. In respect of some kinds of cancer, particularly uterine, its nature may depend upon the extent to which the region is served by special departments or special hospitals at which radium treatment is given, and it results from this that local effort against cancer is now likely to be particularly active in connexion with those institutions to which supplies of radium have been allotted by the Radium Commission from the purchases of the Radium Trust. The kind of local action to which I refer in this summary way was lately reviewed rather fully in one of the memorandums (Cancer VII) of the Departmental Committee on Cancer at the Ministry of Health, the issue of which followed the transfer to the larger public health authorities of the many cases of cancer which were previously the responsibility of the Poor Law Institutions.

It has also been largely with the object of facilitating organized local action that the Departmental Committee has, over a series of years, recommended official circulars, and produced monographs on the causes and treatment of cancer of accessible sites. We owe to that Committee and its workers the determination of the average natural duration of the different varieties of cancer in different parts of the body, and various assessments of the extent to which appropriate operative treatment is being made use of, and facilities for early diagnosis are available. And its work has been specially useful when it has provided a reasoned judgement, obtained from home and from foreign sources, of the suitability of particular operative and radiological treatments, as judged by survival rates at defined intervals after those treatments. A city or region, therefore, which desires to consolidate its cancer work and settle a local cancer policy can, with the aid of these reports, start its work with some useful measures of the degree of achievement which can be reached by medical, surgical, and radiological skill.

The matters to be considered and co-ordinated at the local nucleus, which in general may be taken to be the Public Health Department, include the sufficiency of the local methods of securing that early and curable cases of cancer come to light at the right time and are treated in the right way. One is apt to forget how many different conditions are included under the name "cancer." Many of them, like cancer of the skin and lip, are so essentially curable that they only contribute a small proportion to cancer mortality. It is within the sphere of practical effort to bring cancer of the breast and cancer of the uterus into the same category. The local question is whether in fact these opportunities for cure are provided and used, and, if not, how the defect can be remedied through combined action of the local agencies available. There is again the large question of the sufficiency of palliative and nursing treatment in advanced cases of cancer, and, finally, the recording and following up of cases to which I have just referred. We can only wish well to those cities in which pioneer work is being done in these directions, not necessarily in the same way but always with the same object.

CONCLUSION

It seems, indeed, even from the random examples I have given, that in dealing with public health policies in regard to particular diseases we can hardly have enough local experiment. No one locality can attempt, or can be expected to attempt, to try out everything new that looks promising. But it is all to the good to find one particular area identifying itself, for example, with a carefully contrived effort to secure immunization against diphtheria,

another specially concerning itself with a policy to prevent heart disease, or a third with cancer. The same applies to many other forms of local effort, none the less valuable because it arises from circumstances that are fortuitous, like a local experiment for the communal treatment of chronic rheumatism and fibrositis, or the establishment of a foot clinic. Always the existence of such particular policies gives life and individuality to the local health work, and particularly when that work is not confined to the statutory public health authorities, but, as in the instances I have given, requires a combined action with which the whole medical profession of the place is identified. The position is much the same with the international work for health which is becoming so important. Uniformity is often desirable, and standardization is often essential to progress. One requires standards for statistics, terminology and for all kinds of purposes of comparison and research. They are essential also to facilitate a common understanding between nations in those instances where success must depend on united effort. But this is something quite different from the view that one should aim at conducting national health work on standardized international patterns or in pursuance of authoritative international doctrines. Room must be left for each community to develop its health policy on individualistic lines according to its own resources and the circumstances and genius of its own population. It is a healthy sign of progress in public health that this process can continue side by side with necessary consolidation and standardizing, and that communities, large and small, should still be ready to work out and apply new policies of their own or to reconsider and recast lines of effort so consecrated by tradition that they usually are not questioned. By so doing they help the world at large, certainly by success if it comes, but often hardly less usefully by learning the lessons of failure if they should be disappointed—which, after all, has been the essence of progress in true art and science from the beginning.

LATE RESULTS OF OPERATIVE TREATMENT IN TOXIC GOITRE

BY

CHARLES S. D. DON, M.D., M.R.C.P.LOND.

ASSISTANT PHYSICIAN, SALFORD ROYAL HOSPITAL

The beneficial results of thyroidectomy in cases of toxic goitre have led to a steadily increasing number of these patients being referred to a surgeon for this operation, while the present pre-operative treatment with iodine has reduced considerably the mortality rate. There are comparatively few follow-up studies, but in the papers published the percentage of good results claimed has varied so widely that the average reader is left with the impression that, assuming the writer to be unbiased, either the criteria for judging results differ or the results depend on the clinic at which the operations took place; the former is probably correct. It seems probable that the average case operated on in the United States is milder than in England; this fact is no doubt the main reason for the difference in the immediate operative mortality rate. The apparent higher incidence of the disease in America cannot be due to earlier recognition by means of the more frequent examination of the basal metabolism, as, in my experience of 1,200 basal metabolic rate estimations in the past three and a half years, this test when properly done prevents many neurotic patients from receiving x-ray treatment. The test has to be repeated several times in these cases.

FOLLOW-UP RESULTS OF VARIOUS OBSERVERS

Before proceeding to give the details of my own cases I wish to mention the follow-up results of operative

treatment of toxic goitres obtained by other observers, but in doing so I make no attempt to include all papers on this subject; a selection of statistics has been made for the sake of comparing them with my own results. Many of these follow-up studies are based on correspondence, others on personal observations; in some the basal metabolic rate has been estimated, but very often this test has not been done. In 1920 Judd¹ found that of 100 cases of Graves's disease, 55 per cent. were cured, while of 100 toxic adenomas, 83 per cent. were cured. After the introduction of iodine the results improved, and Elliott² found that of 72 cases in which the basal metabolic rate was done at the time of the follow-up, 63 were considered satisfactory; out of a total of 100 cases examined then, however, 46 showed signs of definite visceral injury due to the disease or to the operation. Dunhill³ states that of 185 patients about whom he received information, 167 were able to lead their normal lives, 4 died some time after operation, including 1 from high blood pressure. Walton⁴ claimed that 55 per cent. of his cases were cured and that a further 26 per cent. were improved. Morley⁵ assessed the results in 151 cases, regarding them as "very good" in 98 cases, and "good" in a further 37 cases. Smith, Clute, and Strieder⁶ consider that 92 of 100 cases examined were cured, while Clarke and Black⁷ report that of 64 cases in which basal metabolic rates were done at the follow-up, only 4 had a reading of over plus 20 per cent. Many of their cases were mild, 30 out of 62 having basal metabolic rates not exceeding plus 30 per cent. before operation. Thompson, Morris, and Thompson⁸ state that of 190 cases followed up, 19.5 per cent. were not cured, and in their opinion the toxic symptoms seen in this group had, in most cases, persisted since operation; in 2 patients the symptoms subsided after operation, but recurred later. Crile⁹ gives the number of his recurrences in cases of hyperthyroidism as 4.4 per cent.

A FOLLOW-UP OF FORTY-TWO CASES

In this contribution the details of 42 patients who were operated on either by Mr. G. Jefferson or by Mr. Garnett Wright are given, but before presenting details I wish to mention the method of examination.

A list of patients who had had operations for toxic goitre was compiled up to March 31st, 1930. These were circularized and asked to come into hospital for one night. A large number refused or did not reply. In many cases the information was volunteered that the patient was feeling quite well; all of these persons were excluded, but the 42 who accepted were admitted to the ward in the evening, a clinical examination, a basal metabolic rate estimation, an estimation of blood pressure, and an electrocardiographic examination being carried out on the following morning. The electrocardiographic results will form the subject of another paper, but at this point I may observe that an inverted T in Lead III was present in many of the follow-up cases. In some, myocardial damage present before operation was discovered to have cleared up completely at the follow-up; in others I had reason to believe that this had progressed since operation, in spite of an apparent complete relief of toxic manifestations as a result of operation. The other points to which particular attention was paid were the presence or absence of nervousness, tremor, exophthalmos, and the rate and rhythm of the pulse. Of these 42 patients, 38 were suffering from Graves's disease and 4 from toxic adenoma. In some cases, the basal metabolic rate was a little high compared with the pulse rate, but it was found that any untoward incident caused a surprising increase in the pulse rate (for example, the electrocardiographic proceedings), while the patient admitted that he was still nervous. A few patients had never had a basal

metabolic rate estimation done before; for this reason I made them breathe into the Douglas bag and accustom themselves to it on the night of their admission, the test being done next morning. The basal metabolic rates were done by the Douglas bag and Haldane gas analysis apparatus. The average interval between operation and the follow-up examination was a little over three years, the longest interval being eight years, the shortest one year. The majority of these patients were women, there being only five men.

RESULTS OF EXAMINATION

The details of the examination of these patients were as follows:

Exophthalmos.—When present, this is one of the most outstanding features of the disease, yet in some cases where I had not seen the patient previously I found difficulty in deciding the previous degree; in these cases I accepted the patient's statement on the amount of improvement. Before operation, exophthalmos was present in 29 of the 42 cases; in 18 of these it was well marked, in 11 it was slight, and in the remaining 13 there was not any unusual prominence of the eyes. In some cases it is possible that another observer might have disagreed with my classification, but it will suffice to give an impression of the result of operative procedures on this sign. At the time of the follow-up examination, of the 18 cases in which it had been well marked it was absent in 5, much improved in 11, and in 2 there was no change, but, in these, toxic symptoms were still present, the basal metabolic rate being plus 20 per cent. and plus 31 per cent. respectively. Of the 11 cases which had shown, prior to operation, exophthalmos in a milder degree, it had disappeared in 4, but still persisted in a milder degree in 7 cases. In one patient (Case 7) the exophthalmos, which was well marked before operation, increased during the year, following a subtotal thyroidectomy, and at the present time (three years later) the exophthalmos is approximately similar in degree to that before the operation. This patient's basal metabolism was normal three weeks after operation, but each of four basal metabolisms estimated since have been about plus 20 per cent. I have not seen a case in which exophthalmos which was absent before operation has made its appearance since. Zimmermann¹⁰ reports eight cases in which exophthalmos became more marked after a thyroidectomy had relieved the actual hyperthyroidism.

Nervousness and Tremor.—These were present to a greater or lesser degree in every case before operation, and were greatly improved in 39 of the 42 cases, though they were still present in 17 of these 39 patients. Of these 17 persons, 10 had basal metabolisms of or below plus 13 per cent., but in 7 it was still increased above plus 13 per cent. The remark made by Moschowitz¹¹ might be applied to those cases with normal metabolisms yet still somewhat nervous: "Thyroidectomy therefore removes only the hyperthyroidism and the increase of basal metabolism; the personality of the constitution remains, a potential, given enough stimulus, to flare up into the more intense forms of Graves's syndrome." In the 3 cases in which there was no improvement, the basal metabolisms were still plus 47 per cent., plus 42 per cent., and plus 59 per cent. respectively.

The Blood Pressure.—Unfortunately there are very few of the cases in which a record has been made of the blood pressure before operation. This is all the more regrettable, as Clarke and Black⁷ found that among their cases in which blood pressure readings were taken before operation and at the time of the follow-up, although the average systolic pressure had only risen 2 mm. Hg and the diastolic 9 mm. Hg, in certain cases the blood pressure had definitely increased. They mention that at the time

of discharge after operation 7 patients had a blood pressure of over 150 mm. Hg. while at the follow-up 15 cases had blood pressures of over 150 mm. Hg. Among the cases I investigated, the blood pressure was taken on 35 patients at the time of the follow-up; the average systolic pressure was 147 mm. Hg and the average diastolic 88.3 mm. Hg (the normal¹² for the average age, 36, is systolic 126 mm. Hg, diastolic 90 mm. Hg). If the patients are divided into two groups—namely, those above and those below 41 years of age—then the average systolic pressure for the group over 41 years is 153 mm. Hg and the diastolic 89 mm. Hg. The average age was 45, and on that basis it should be systolic 129 mm. Hg, diastolic 91 mm. Hg. For the group below 41 years the average systolic is 142 mm. Hg and the diastolic 88 mm. Hg; the average age was 28, and the normal reading for the age is systolic 122 mm. Hg, diastolic 87 mm. Hg. Among these 35 cases there are 14 patients with a systolic blood pressure of 150 mm. Hg or over, and of these 7 are below 41 years. It seems clear, therefore, that the frequency of the increased blood pressure cannot be attributed to the age of the patient. It was evident, also, that the height of the blood pressure had no relation to the number of years which had elapsed since thyroidectomy was performed. It did not seem to bear any constant relation to the persistence of the toxicity. As there are so few records of pre-operative blood pressures, no further deductions can be drawn.

The Weight.—The successful operative treatment of this disease usually results in an increase in weight, the amount depending somewhat on the loss of weight which occurred before operation. An accurate record of the weight before operation is only available in 27 cases; 23 have gained since their discharge from hospital, the amount varying from 2 lb. to 48 lb., with an average of 16 lb. Four patients have lost weight, varying from 2 to 5 lb.; 2 of these had normal metabolisms, and 2 have increased metabolic rates of plus 24 per cent. and plus 25 per cent. respectively.

The Basal Metabolic Rate.—The main purpose of this investigation was the estimation of the basal metabolic rate. I regard this as the most important single piece of evidence in assessing the condition of the patient; errors may occur, not through inaccuracy in technique but because these patients are emotional, and any marked variation in the result as compared with previous readings, or any discrepancy between the apparent clinical condition of the patient and this test, must be viewed with suspicion and checked by a further metabolic rate estimation.

Of the 42 cases described, a basal metabolic test was done before operation in 23 patients, in 17 of whom the rate was above plus 40 per cent. Rates below plus 30 per cent. were found in 4 patients, of whom 2 were adenoma cases; 1 was a recurrence and 1 (Case 38) was a case in which the rate was lower than was expected from the symptoms and signs. The remaining patients were operated on before I began to do the basal metabolic rates. These pre-operative basal metabolic rates are mentioned in order to emphasize the fact that most of the 42 cases were of the type with well-marked toxic signs, which, I believe, is operated on by surgeons in this country after medical treatment, and, in some cases, x-ray treatment, have proved inadequate. The immediate effect of the operation was to cause a fall in the rate in 21 of the 23 cases, while in 2 (Cases 3 and 38) there was a rise. In those cases in which the rate fell it did not necessarily approximate to normal at once, being at times below and at other times above normal limits. The actual details are not given, as the subject was discussed in an earlier contribution.¹³ The failure of the basal metabolic rate to become normal at once in many cases supports the view, re-stated by Pemberton,¹⁴ that the surgeon cannot judge how much

gland it is necessary to remove in order to bring the rate to normal, and that the operation breaks a link in the chain of etiological factors which permits readjustment to normal in the physiology of the gland. Of 7 patients who had had thyroidectomy and had later died of another disease, in 6 Pemberton found regressive changes in the histological structure of the gland, and this he regards as supporting his contention.

At the time of the follow-up a basal metabolism was done in all 42 cases; the longest time since operation was eight years, the shortest one year, with an average of just over three years. These figures are presented in the accompanying table, and it will be seen that in 13 cases the rate is still above plus 13 per cent., a high

The Basal Metabolic Rates at the Follow-up compared in some Cases with those taken before Operation

Case	B.M.R. at the Follow-up	B.M.R. before Operation	Operation
1	Plus 26 %	—	Right lobectomy
2	Plus 7 %	Plus 58 %	Subtotal thyroidectomy
3	Plus 24 %	Plus 23 %	Right lobectomy
4	Plus 5 %	Plus 61 %	Subtotal thyroidectomy
5	Minus 5 %	—	Right lobectomy
6	Plus 19 %	—	Ligation of arteries
7	Plus 20 %	Plus 55 %	Subtotal thyroidectomy
8	Plus 5 %	Plus 51 %	Ditto
9	0	—	Ligation of arteries
10	Plus 13 %	Plus 60 %	Subtotal thyroidectomy
11	Minus 27 %	Plus 65 %	Ditto
12	Plus 6 %	—	Ditto
13	0	Plus 46 %	Ditto
14	0	—	Ditto
15	Plus 4 %	Plus 41 %	Ditto
16	Plus 25 %	—	Ditto
17	Minus 22 %	Plus 52 %	Ditto
18	0	Plus 61 %	Ditto
19	Plus 22 %	—	Ditto
20	Minus 7 %	Plus 27 %	Ditto
21	Plus 2 %	—	Ditto
22	Plus 3 %	Plus 57 %	Ditto
23	Plus 47 %	—	Right lobectomy
24	Plus 31 %	Plus 24 %	Subtotal thyroidectomy
25	0	Plus 64 %	Ditto
26	Plus 12 %	—	Ditto
27	Plus 5 %	Plus 47 %	Ditto
28	Minus 12 %	—	Ditto
29	Plus 42 %	—	Right lobectomy
30	Plus 8 %	—	Subtotal thyroidectomy
31	Plus 6 %	Plus 67 %	Right lobectomy
32	Plus 2 %	—	Subtotal thyroidectomy
33	Minus 2 %	Plus 59 %	Ditto
34	Plus 59 %	—	Ditto
35	Plus 17 %	Plus 31 %	Left lobectomy
36	Plus 19 %	—	Right lobectomy
37	Plus 24 %	—	Left lobectomy
38	Plus 7 %	Plus 14 %	Subtotal thyroidectomy
39	0	Plus 33 %	Ditto
40	Minus 16 %	Plus 64 %	Ditto
41	Minus 9 %	Plus 45 %	Ditto
42	Minus 1 %	—	Right lobectomy

percentage, but in 7 of these the operation consisted of the removal of one lobe; in 1 only the arteries were tied, while in the other 5 cases the operation was a subtotal thyroidectomy, and the persistence of toxic symptoms is surprising. In 2 of the lobectomies (Cases 3 and 35) the operation consisted of the removal of an adenoma from one lobe, but the persistence of toxic symptoms suggests that smaller adenomas are still present in the remaining lobe. The patient in Case 6 (the ligation) had auricular fibrillation with marked oedema and ascites. As she did not show sufficient improvement after ligation to warrant a thyroidectomy she was sent home. She had signs of cardiac failure for twelve months, after which she gradually improved, and at the time of writing (five years after operation) she has no oedema, and is carrying on a fair share of her household duties. The patient in Case 34 stated that her operation effected a complete cure, but the excitement of escaping out of her burning house caused an immediate recurrence; there is no enlargement of the thyroid. The other 4 patients with toxic symptoms who had subtotal thyroidectomies were considerably improved by the operation; the present toxicity is a persistence of symptoms and not a recurrence. In the remaining 29 cases the basal metabolic rate is within normal limits in 25, while in 4 it is below minus 10 per cent. Clarke and Black⁷ found that 7 of their 64 patients had basal metabolic rates between minus 10 per cent. and minus 30 per cent. Smith, Clute, and Strieder⁸ mention that in 19 of their 100 cases there was a basal metabolic rate below minus 10 per cent., while in 15 of these there was definite evidence of myxoedema; they attribute this to the pre-operative use of iodine. Of my 4 patients with a basal metabolic rate below minus 10 per cent., 2 showed definite evidence of myxoedema—namely, Nos. 11 and 40. The patient in Case 40 had a rate of minus 16 per cent., this in spite of 6 grains of dry thyroid daily. Immediately after operation her rate was minus 18 per cent.

There were 4 patients who still had auricular fibrillation. In 2 of these the pulse rate was over 100; in the third it was 68, and in the fourth it was 72. One of them, patient No. 34, has developed fibrillation since her operation, which was a subtotal thyroidectomy, but as the basal metabolic rate is at present plus 59 per cent. it is to be expected.

CONCLUSIONS

The disappearance of exophthalmos does not of necessity follow the complete relief of hyperthyroidism, yet a diminution of toxicity is usually followed by an improvement in this sign, which may disappear completely in cases which we regard as cured. Exophthalmos is likely to persist in those cases in which the metabolism has not dropped to within normal limits. Nervousness is invariably improved by a successful operation, but the patient may continue to be nervous in spite of a normal metabolic rate.

The majority of patients gain weight after operation, yet there are occasional cases in which an otherwise successful result is not accompanied by any increase, and this is seen even in a patient in whom there was a considerable loss of weight before operation. Failure to gain weight is occasionally associated with glycosuria, which is sometimes seen in Graves's disease, and may become more severe after operation.

Relief of hyperthyroidism and a lowering of the metabolic rate to normal limits usually follow a subtotal thyroidectomy; there are cases in which the metabolic rate remains above normal limits in which no increase of thyroid tissue can be made out, and in which no focus of sepsis can be discovered in any part of the body; worry and too little leisure are probably the factors which prevent a good result in these cases. Crile⁹ attri-

butes these failures to the presence of septic foci in other parts of the body, or to worry and overwork.

The removal of one lobe is followed by improvement, and patients Nos. 31, 36, and 42 were types with exophthalmic goitre in whom relief of hyperthyroidism has followed this operation. A satisfactory result has also followed ligation in two cases, and it will be readily understood that the improvement which ultimately followed was the cause of the complete operation not being undertaken at a later date. Ligation alone is not regarded as a curative measure, and while it is of interest to note that a normal metabolic rate can follow this simple procedure, it is probable that the operation had little effect on the ultimate outcome.

If the clinical condition of the 42 patients, their ability to do work, and the rhythm and rate of the pulse are considered in conjunction with the basal metabolic rate findings, the result of operative treatment has been good in 28 cases, improvement in 9, while in 5 the operation has been of little value.

I am indebted to Mr. G. Jefferson and to Mr. Garnett Wright for allowing me to examine these patients, and to Dr. G. Murray for some very useful criticisms.

REFERENCES

- 1 Judd, E. S.: *Annals of Surgery*, 1920, lxxii, 145.
- 2 Elliott, C. A.: *Journ. Amer. Med. Assoc.*, 1927, lxxxix, 519.
- 3 Dunhill, T. P.: *Practitioner*, 1930, December, 672.
- 4 Walton, A. J.: *British Medical Journal*, 1928, i, 83.
- 5 Morley, J.: *Ibid.*, 1931, i, 450.
- 6 Smith, L. W., Clute, H. M., and Strieder, J. W.: *Surg., Gynecol. and Obstet.*, 1928, xvi, 325.
- 7 Clarke, N. E., and Black, I.: *Arch. Int. Med.*, 1930, xvi, 266.
- 8 Thompson, W. O., Morris, A. E., and Thompson, P. K.: *Ibid.*, 1930, xvi, 946.
- 9 Crile, G. W.: *Practitioner*, 1930, December, 661.
- 10 Zimmermann, L. M.: *Amer. Journ. Med. Sci.*, 1929, clxxviii, 92.
- 11 Moschowitz, E.: *Arch. Int. Med.*, 1930, xvi, 627.
- 12 Figures given by the American Institute of Medicine.
- 13 Don, C. S. D.: *British Medical Journal*, 1929, i, 1108.
- 14 Pemberton, J. de J.: *Journ. Amer. Med. Assoc.*, 1930, xciv, 1483.

TREATMENT OF CARCINOMA OF THE OESOPHAGUS*

BY

E. MUSGRAVE WOODMAN, F.R.C.S.

LECTURER ON DISEASES OF THE EAR AND THROAT, UNIVERSITY OF BIRMINGHAM; SURGEON EAR AND THROAT DEPARTMENT, GENERAL HOSPITAL, BIRMINGHAM

In this day of scientific progress we are faced with a disease, insidious in onset, slow in its development, relentless and remorseless in its course, and finally fatal in its issue. Against this disease we must admit that progress has been slow and almost unperceivable, and yet a great deal of work has been done, and many hopes have been raised only to be quickly dashed to the ground. With this admission we will first review briefly the principal difficulties, then proceed to a critical survey of the existing methods of treatment, and finally suggest certain points on which discussion would seem to be particularly valuable and indicate certain directions in which further research is urgently called for.

The principal difficulty lies in the anatomical formation of the oesophagus. Suspended at the entrance, where it is attached to the back of the cricoid cartilage, and fixed at the lower end, where it traverses the diaphragm, it remains completely mobile in the centre. This can be demonstrated by passing a tube into the oesophagus and viewing it on the x-ray screen, when the end of the tube can be freely moved from side to side.

In the cervical region the gullet lies in intimate contact with the trachea, into which perforation readily takes place. In the central zone the left bronchus is easily

* Read in opening a discussion in the Section of Oto-Rhino-Laryngology at the Annual Meeting of the British Medical Association, Eastbourne, 1931.