that by these movements the child is sucking in quantities of liquor amnii by which it will be drowned. Surely the child's mouth and nose are pressed tightly against the wall of the vagina, and not free in a bath of fluid, so that literally no liquor amnii can be inspired at this stage of delivery. It is a fallacy that the child can drown himself during extraction of the head, and therefore the inspiratory movements should not influence the accoucheur during his efforts to deliver the head.

The first point to try to be sure of is whether or not the child's head is flexed as it lies in the antero-posterior direction—that is, with its back facing anteriorly. Difficulties will depend more upon absence of flexion than size of head and pelvis; in fact, where the pelvis is normal incomplete flexion is the cause of nearly all difficulty. When, therefore, can we *suspect* that flexion will be imperfect? First, in all cases in which legs and arms have been extended, then in most cases in which the child's body has been pulled down by the contracting uterus, and, rarely, in some cases of flat pelvis. We can, however, *expect* complete flexion in all multigravida where legs and arms have been normally flexed, in all small or premature babies, and in nearly all cases in which the delivery of the child's body has been accompanied by firm pressure on the head through the fundus of the uterus. The last contingency implies the necessity in all cases of instructing the nurse to press firmly on the fundus during the delivery of the body.

THREE CLINICAL TYPES

In practice, therefore, we have three clinical types. The first is where a multipara has delivered herself of the legs, body, and arms normally and easily. Here we may assume the head will be flexed and give no trouble. The direction is to deliver by the simplest method, which is to grasp the feet in the right hand and lift them gently and gradually up to the mother's abdomen, while the left hand, or a nurse, presses firmly above the symphysis. No other manipulation, or grasp, of the child's body is necessary.

The second clinical type is the primigravida who has been slow and "difficult," possibly with some slight general contraction of the pelvis. Here the head *may* not be fully flexed, and, therefore, for these patients it is well first to pass a finger up to the chin and feel whether it is pressed well down on the sternum. If it is flexed, deliver the child by a grip which will allow some harmless traction—that is, by the so-called Prague manoeuvre. Take the feet in the right hand and place the first and second fingers over the child's shoulders. Traction is made on the shoulders, while the feet are carried to the mother's abdomen as the head is pulled down, while, again, the nurse presses firmly above the symphysis. Care must be taken to deliver the chin and face slowly from under the perineum; in fact, as soon as the mouth and nose are liberated there should be a pause for the sake of both the head and the perineum.

The third clinical type is that in which we can assume that the head is extended; for example, where we have already brought down arms and legs from the extended position. It is also likely to be extended if it does not descend well through the brim, and thus appears to be high up with stretching of the neck. Here the "jaw traction" method should be adopted. The right hand is passed into the pelvic cavity so that the index finger slips readily over the lower jaw into the mouth, the child's body lying astride on the right forearm. The first and second fingers of the left hand are placed over the shoulders. Before there is any traction made on the shoulders the head should be flexed by gentle traction on the lower jaw and simultaneous firm pressure on the fundus. Usually the head can be

made to flex by this manoeuvre, and then firm traction is made on the shoulders, which are, as the head descends, carried towards the mother's abdomen.

Some accoucheurs prefer to deliver all difficult cases by forceps, but of this I, personally, cannot speak from experience.

THE PROGRESS OF ANAESTHESIA*

BY

R. J. CLAUSEN, M.C., M.B., B.S.

It is no doubt fairly generally believed that very considerable progress has been made in recent years in the art or science of anaesthesia, and that great benefit has resulted therefrom to those undergoing surgical operations. New drugs, machines, and methods have been introduced from time to time with a greater or less flourish of trumpets, the echoes of which have occasionally reached the public through the daily press. It has in fact occurred to me on more than one occasion that a patient has remarked in the course of preliminary conversation how much these things have improved since the occasion of a previous operation, perhaps ten or twenty years ago, and what wonderful people we doctors are.

Such a state of mind is doubtless very beneficial to the patient, but the flattery though pleasing to us seems hardly justified if, as may well be the case, nothing more remarkable is about to be given than ether or chloroform on a mask, perhaps preceded by a hypodermic injection of 1/100 grain of atropine. However, there is no doubt that the resources at our command have considerably increased, and the object of this paper is to suggest that full advantage has not been taken of them, to inquire into the causes of this neglect, and, if possible, to suggest remedies for it.

It will be obvious that without a very exhaustive inquiry it is impossible to assess the state or progress of anaesthesia as a whole throughout the land with any degree of accuracy, and an incomplete inquiry may be as misleading as none at all or even more so. For operative surgery is carried out not only in the accessible city hospitals but also in numberless small hospitals, nursing homes, and private houses scattered over the country. An example of an incomplete inquiry was provided in a recent number of a periodical devoted to anaesthesia and analgesia, in which a writer made a rather ambitious attempt to survey world conditions. Such an article must be either very general or quite complete, but the author made the mistake of mentioning a few names of individuals; I was mentioned, for instance, as having the only large experience of ethylene in this country, but I am sure that the President of this Section, to name only one other, has a very much greater experience of this anaesthetic than I can claim.

For the purposes of this paper, therefore, I have not attempted any inquiry whatever, and it follows that any criticisms I shall make may be entirely unjustified. A good deal of information may, however, be gathered and inferences drawn from articles and correspondence in the journals, conversation with colleagues, the equipment of cottage hospitals and nursing homes, and so on. Further, to establish my credentials at any rate as regards London, and to avoid any suggestion of individual prejudice, I may mention that during the last ten years I have held resident or honorary anaesthetic appointments in nine hospitals, of which two have medical schools attached. It should also be understood that conditions in this country alone

* Read in opening a discussion in the Section of Anaesthetics at the Annual Meeting of the British Medical Association, Eastbourne, 1931.

are under consideration, and that no attempt is made to deal with anaesthesia abroad. It is probable that the most outstanding differences in the practice of other countries arise from the fact that in them the majority of operative work is carried out in large hospitals or clinics, and very little in private houses or nursing homes.

In the large teaching hospitals here, which are of course the most easily accessible for inquiring visitors, the standard is probably fairly high, but the adoption of intricate or experimental methods may be hampered by the necessity for teaching general principles and routine methods to students, who cannot be made fully conversant with all new developments. On the other hand, there is at times an unfortunate tendency for some new method or technique to be exclusively used in the hospital of its origin, with the result that students acquire an exaggerated belief in the benefits of this method and a lamentable lack of knowledge of others. As in other branches of practice, of course, the newly qualified man is only beginning to learn, and he may have to forget a good deal of what he saw and learnt as a student; but in the brief time that he then has he should be enabled to acquire a general foundation on which to develop a more specialized knowledge later, should the occasion arise.

Before proceeding to a more detailed criticism of our present methods it would be as well to point out that although the subject may appear to be of most interest to this Section, yet actually it concerns the surgeons more closely. For as Dr. Torrance Thomson recently pointed out in the *British Journal of Anaesthesia*, the surgeon must realize the importance of our work if progress is to be made, since without his interest and co-operation we can make but little advance, notably in the field of gas-oxygen anaesthesia, or any local methods.

SOME CRITICISMS AND SUGGESTIONS

Dealing, then, with our present methods, the following criticisms and suggestions are put forward, though naturally in the space of such a paper as this it is not possible to go into very great detail.

I

Insufficient individual attention is given to patients, particularly in hospital, but also in private practice, since operating theatre habits and customs are acquired in hospital by physicians, surgeons, and nurses.

The saying may be familiar that "routine anaesthesia is bad anaesthesia," and this may be correct, but nevertheless a good deal of routine is useful and necessary in our work, to enable it to be smoothly and easily done. Probably most anaesthetists adopt and perfect some standard technique, which tends to be applied to every patient, perhaps without great regard to their needs or desires. It is probably true that for operations of reasonable length—say, under an hour's duration—on healthy persons and where little shock is expected, the choice of anaesthetic makes little or no difference to the recovery of the patient and but little to the immediate discomfort of the operation. However, a large proportion of our cases do not come into this favourable category, and if the scope and advantages of new methods are not realized these less fortunate patients may suffer.

Besides the lack of adaptation of anaesthetic methods to individual cases there is often a striking lack of regard for the patient as an individual. Both failings are due to or encouraged by the system in large hospitals, where the anaesthetist is confronted with a long list of major operations of all descriptions, on patients he has never seen before and knows nothing of, while conditions are not improved by a general atmosphere of hurry and an impatient surgeon striding from room to room exclaiming

"Let's get on with it." Probably arising from these conditions there is sometimes a tendency to regard an anaesthetic as a standardized article which can be applied indiscriminately to any patient, and an anaesthetist may be asked to give his services, no other information being given him about the patient than the nature of the operation it is proposed to perform.

It is interesting to see to what extent the individual patient may be lost sight of in hospital; in the course of instructing students I have often asked, "Who is the most important person in the theatre?" but needless to say I have seldom had the correct reply. A friend with whom I was discussing this subject made the illuminating remark that little else was to be expected so long as the architects designed hospitals with a magnificent front door for the lay committee, staff, and students, whilst patients entered by an underground passage at the back. I do not suggest that a radical reform of hospital architecture will greatly benefit our work, but I cannot help feeling that there are many perhaps apparently trivial points in the treatment of patients which could be greatly improved.

For example, it is, I suppose, laid down somewhere in the regulations that the head must be low when anaesthesia is induced, and consequently a nurse will often take away a patient's pillow and leave him nearly flat on a trolley or table in a position of great discomfort at a time when every effort should be directed towards making him as easy and comfortable as possible. Or, in a hospital which I visit, though a sorbo mattress had been provided for the table it was very difficult to get it used, the reason being that it made the movement of the patient into the lithotomy position a little more awkward, so instead a thin folded blanket was used on the metal-topped table.

Just as every surgeon should undergo an operation or two, and every anaesthetist a few anaesthetics, for instructional purposes, so it would be of value in the training of theatre nurses and sisters if they were dressed, towelled, etc., as for operation and placed in the appropriate position on the table for an hour or so. The incidence of post-operative headache, backache, and even of chest complications might be thereby materially lessened. Another striking example of the way in which the individual can be overlooked, with results that can only be detrimental, is shown by enthusiastic residents and others, who often indulge in vivid discussions in the presence of the next patient while perhaps still wearing bloodstained garments; I have seen a house surgeon actually brandish a fresh prostate gland dripping with blood over the trolley on which a conscious patient was waiting, while he discussed its points with the surgeon.

While the patient may thus be lost sight of, a tendency sometimes appears for the theatre to become a stage on which the star performer displays his skill, supported by leading lady, minor characters and chorus; for some obscure reason it often seems to be expected that the anaesthetist should supply the comic relief, while in addition he is expected to be a sort of stage hand and property man, manipulating the lights, table, suction pump and so on.

In only one textbook on anaesthesia so far as I know is there a chapter dealing with the subject from the patient's view-point, and that book is not an English one. The matter, however, seems one of importance and deserving of attention in the teaching of anaesthesia.

II

The use of mask and drop-bottle methods with chloroform and ether is still too prevalent; the ritual act of dropping liquid on a mask is however firmly established, and alternatives may still be regarded with suspicion.

The mask and drop-bottle have advantages, of course—for example, simplicity, portability, ease of control and so on. Their use, however, should be restricted if possible to induction, and if chloroform or other anaesthesia is to be maintained for long periods it should be with vapour methods and some amount of re-breathing, preferably with oxygen, the value of which cannot be too highly stressed.

It is not uncommon for protests to be made against the growing complication of anaesthetic procedures, and for the claim to be advanced that the simplest methods are necessarily the best. This may be so in some cases, but actually the best method is that which ultimately gives the best results to the patient. Naturally, if a man arrives with an elaborate machine which takes a long time to erect, wastes everyone's time, and then gives a bad anaesthetic, he deserves some censure; for, besides his other faults, he may have brought a good method into disrepute. On the other hand, if he only spends his own time and his results are good, no surgeon has a right to quarrel with his methods, for we do not attempt to dictate what tools the operator shall use. There are those who support the plea for simplicity by using but few and simple instruments, but even these possibly forget the enormous complication of preparation, sterilization, etc., which precedes the actual entry of the patient.

Open ether, once claimed as the safest and best anaesthetic, has been described as a method invented by amateurs for amateurs; it still has many supporters, though it is probable that as most successfully practised it is far from "open." I can remember the time when the method was introduced into my hospital, which may have been a little backward in this respect, as the latest novelty from America. Its success was rapid and its vogue widespread, and closed methods which had been used with but few exceptions since the introduction of ether were largely abandoned for some years; for the open method first showed how ether could give complete relaxation with relative safety and without limitation of oxygen.

As a result of the work of Haldane, Henderson, and others, the value of re-breathing and the dangers of prolonged open etherization first came to be recognized in America, where a return to closed methods with re-breathing was initiated which later spread to this country. Open ether, however, will die hard, for apart from its actual advantages there is a frank and reassuring sound about its name which cannot fail to be attractive.

III

Local, regional, and spinal methods are not exploited as fully as they might be.

It is commonly said that local anaesthesia is widely used on the Continent because there are no good anaesthetists there, but the converse may well be true, that general narcosis is often unnecessarily used here because few people trouble to master the technique of other methods. It is also said that the people of this country are by their temperament unsuited to local methods, but this is probably in the main not true; at any rate, the behaviour of patients under spinal is usually good and helpful, though naturally the whole theatre method and organization must be altered when dealing with conscious patients.

It is a striking fact how many persons, particularly women, will say, "Oh, doctor, I don't mind the idea of the operation, but I do dread the anaesthetic." Of course, many of these might change their minds when actually confronted with an operation in the conscious state, but the benefits conferred by intelligent pre-medication must not be forgotten in these circumstances.

I have heard the methods of anaesthesia in a Continental clinic summarily dismissed as "a combination of local and straps," but I have often thought that an intel-

ligent use of suitable straps would serve to maintain the patient in some of the fantastic operating positions very much better than the makeshift of pillows and sandbags which we use, and would thereby facilitate the work of the anaesthetist.

Spinal analgesia is being fully dealt with in another paper, so I will not enter into any detail. It is a topic which is constantly cropping up, for the method is re-discovered, written up and more or less widely adopted every few years, being presumably forgotten in the intervals; the present, needless to say, is a "boom" period. While not advocating its wholesale use throughout the body, as do some, or its routine use for abdominal work, I believe it to be a very valuable aid in many difficult cases, and should be maintained in an effective state, so as to be available when required.

IV

The advantages of gas anaesthesia, and notably ethylene, are not adequately utilized.

Nitrous oxide and oxygen anaesthesia received an artificial impetus in the later stages of the war, having proved so useful in the repeated anaesthesias on shocked and debilitated patients then met with. Later, however, the method was largely discredited in this country owing to its indiscriminate application to unsuitable cases, with inadequate preparation and inefficient apparatus, while its necessary limitations were not realized by operators. I fully agree with the authors of a recent textbook on anaesthesia, who say that "the apparatus available in England has tended to hinder the effective adoption of the method in this country."

It is unfortunate that the recent revival of gas-oxygen here has been attended by, if not actually due to, the use of imported machines, but there can be no doubt that these show considerable advantages, except perhaps in price. A similar lack of enterprise is shown by our engineering firms with regard to electrical suction and pressure pumps for surgical and anaesthetic use, for most of those sold by the instrument dealers are of foreign make.

As regards gas apparatus, there are those who claim that they can give a good anaesthetic with no more than a cylinder of each gas, connected to a bag and mask, or who criticize certain machines on the ground of their complication of dials, gauges, etc., which distract attention from the one essential—namely, the patient. However elaborate the machinery used the condition of the patient must remain the ultimate indicator and control, but the more refined the apparatus used—that is, the more accurately it is capable of giving and maintaining any desired mixture—the better will it serve both novice and expert. In this connexion, I have not yet heard an anaesthetist, or a surgeon, protest at the complication of his motor car, or complain that the number of dials and "gadgets" on the dash interfere with his attention to driving.

Wherever possible, in both major and minor surgery, gas-oxygen should be preferred to ether alone, though in many cases small amounts of the latter may be used in addition, while, where circumstances do not prevent it, ethylene should often replace nitrous oxide. In midwifery particularly, gas-oxygen is undoubtedly the ideal anaesthetic, and should be much more widely used. There are of course gas enthusiasts who will not tolerate even the smallest amount of ether, just as there are local exponents who have no objection to injecting unlimited drugs but will not hear of giving any by inhalation. Personally I see no reason why the fullest use should not be made of combined methods, which have many advantages.

Ethylene is usually condemned on the ground of explosion risk, but in view of the many risks we let pass unnoticed this shows a curious lack of proportion. For example, there

is now an enormous death rate from the entirely preventable disease of "motoring," but I believe I am right in saying that neither this Association nor the profession as a whole has taken any steps to deal with it. Actually I believe the probability of explosion with ethylene is little if any greater than with gas-oxygen-ether, or oxygen-ether, and at present fires, explosions and fatalities are by no means unknown in our theatres; but in certain circumstances the resulting explosion with ethylene would certainly be more severe and deadly.

The widest field for gas anaesthesia has been in the past its use for dental extractions in the chair, and though this is now smaller owing to the increase of local and regional methods, yet there are many cases in which it will be preferred. In patients able to co-operate and in the absence of nasal obstruction or gross deformity it should always be given by the nasal route and the mouth properly packed off before extraction is begun. The advantages of this method—namely, comfort for the patient, speed of induction, safety, absence of cyanosis, and prolongation of anaesthesia if required—are so manifest that it is surprising that other methods are still tolerated and that full face mask induction is still taught as the routine.

For long oral operations, as for other head operations, some form of intubation is required, preferably using gas-oxygen, at any rate for maintenance, but it is to be hoped that the vogue for the endotracheal insufflation of gas mixtures in all cases has now passed, and that this is no longer regarded as providing a short cut to good anaesthesia. Gas-oxygen, however, depends for its success on free breathing and exclusion of air, and though in general surgery these may as a rule be obtained with the ordinary mask, yet there may be some occasions when intubation of the larynx may be desirable to ensure these conditions.

V

The advantages, or rather the necessity, of adequate pre-medication are not fully realized.

It often seems to be thought that all that is possible in this respect has been done when 1/100 grain of atropine has been given, and this routine is often hard to break away from. There are, I believe, two reasons for giving atropine before anaesthesia: the first theoretical, because it prevents vagus inhibition of the heart; the second practical, because it dries up the mucous and salivary secretions of the mouth respiratory tract. I am not in a position to discuss the former, and suspect that the latter is the only one of benefit. The same result, drying of the mouth, is obtained when morphia alone is given, though possibly not to the same extent, and it is probable that atropine could often be omitted with advantage; it has no sedative effect but on the contrary a stimulant one, and may facilitate the onset of shock instead of delaying it as does morphia.

Morphia or its equivalent is essential if good results are to be obtained from gas-oxygen, and to get a good effect it may be given in a divided dose. Further, in long operations an additional injection may be necessary, while where it is known that it will be needed after operation it is often useful to give it before the end in order that it may have taken effect before the gas anaesthesia is over.

In a hospital where both are given, there is a striking contrast between the placid and quiet morphia patients and those who have had atropine only, the latter being bright-eyed and apprehensive. I believe that no patient should come to general or local anaesthesia for any serious operation without preliminary sedatives of the morphia type or an adequate dose of one of the "basal anaesthetics" unless there are real contraindications, of which, however, there can be but few. There is, however, a strong prejudice against preliminary morphia, dating probably from

the days when induction with open chloroform was the rule. In this case it is undesirable, for the combined effect so depresses respiration that progress becomes difficult or impossible, though probably this could be overcome by the use of carbon dioxide.

VI

The claim has been made that by the use of carbon dioxide to increase lung ventilation after anaesthesia the respiratory complications which have been too common in the past can with certainty be entirely eliminated. Likewise it is stated by some that by the routine use of intravenous glucose vomiting after anaesthesia, another of the bugbears of the past, has been abolished.

Though both claims may appear somewhat exaggerated they are doubtless well founded, and the more extensive use of both treatments would appear desirable. Carbon dioxide particularly might be more widely employed, since apparatus is available which renders its administration in suitable mixtures automatic, and which could safely be used in the wards by ordinary nurses.

VII

Perhaps the most serious failing in our anaesthetic procedures is the fact that in spite of the frequent, or rather continuous, discussions as to the merits of various methods, no real attempt has ever been made so far as I know to investigate this problem in the only place and way in which it could be effectively done—that is, by observation of a large number of cases operated on with different anaesthetics but under otherwise similar conditions.

A striking example of this neglect was provided some few years ago when workers at a large teaching hospital, in an attempt to improve on ether and to reduce its after-effects, introduced an artificially impure ether as a new anaesthetic. Though the conditions were ideal for a proper investigation, since large numbers of operations were done by the same men under conditions identical but for the anaesthetic used, yet no attempt was made to carry one out, and the numbers of cases reported by the partisans of rival methods were too small to be convincing. Eventually the manufacture of the product ceased, and there may be many here who have never heard of it.

Further, more recent examples might be quoted—such as spinal analgesia with percaine, which has been accused of causing an undue incidence of severe headache as compared with other agents; or the various barbituric acid derivatives. To base a comparison between three of these drugs upon an experience of 262 cases with one and of 23 and 15 with the others, seems to show a certain lack of sense of proportion to which medical men would sometimes appear particularly liable, unless indeed the results with the smaller numbers were uniformly and convincingly disastrous. I must plead guilty to the same crime myself, for I drew some conclusions as to the value of ethylene without control records of nitrous oxide cases. This series was begun, but was given up owing to the difficulty of collecting enough material while working single-handed in various scattered places.

Inhalation anaesthesia, however, is largely a personal matter and is more of an art than a science, and whatever method is adopted by a skilled administrator and used consistently, the results will be apparently the same in most cases; hence the wide diversity of agents, apparatus, and methods. This apparent similarity might be dispelled were adequate data available for the various methods, and in any case it does not apply to the large category of poor risks, in which the choice of anaesthetic may make a large difference in prognosis.

If what are sometimes regarded as special methods, such as gas-oxygen or spinal, are reserved for desperate cases,

not only is insufficient practice obtained with them to obtain the best results but they may come into disrepute owing to the high mortality apparently resulting.

REASONS FOR SHORTCOMINGS

Such then are, in brief, the shortcomings, real or imagined, of our anaesthetic procedure, but before considering remedies it is necessary to investigate the causes.

These fall into two main categories, traditional and economic, which affect both institutions and individuals. The force of habit, custom, or tradition is fairly strong in most of us, and we probably find it easier to go on doing things in the same way when a good procedure is established. Further, there may be strong national traditions, as, for example, the use of chloroform in Scotland, which we are told was until recently almost universal; or local ones, as a result of which one method may be taught and used almost to the exclusion of all others. Hospital routine or tradition may be reinforced by the conservatism of theatre sisters and nurses, which though it may appear unimportant, can yet provide an effective opposition or a damaging passive resistance. Surgeons, again, who have operated with ease and success for a number of years are naturally prone to look askance at new methods, which may not give the same ease of access as the accustomed one, and may at first take longer in induction. The tradition of haste and hurry which still overshadows some of our theatres is in part a relic of the old pre-anaesthetic days when speed was essential, and in part the result of present economic conditions.

Hospital finances are not as a rule too secure, and committees and secretaries naturally hesitate to order new and possibly expensive machines which may not show any apparent benefit and may be discarded after a few years with the change of fashions or of medical staff. The latter, again, are not likely to be unanimous in their needs, and this may lead to unnecessary reduplication of apparatus.

As we know, the financial position of the individual anaesthetist is in most cases more precarious even than that of the hospitals, if such a thing is conceivable, and the same considerations as to apparatus apply even more strongly to him. His troubles are complicated by the fact that his private work, from which his living is derived, is carried out in nursing homes scattered over a possibly wide area and quite devoid of equipment, even perhaps without a lift. Obviously, such conditions tend to portability and simplicity or crudity of apparatus, and do not favour gas-oxygen.

Hospital practice, which may form the great bulk of his work, is almost entirely unremunerative, though the extension of the system of paying beds may alter this. Though many hospitals when advertising for anaesthetists insist that these should be specialists and not engaged in general practice, yet the possession of such an appointment does not guarantee any financial return whatever. It is therefore hardly surprising if anaesthetists limit their hospital attendances to their specified operation days, and show little inclination to embark upon extensive researches, while on these days conditions seldom favour new or experimental work.

REMEDIES PROPOSED

If progress is to be achieved it must be through the recognition by all concerned in the work of operative surgery, of the importance of anaesthesia as a subject, and of the fact that its technique is not necessarily perfect and stabilized. The status of our subject must be maintained and improved, and in at least one respect—that is, the time factor—it must take precedence in the theatre; for whatever anaesthetic procedure is adopted it must be completed before the operative procedure is begun. In another

respect there is at times room for improvement, and that is in the standard of asepsis, which must be as rigid as possible where spinal, intravenous or other injections are made.

In teaching hospitals at least some attempt should be made by classification and investigation of records to evaluate the advantages and assess the risks of the various methods. This could well be done in the Surgical Units, where conditions should be more favourable, and where an additional anaesthetic assistant might be appointed.

The position of the amaesthetist as a specialist must be made secure and his services fully utilized in both "normal" and difficult cases, while the question of remuneration for hospital services needs to be considered.

Perhaps most important of all, the demon of hurry and hustle needs to be finally exorcized, and the tyranny of long lists broken.

INDICATIONS FOR PHRENIC AVULSION IN BRONCHIECTASIS

BY

W. G. OAKLEY, M.B., M.R.C.P.

Unilateral diaphragmatic paralysis by interruption of the corresponding phrenic nerve is the most recent of the principal methods of securing, by therapeutic collapse, that combination of rest and compression which plays so large a part in the successful treatment of chronic pulmonary diseases. With this object it may be used alone, or in conjunction with such methods as artificial pneumothorax, multiple intercostal neurectomy, and thoracoplasty.

THE RADICAL OPERATION

The operative procedure now generally employed consists of avulsion or exaeresis of the phrenic nerve through a small incision above the clavicle, for a distance sufficient to ensure breaking its connexion with the supra-pleural plexus and all accessory fibres. Although usually successful, this method has been known to fail in the production of a permanent paralysis, even after as much as seven inches of nerve have been avulsed.

Partial Operations

In addition to these permanent methods a number of operations have been devised by which a temporary paralysis may be effected in order to permit of healing of a tuberculous lesion and of a subsequent return of function of the diaphragm. These methods include crushing, freezing, alcohol injection, and section with immediate suture of the nerve, and are generally considered less satisfactory than the permanent operations. In bilateral basal bronchiectasis, however, the crushing operation has been successfully combined with phrenic avulsion in America by Matson in the following way. The phrenic nerve is crushed on the worse side, and the effect of a temporary paralysis on cough and sputum observed. If satisfactory improvement results, then the opposite nerve is crushed or avulsed, and the paralysis is later made permanent.

Complications of Radical Operation

The complications of the radical operation are more formidable in theory than in practice, and fall into three main groups: (a) those due to failure in technique; (b) those due to adhesions in the course of the nerve; and (c) reflex disturbances.

In the first group haemorrhage constitutes the most frequent and serious of all complications, and results from injury either to the pericardio-phrenic artery (which accompanies the nerve and venae comites in the same