place into the other lung, or the lesion may reach the surface, causing a pleurisy and adhesion, which will seriously interfere with a subsequent artificial pneumothorax.

In the diagnosis of early phthisis, it is far better to have a negative sputum and a negative skiagram than a missed case. Radiology plays an important part in the diagnosis of the initial infiltration, but it is not the deciding factor, for the skiagram of this lesion may bear a close resemblance to a small pneumonic area, a lung abscess, a new growth, or a gumma. It is only by taking into account the clinical history, the physical signs or the absence of physical signs, the pathological examination of the sputum, and, sometimes, of the blood for leucocytosis or Wassermann reaction, that the x-ray picture can be correctly interpreted. The pathologist, the radiologist, and the physician, when united, are becoming more and more powerful in the field of diagnosis of pulmonary disease. It is to be remembered that this new teaching was established by a physician, working hand-in-hand with his radiological and pathological departments.

TREATMENT

There is no doubt that a certain number of these initial infiltrations will heal and become fibrotic if the patient is given general sanatorium treatment only; on the other hand, the majority of them tend to spread if so treated, and the difficulty is that in a given case it is not possible to say whether the patient will do well on general treatment only. I have seen these cases treated by rest alone;

in the majority this line of treatment has been followed by a spread of the disease, and I have wished that artificial pneumothorax had been induced at the start. This is the best form of therapy, and the longer it is delayed the greater is the possibility of a spread taking place which may jeopardize the future artificial pneumothorax. At the same time it must be admitted that artificial pneumothorax does not arrest every case, particularly if cavitation has already occurred, but the sooner it is induced the greater is the chance of success. Because of the success of early artificial pneumothorax it is imperative to diagnose intact initial infiltration before cavitation or spread has taken place.

In my previous publication on this subject I translated Redecher's "Frühinfiltrate" as "early infiltration." Zimmerli and Simmonds,⁵ in a recent paper, suggest that "initial infiltration" is a better term, because the infiltration is the initial lesion of phthisis rather than the earliest manifestation of pulmonary tuberculosis in its broadest sense, which must include such lesions as Ghon's primary focus and complex. In this paper I have adopted the term "initial infiltration."

The two skiagrams reproduced (Figs. 1 and 2, Special Plate) show the typical radiological appearance of the initial infiltration. My thanks are due to Dr. V. Sparkes for the skiagrams.

REFERENCES

- ¹ Kayer-Petersen, T. E.: *Beitr. z. Klin. d. Tuberk.*, 1928, lxx, 132.
- ² Assmann, H.: *Ibid.*, 1924, lx, 527.
- ³ Lydtin, K.: *Zeit. f. Tuberk.*, 1926, xlv, 273.
- ⁴ Redecher, F., and Walter, O.: *Entstehung und Entwicklung der Lungenschwindsucht des Erwachsenen*, Leipzig, 1929.
- ⁵ Zimmerli, E., and Simmonds, F. A. H.: *Tubercle*, 1930, March-May.

A CASE OF MULTIPLE OSTEochondROMATA WITH OBSTRUCTED LABOUR

BY

E. J. BLACKABY, M.R.C.S., D.T.M. AND H.

DISTRICT MEDICAL OFFICER, NICOSIA, CYPRUS

(With Special Plate)

Bony tumours of the pelvis obstructing labour are rare, and it is uncommon to find multiple affections of this structure in the condition known as "multiple osteochondromata." The case described below is a combination of these two circumstances.

The patient was a woman of 28, well developed though of short stature (5 feet). She was married in 1928, and in the following year had a five months' miscarriage. On August 23rd, 1930, she was admitted to hospital in labour, which had been in progress for over twenty-four hours. She said that there had been slight loss of blood since the pains began, and that the membranes had ruptured. On admission her pains were frequent and strong.

On abdominal examination the lie was longitudinal, with the head above the pelvic brim. There was a definite overlap of the foetal occiput above the symphysis pubis, and the head was very freely movable between the pains. It was quite impossible to press the head down into the pelvis and to make it engage. The pubic bones seemed abnormal, but no definite opinion could be arrived at owing to the overlapping of the head. The foetal heart sounds could not be heard.

When the vagina was examined the os was fully dilated, and the presenting part was high and difficult to feel. No abnormality could be made out in the pelvic cavity. The pelvic measurements were: interspinous 9½ inches, intercrystal 11 inches, and external conjugate 7¾ inches. After the vaginal examination was completed, a large bony tumour of irregular outline was noticed on the inner side of the right femur at its lower end. It felt like a large exostosis, and, according to the history, had been present since early childhood. A further examination of the bones showed similar, but

smaller, masses on the opposite femur, the head and lower end of the right tibia, and the head of the left fibula. It seemed probable, therefore, that the abnormality felt on the pubic bones, preventing descent of the head, was a bony growth of similar nature. In spite of strong pains there was no advance of the foetal head, and a Caesarean section was considered the best method of delivery. This was accordingly performed, and the woman was delivered of a living and healthy male child, which weighed nine pounds. There followed an uncomplicated puerperium, and the patient was discharged from hospital on September 6th.

DESCRIPTION OF RADIOGRAMS

Radiograms taken afterwards are shown on the Special Plate. The pelvis (Fig. 1), besides showing growths of bone on its outer surface, exhibits a tumour of the superior ramus of each pubic bone on either side of the symphysis, narrowing the pelvic inlet and rendering it somewhat asymmetrical. This photograph also shows involvement of the upper ends of both femora by the disease. The right knee region (Fig. 2) shows multiple masses on the lower end of the femur and on the upper ends of the tibia and fibula. The left knee region presented similar but less extensive changes. These leg conditions caused no disability in walking. Screening of the upper limbs revealed osteochondromatous growths in the following situations: (1) upper end of the humerus on both sides; (2) upper end of the left radius; (3) distal end of the fifth metacarpal on the left hand.

TREATMENT

Treatment for so widespread a condition seemed impracticable, especially as it is said that these tumours do not grow after adult life is reached, but the patient was advised that a future pregnancy would necessitate another Caesarean delivery. The fact that she lives within easy distance of hospital influenced one against any attempt at sterilization by ligaturing and dividing the tubes at the time of operation.

COMMENTARY

This type of bone tumour starts as an enchondroma,¹ which ossifies later, near the extremities of the long bones of the limbs. According to Thompson and Miles,² it occurs neither in membrane bones nor in bones entirely formed from cartilage; this is borne out in this case. These authors mention involvement of the iliac crest, and Chaumet³ also refers to a case in which a mass had formed on the anterior superior iliac spine, but no mention is made by them of the os pubis as a site. Roughly, as has been observed in other cases, there was a tendency to symmetry of the outgrowths in this particular patient. Heredity⁴ is stated to play a part in the production of the condition. This patient's parents died "many years ago," and no cause of death is known. She has two brothers and a married sister, who is childless, but it was impossible to get them to come for examination; however, as far as can be ascertained, they have no deformity. The child was examined in November, 1930, and showed no signs of juxta-articular tumours.

My thanks are due to Dr. Coureas of Nicosia for taking the radiograms which are published as illustrations of this case.

REFERENCES

- ¹ Tubby: *Deformities, including Diseases of Bones and Joints*, Section IX, Chapter 2.
- ² Thompson and Miles: *Manual of Surgery*, p. 483.
- ³ Chaumet: *Traité de Radiodiagnostic*, p. 40.
- ⁴ Choyce: *System of Surgery*, vol. iii, p. 777.

THE HEALTH OF THE WORKER IN CHROMIUM PLATING

BY

HUBERT B. TRUMPER, B.A., B.CHIR.CANTAB.

MEDICAL OFFICER TO THE ASSOCIATION OF BRITISH
CHROMIUM DEPOSITORS

Workers in this industry suffer disablement, usually temporary, from one or more of the following effects of exposure to chromic acid or its salts: (1) dermatitis; (2) ulceration of the hands, arms, or feet; (3) ulceration of the cartilaginous part of the septum of the nose; (4) vomiting after food.

The process of chromium plating consists in wiring the articles on to a frame ready for the plating vat, the actual plating, which lasts anything up to fifteen minutes, the unwiring, swilling, and polishing of the plated articles. A relatively large current is used in the plating tank, from 500 to 1,000 amperes and 4 to 10 volts. The solution contains 50 per cent. of chromic acid, and during the working of the tank reddish-brown fumes rise up into the air, to be sucked off by exhaust fans. Analysis reveals that these fumes contain upwards of 60 per cent. chromic acid, forced up in the form of a spray by the evolution of hydrogen at the cathode. The polishing of the work is done on a calico mop impregnated with either white or green "compo." The abrasive in the white compo is fused alumina; in the green it is a green chrome oxide (Cr_2O_3). Both these are held as a bond by high-class stearine. Here also the dust from the polishing is sucked off by exhaust fans. Employees actually working on the plating tanks are, as might be supposed, most liable to trouble, but any or all of the conditions mentioned above may occur at any stage of the plating process, or even in persons working in the same shop.

There are at least four main factors in the production of these conditions: first, a septic process in the body of the worker; secondly, inefficient exhaust apparatus;

thirdly, a moisture-laden atmosphere, a sweat-laden skin, or both; and, finally, carelessness on the part of the operator.

PREVENTION OF DISABLEMENT

Persons to be employed in this industry should be medically examined before commencing on this process to prevent those with such obvious septic foci as foul teeth or tonsils from undertaking work which will inevitably lead to disablement. Sepsis of the nose, mouth, and pharynx is the commonest predisposing factor. The exhaust system of the tank should be sufficiently powerful to draw the smoke from the cigarette at the lips of a man standing by the tank. Scrupulous cleanliness on the part of the operator is, happily, as unnecessary as it is difficult to obtain. It is important, however, that wash basins, with a good supply of really dry towels, should be provided. In addition, the skin must be protected with some form of grease. This is of great assistance in preserving the health of the horn cells. The most satisfactory is a modified Lassar's paste with a preponderant base of pure vaseline. This should be smeared on the hands and arms before work is commenced, and a small piece inserted into each nostril, which quickly spreads over the exposed mucosa. In one large plant the operators use a specially prepared soap free of alkali and containing a proportion of free lanoline. This extra lanoline is left as a thin deposit on the skin after the hands and arms have been washed. Rubber gauntlets are compulsory, and are worn where contamination with the solution is inevitable. If they are worn continuously the additional perspiration produced more than counteracts the beneficial effect of the protection. Respirators are not of proved value. In a large plant the floor quickly becomes awash with drippings from the tanks and swills. This is a potent factor in the causation of trouble, and can be obviated by covering the floor with fresh sawdust each morning and sweeping clear at the conclusion of the day's work. Carelessness on the part of the operator can be avoided only by intelligent supervision.

TREATMENT

Dermatitis.—The usual positions for this form of dermatitis are the hands, arms, face, and occasionally the chest. The onset is sudden, but it is very unusual for an attack to occur until the operator has been in contact with the plating plant for at least six months. A severe case presents a horrifying appearance. The face is intensely red, and swollen out of all recognition. I have seen four cases in which oedema was so severe as to cause temporary blindness. Where the arms are affected they, too, are red and swollen, and the affected parts itch intensely, or may even be excruciatingly painful. The conjunctivae are injected, but the mucosa of the mouth seems to be unaffected, save that some patients complain of intense tingling of the tongue. At this stage the patient should be put to bed and the affected parts covered with lint soaked in double strength calamine lotion. The dressing should be changed frequently. Morphine is sometimes necessary to allay the irritation. Twenty-four hours later the oedema has subsided; and the affected skin is dusky in colour, with clusters of brick-red spots separated by crusted effusion from the contracting corium. The skin should now be scrubbed with gauze soaked in spirit, carefully dried, and swathed in lint plentifully spread with Lassar's paste. This procedure should be carried out twice daily. As soon as all irritation has ceased, usually in two days, the patient should resume work in the plating shop, being dressed by the first-aid official night and morning. The majority are clear of the dermatitis in a week. If patients remain

H. V. MORLOCK: THE DIAGNOSIS OF EARLY PHTHISIS

FIG. 1.—Female aged 25; the only symptom was cough of seven weeks' duration. Sputum T.B. positive. The skiagram shows a typical initial infiltration in the right mid field.

FIG. 2.—Female aged 28. Skiagram taken two months after onset of cough following a short febrile attack; shows a typical initial infiltration in the right infraclavicular region.

E. J. BLACKABY: MULTIPLE OSTEOCHONDROMATA WITH OBSTRUCTED LABOUR

FIG. 1.—Radiogram of the pelvis showing growths of bone on its outer surface and a tumour of the superior ramus on either side of the symphysis.

FIG. 2.—Right knee. Multiple bony masses on the lower end of femur and on the upper ends of the tibia and fibula.