twice done tendon-lengthening operations for this type of deformity. Deformity resulting from fibrosis of the ulcer bed generally produces talipes equinus and calcaneus; (d) mental changes, chiefly melancholia, neurasthenia, and hebeteude.

COMMENTARY

This is a very formidable list of complications to which rarities can still be added. They all result from loss of balance between arterial and venous systems of the leg. This balance can be restored in two ways—by lying down and by reinforcing the skin with an elastic support. In the cases due to varicose veins a great improvement can be made by obliterating, and in bad cases also ligating, the varices, so that 75 per cent. of ulcer cases can be cured, and remain cured, without any support but the elasticity of the patient's own skin.

Ulcers of the leg will always be with us till we find a way of preventing femoral thrombosis after operations, fractures, and childbirth; but they will diminish in number because of the simplicity of the injection treatment for early varicose veins. By this method of treatment we have a certain way of curing all gravitational ulcers. The treatment of small ulcers, up to 4 square inches, is a very simple matter, quickly accomplished, with a minimum of trouble. These ulcers can be cured by other methods, I admit, but generally at the expense of much trouble and time to patient and surgeon. The larger ulcers are generally acknowledged to be incurable, but they can be readily cleared up by this method, even when of the gigantic dimensions of 180 square inches ($1\frac{1}{4}$ square feet!), the largest ulcer I have yet treated since I have kept statistics. I have now 525 cases of cured ulcer of a total area of a little over 25 square feet.

One of the most wonderful things in nature is surely to see the ulcer which started a quarter to half a century ago (my longest case is fifty-four years) begin to throw up its granulations, flatten out its margins, pull in its edges, and epithelialize in the way described so well by the "Father of Surgery"—John Hunter—many years ago.

A CASE OF ACHALASIA OF THE CARDIA

BY

H. B. BUTLER, F.R.C.S.ED.

SURGEON, ROYAL SURREY COUNTY HOSPITAL, GUILDFORD

(With Special Plate)

The following is the history of a male, B. K., aged 35, who was sent to the x-ray department of the Royal Surrey County Hospital with a probable diagnosis of pyloric obstruction.

There was a history of difficulty in "swallowing" for the past two years, much worse during the last four months. He had lost about a stone during the latter period, but though rather emaciated his physical condition was fairly good. He was of the opinion that "all" his food returned after having been retained for some hours; it gradually worked up, but he did not actually vomit. He got a feeling of tightness and great discomfort in the chest until the food was regurgitated. He had been given a barium meal at home, six hours before being seen at hospital, and when he was examined on the screen it was seen that this meal was still in the greatly distended oesophagus, none being seen in the stomach or elsewhere. The obstruction was at the cardia.

The oesophagus was evacuated, and then a soft mercury tube was passed into the stomach without any obstruction being encountered. Immediately after the passage of the tube he was given half a pint of beaten egg and milk, which he swallowed with ease and evident pleasure, saying that it was

the first food he had felt entering his stomach for months. He was admitted to hospital to be taught to pass the tube himself, which he soon accomplished, passing it before every feed; and he was discharged in about a week, able to take all soft food and doing well.

On June 14th, 1929, he came again, and complained of difficulty in swallowing, but as the tube was seen to pass easily into his stomach he was told to persevere (see Plate, Fig. 1). On July 27th he was readmitted, his condition then being nearly as bad as when first seen. Fig. 2 shows an immediate meal distending the oesophagus (July 27th). Fig. 3, taken on July 31st, shows a mercury tube, which he was able himself to pass into his stomach. Fig. 4 shows a bread-and-milk barium meal, the consistency of cream, taken immediately after the successful passage of the tube, and demonstrates the fact that the achalasia was immediately present, even directly after dilatation with a large-sized tube.

No obvious cause for the condition could be discovered, and the patient begged for "an operation, and be done with it." I had had no experience of dilatation with an inflatable bag, and felt that operation offered the best chance of success.

Operation was performed on August 1st. No cancer or ulcer of the stomach or duodenum was found, nor any lesion of the gall-bladder. A transverse incision in the anterior wall allowed three fingers to be introduced into the stomach, but to my surprise I had great difficulty in locating the cardia. (In this connexion I was afterwards interested to read in a paper published by the Mayo Clinic: "The silk thread is just as valuable a guide to manual dilatation from below as it is to dilatation from above, by means of the hydrostatic dilator.") When the cardia was found, one finger was introduced, then a second, and then a third, before there was any sensation that further dilatation would be dangerous. There was no palpable hypertrophy. The stomach and the abdominal wound were then closed. The patient made an uninterrupted recovery, and left the hospital within three weeks.

As I had no report about this patient I wrote to him, and received a reply dated April 17th, 1931: "I am pleased to inform you that I am still going strong and can eat anything that comes along, from beef-steaks, eggs, and bacon to meat pies, in fact I have not found anything yet that I cannot swallow. Twelve months ago I secured light employment and have continued to work regular ever since, and have had no need to visit my doctor."

There seems to be little doubt that, in the large majority of cases, a cure can be obtained by forced dilatation. In approximately 700 cases at the Mayo Clinic dilatation has been carried out readily and satisfactorily from above, by means of the bag, only three requiring dilatation from below.¹ On the other hand, Hurst² finds that the far simpler treatment, by means of the mercury tube, is effective in the majority of cases, and only reports one case in which no lasting improvement occurred. In this case forcible dilatation by the bag proved equally unsuccessful, and the patient was operated on by Rowlands, who divided the cardiac sphincter in exactly the same way as the pyloric sphincter is divided in Rammstedt's operation for the hypertrophic pyloric stenosis of infants, the operation curing the condition, which had been present for seven years. If treatment is undertaken by means of a bag, it would obviously be an advantage if the bag were opaque to x rays, so that dilatation could be carried out under direct observation and its correct position verified.

A. M. J. Wright, in the *Medical Annual*, 1930, extracting from H. P. Mosher's Semon Lecture, says: "The author employs dilatation with an inflatable bag under manometer control, the bag having barium implanted in four strips in its wall, so that the dilatation can be carried out under direct observation with the x-ray screen."³

REFERENCES

- ¹ Mayo Clinic Collected Papers, 1928, p. 3.
- ² Hurst: *Medical Essays and Addresses*.
- ³ *Journ. Laryngol. and Otol.*, March, 1930, 161.

H. B. BUTLER: ACHALASIA OF THE CARDIA

FIG. 1.—Radiograph showing passage of tube, June 14, 1929

FIG. 2.—Showing an immediate meal distending the esophagus, July 27, 1929.

FIG. 3.—Showing a mercury tube which the patient himself was able to pass into his stomach, July 31, 1929.

FIG. 4.—Showing a bread-and-milk barium meal, the consistency of thick cream, taken immediately after the successful passage of the tube.