Observations

ON

THE TREATMENT OF SEPTICAEMIA*

BY

SIR THOMAS HORDER, Br., M.D.

PHYSICIAN TO ST. BARTHOLOMEW'S HOSPITAL

It is not easy to define the condition known as septicaemia, but the most satisfactory description is: a general infection associated with bacteria in the blood, and their multiplication there. The temporary bacteraemia which appears in the course of pneumonia, typhoid fever, and other such infections may be excluded from the discussion, together with terminal and agonal conditions, and septic endocarditis, which is a special variety of focal sepsis associated with an overflow of bacteria from the endocardial lesion. On the other hand, bacteria may not be found in the blood for one or more of several reasons, and yet a case may be septicaemic; the concept of septicaemia is therefore essentially clinical rather than bacteriological. To escape some of these difficulties changes in terminology have been suggested. Libman declines to use the term septicaemia, but proposes no substitute; Ryle suggests "streptococcal fever" and "staphylococcal fever," and there is much to be said for this innovation. Since the problem centres round a general infection by the two common pyogenic organisms, *Streptococcus pyogenes* and *Staphylococcus aureus*, I suggest that the discussion may well be limited to these.

With the recognition of the nature of the disease the earliest efforts lay in the direction of expectant measures. Then came efforts at chemotherapy, the use of mercury perchloride intravenously being the chief of these, but such crude chemicals as carbolic acid and formalin were also tried. Failures, some disasters, and the arrival of immunology, led to bacteriotherapy—first serums, then vaccines, then sensitized vaccines, and lastly immunotransfusions. Disappointments followed, and there was a swing back of the pendulum to chemotherapy. In the meantime, measures based on protein and chemical "shock" were introduced, and are still on trial. Finally, expectant treatment has again been emphasized, as being the main standby in the majority of cases. The great lack of uniformity in practice demonstrates the absence of convincing methods of therapy.

EXPECTANT THERAPY

Expectant measures aim at getting control of the disease, rather than "curing" it; even to-day we do not "cure" tuberculosis, we "control" it. The methods of control should be instituted at the earliest possible moment, and should be exploited to the full. It can scarcely be doubted that, taken together, they are of greater service to the patient than any "specific" remedy as yet known. If it is possible to give "institutional" rather than "home" treatment, this should be undertaken. Comprised under this heading are rest and fresh air; the patient should be treated in the open air whenever possible. Food should not be stinted, but be given up to the limit of the digestive capacity, and water should be taken in abundance. Alcohol is of service, particularly in patients who have been accustomed to it previously.

Other factors in the treatment include: sunshine, both natural and artificial; eliminative measures for bowels and skin, such as hydrotherapy; the administration of alkalis and glucose to combat acidosis; haematinic drugs

to control anaemia; and nucleic acid with other leucocytic stimulants. A useful formula for intramuscular injection is: sodium cacodylate 1 to 3 grains in 20 minims of a saturated solution of nucleic acid; this may be injected once or twice in the twenty-four hours. Sleep must be secured, and pain relieved. Paget, who suffered from an attack of septicaemia, paid a glowing tribute to morphine. Pyrexia is to be controlled by ample ventilation of the skin and hydrotherapy. Good nursing is imperative, and the personality of the doctor, who should be encouraging, resourceful, and confident, is a great adjunct to success. Although expectant measures lack the dramatic element, and are consequently not popular, they are the only ones which have stood the test of experience.

SPECIFIC TREATMENT

Bacteriotherapy.—Serums have been disappointing, but the theory is so attractive that they are still the most popular remedy on specific lines. "Bactericidal" serums were anti-endotoxic in the main, but their content in antibody has never been such as to satisfy the immunologist. Recently, efforts have been directed towards the production of anti-exotoxin serums; and these "anti-toxic" serums appear already to be more helpful, especially in the case of the streptococcus. They should be used early and in adequate doses. "Sensitized" vaccines (Besredka) seem to be successful in some acute streptococcal cases. Antigen therapy is of little or no proved service; "immunogens," which are ecto-antigens, have scarcely yet had a good trial. Immuno-transfusions have been on trial now for ten years, and it is doubtful if any good effects reported are due to a specific increase in the bactericidal powers of the blood so much as to non-specific aid given by the transfusion. So long as the immunological factors in the interaction between the tissues and the infecting organism are still so largely undetermined, there must needs be doubt as to the best type of specific remedy, but in general infections (as compared with local) there is an *a priori* bias in favour of passive, rather than active, methods. Finally, bacteriophages are on trial and have been commercialized.

Chemotherapy.—During and immediately after the war the chlorine group was in favour. More recently mercurial treatment has been revived, and especially in combination with aniline dyes. The favourite at present is mercurochrome, 10 to 15 c.cm. of a 0.5 per cent. solution being given by the intravenous route on alternate, or even on successive, days. The dangers are thrombosis and renal insufficiency, but successes are recorded from time to time, which may be due to "chemical shock." Novarsenobillon and other salvarsan equivalents have been used freely, following their success in syphilis. The ideal blood disinfectant must diffuse rapidly and widely; it must be non-toxic to the tissue cells, and must retain its bactericidal power over a lengthy period. We need not despair of finding something even better than mercurochrome.

Blood Transfusions.—These are definitely useful, but chiefly to combat the anaemia due to haemolysis.

Shock Therapy.—As regards protein shock, the indications are difficult; probably it should be reserved for the later stages of the disease, and, even then, it should be adopted with caution. Chemical shock treatment is best left to those who understand the teaching upon which the practice is based.

Surgery.—In all cases close co-operation between physician and surgeon is essential. Early in the case this applies especially to the focus of primary infection, if this is known; later, and throughout the course of the disease, it is concerned with the formation of metastases—whether in or around joints, bones, serous membranes, or

* A synopsis of the remarks made in opening a discussion at a joint meeting of the Sections of Medicine and Surgery at the Annual Meeting of the British Medical Association, Eastbourne, 1931.

solid organs. Thorough survey of the patient, almost daily, for these metastases must be made. Effusions are to be dealt with by aspiration for preference, but if this is unsuccessful, then incision and drainage become necessary. Great exercise of judgement is required at both these stages of the disease. Surgery is indicated more often, and more early, in staphylococcal than in streptococcal infections. The favourite sites for staphylococcal metastases should be passed in review again and again.

GENERAL REMARKS

The individual case is to be treated on its merits, due consideration being given meanwhile to any general specific remedy offered by the immunologist, and applicable to the particular infection concerned. The physician should work hand in hand with the bacteriologist, but should not take the cue entirely from him in matters of treatment. Close liaison with the surgeon is a *sine qua non* for success. Expectant measures should be mobilized promptly and thoroughly. All the therapeutic "eggs" should never be put into one basket. A plan of campaign appropriate to the case in point should be decided upon and steadily continued, and the patient should not be disturbed by too many treatments. Panic on the part of the friends is a frequent deterrent, but is pardonable; panic on the part of the medical adviser is sometimes fatal and is unforgivable.

SURGICAL ASPECTS OF SEPTICAEMIA *

BY

D. P. D. WILKIE, M.D., CH.M., F.R.C.S.

PROFESSOR OF SURGERY IN THE UNIVERSITY OF EDINBURGH

The question under discussion to-day, whilst of vital importance and wide interest, is, at the same time, one on which fruitful debate is difficult. Many of us have had considerable experience, of an occasional and isolated nature, of cases of septicaemia; few have had the opportunity of treating a series on truly scientific lines. The features peculiar to the individual case, the intermittent and irregular incidence of cases, our own changing ideas of the value of certain lines of treatment, all tend to render judgement difficult. Further, we are so apt to enthroned in the sunshine of our memory the apparently striking success, and to allow the many failures to slip into the quiet shadows of forgetfulness.

The surgeon may be called on to deal with septicaemia under three sets of circumstances:

- (1) Following an infected accidental wound;
- (2) Accompanying an idiopathic infection, such as acute osteomyelitis, cellulitis, or phlebitis;
- (3) Following operation on an infected area.

In (1) and (2) active surgical intervention may be called for, but nicety of judgement is necessary in determining how much must be done, and when. In many cases it will be found that active surgery must be held in restraint until such time as the indications for intervention have become abundantly clear—in other words, the knife may be a lethal weapon if prematurely or incautiously applied (for example, in the infected puncture wound too hasty incision may induce septicaemia by bursting imperfectly formed natural barriers and throwing open to micro-organisms a pathway into the blood stream). Respect for, and aid to, the formation of such barriers are essential criteria of the successful treatment of such infected wounds.

* Introductory remarks in a joint discussion in the Sections of Medicine and Surgery at the Annual Meeting of the British Medical Association, Eastbourne, 1931.

In the common streptococcal infected wound of the finger or hand, with rapidly developing lymphangitis, the judicious use of passive congestion and administration of antiserum should always precede any treatment by incision. The value of Bier's passive congestion is not sufficiently recognized. Rightly used, I believe that it is a very real help in preventing and limiting blood infection. Bier laid no claim to be the originator of the idea of passive venous congestion as a therapeutic measure. He has, however, fully earned lasting credit for demonstrating how by this means a local infection, which was tending to invade the blood stream, might be hemmed in and held up until such time as local defences had been mobilized, and the general protective mechanism had been stirred to action. Rightly applied, a Bier's bandage will always relieve pain. If the pain persists or is increased, the bandage should be removed and reapplied more lightly. It should be worn for twenty-two hours out of the twenty-four.

Many deaths have resulted from a precipitate use of the bistoury, no time having been allowed for a leucocytic encirclement to develop and a natural localization of the contest to occur. Only when dangerous tension has arisen, or dead or moribund cells have accumulated in the shape of pus, can incision be helpful.

In certain tissues, such as bone and deep cellular planes, where rigid walls prevent the deployment of natural resistant forces, early incision may be of the utmost value in permitting expansion and the entrance of protective elements. Total excision of the focus—for example, in anthrax pustule, in carbuncle, and in the very earliest stages of an infected punctured wound—may sometimes be justified; as a general rule, however, surgery must exercise a watching brief, prepared to step in when, and only when, nature cries for help.

When we consider the various means by which the patient's resistance to the invading germ may be supplemented we turn first to antiserums. Varied opinions will doubtless be expressed as to the value of anti-streptococcal serum. As with anti-tetanic serum, large doses (60 c.cm.) must be given early and repeated daily. My experience has been that it is in the *very earliest stages* of the general infection, before cellular damage to the organs has occurred, that the help from passive immunity is great and real. I have no hesitation in placing it first in importance, just as I believe it should take precedence in time of all other measures.

Copious quantities of fluid must be given, usually by mouth. In exceptional cases where vomiting occurs fluid must be given rectally or subcutaneously. If an abscess develops at the site of subcutaneous injection, it will be a matter for rejoicing. With the fluid, alkalis and glucose must be freely given. As a rule they are both welcomed by the patient, which is significant.

Immuno-transfusion should certainly be considered in the weakly patient, and I have seen it followed by remarkable betterment in a case in which blood infection had supervened in a feeble and anaemic patient. How much of the benefit was due to the introduction of specific antibodies and how much simply to the supply of healthy blood was impossible to determine.

The production of a really efficient anti-staphylococcal serum is a real need, and I trust that the antitoxic serum now on trial may prove to have for the human subject as true and specific a protective action as it has been proved to possess for animals.

The use of vaccines in a fulminating blood infection I have always regarded with scepticism and, along with protein shock therapy, I have left to those who can visualize their action and believe in their efficacy. Sensitized vaccines have given favourable results in some hands, and they undoubtedly deserve an extended trial.