Observations ON TRIGEMINAL NEURALGIA*

BY

GEOFFREY JEFFERSON, M.S.(Lond.), F.R.C.S.

HONORARY NEUROLOGICAL SURGEON, MANCHESTER ROYAL INFIRMARY;
HONORARY SURGEON, SALFORD ROYAL HOSPITAL.

In introducing the subject of paroxysmal trigeminal neuralgia I shall omit any references to the nature of a disease the causes of which are as unknown to us as they were to Tudor physicians. But it will be necessary at the outset to make some allusion to the character of the paroxysms, because adequate treatment depends on the selection of the right cases. Not uncommonly patients are referred as "neuralgias" who undoubtedly have pain in the face and head, yet who are not true sufferers from the paroxysmal disorder, and these may be made much worse by procedures very right and proper for the orthodox case. As to these procedures, the present occasion is important because the time has arrived for a reassessment of values in the methods of treating this condition. The claims of alcohol injections have been most ably and convincingly pleaded in this country in particular; but sufficient evidence is now available to show that operation is in general superior to injection, and should therefore be more widely employed. Alcohol has its place in the treatment of neuralgia, but it must be removed from the dominance which it has held. I shall elaborate the logic of these statements later in this discussion.

CHARACTER OF PAROXYSMAL TRIGEMINAL NEURALGIAS

Although the character of the attacks of pain in true paroxysmal neuralgia is familiar enough, yet an outline of the chief features will not come amiss, for in its severe forms it is not a common occurrence, being slightly rarer in my experience than intracranial tumours. If reflex neuralgias from various local causes were added, they would no doubt more than adjust the numerical balance. Clearly, comparative experience must vary, but I believe that I am correct in the main on this point.

Sex.—The predominance of neuralgia in females has long been recognized, but since we do not know the cause of the disease no important conclusion may be drawn, nor is the sexual disparity sufficiently marked to lead in itself to any special deduction.

Unilaterality.—The pain is always confined to one side, and generally to one division at the commencement, either the second or third. Sometimes the pain affects two divisions very early, but, as a rule, it remains limited at the beginning and only spreads after months, or often years, have elapsed. The pain keeps within the confines of the trigeminal field, and spread outside it into the neck or shoulder will make us suspect some other cause, though overflow is sometimes genuine. In some cases it may reappear on the opposite side, limited to a single division, generally the second. As the pain may appear on the opposite side long after it has been successfully treated on the primary side, the sparing of the motor division is a wise precaution at operation. Fortunately, this step in no way increases the likelihood of recurrence, nor is the resultant anaesthesia any the less.

Nature of the Paroxysm.—The pain is always of brief duration, and when the storm has passed the patient is generally comfortable, though sometimes a tingling or stinging sensation endures. But whether this slight middle pain is present or not, the patient lives in dread of a return

of the torment, and even after successful treatment this apprehension of the future persists for a long time. The pain is brought on by a touch, by a breath of wind, by a movement, so that smiling, talking, and eating become a penance. A facial pseudo-palsy develops in some patients; the face becomes almost Parkinsonian in its mask-like appearance, actually a purely voluntary defensive inhibition. The patient sometimes swathes herself in wrappings and brings with her someone to talk for her, a "talker," in the same way that a disabled batsman at cricket may be allowed a "runner." The descriptions of the nature of the pains vary somewhat, but it is clear that neuralgic pains are sharp, and lancinating, like long needle thrusts or stabs. It is probable that the differences in neuralgic pains are differences of description rather than real disparities in essence, and vary with the patient's powers of expression. This point is important, and may be useful in diagnosis, for if we can standardize, as it were, neuralgic pains, we can more easily differentiate between true "neuralgias" and those who suffer from pains in the face of another kind. Certain it is that the pain is essentially different from the more common burning, throbbing, duller, and continuous pain of an inflammatory lesion, and when a person who is thought to have neuralgia uses such terms as the last to describe the pain, we shall rightly suspect that we are faced with something other than true trigeminal neuralgia—very possibly a closed infected antrum with secondary neuralgia. In much the same way that the pain is differently described by different persons, so the expedients to which they resort to mitigate, or to prevent, the attacks vary with individual ingenuity and resources.

Facial Spasms.—The term "tic douloureux" is not a good generic title, as in many patients facial spasm is not an important feature. Often during severe attacks the pain causes irregular contractions of the facial musculature, but they are seldom of a severely spastic or clonic nature. Painful cramp-like facial spasms may occur in cases which are not essentially due to trigeminal neuralgia, spasms which are not stopped until the seventh nerve is blocked. These cases are very rare, but must be distinguished as a separate entity. In trigeminal neuralgia the pain comes first and facial movements are secondary, followed as the pain passes off by lachrymation, salivation, and flushing of the capillaries of the face (a sympathetic phase).

Remissions and Intermissions.—As already mentioned, the painful spasms are in general of brief duration; hence the pain commonly disappears altogether for long periods. In fact, remissions of symptoms are as characteristic of major neuralgia as they are of duodenal ulcer. This is a fundamental characteristic. The pains may disappear for weeks, months, or even years; and here we have the explanation of the fact that so many patients are able to give such long histories of their malady. Generally this is well recognized by the patient, who admits that were it not for the free periods he would have gone out of his mind or sought death. At the same time the knowledge that after a few weeks or months of agony the pain will probably go, and the hope that it will not return, encourage the victim to struggle on with inadequate medicinal protection against his sufferings. With this must be coupled the undeniable fact that doctors themselves too often hesitate to advise radical surgical treatment. I shall refer to this point again later.

The remissions from pain are most irregular, and although in some patients there is a seasonal incidence in spring and autumn, this is by no means general. Cold in my own series has been more beneficial than heat, and there is little hope that a stay in a warm or sheltered watering-place will give the relief so eagerly sought, and sometimes found in other chronic diseases.

* Read in opening a discussion in the Section of Surgery at the Annual Meeting of the British Medical Association, Eastbourne, 1931.

Sites of Election.—The pain of trigeminal neuralgia is generally strictly anatomical in its distribution. In my cases it has commenced rather more often in the second division. Probably the second and third divisions are affected more or less equally, but the slightly greater incidence in the second division in my own series is worth noting, because it used to be taught that neuralgia began in the third division and spread upwards to the second. This is certainly not the rule.

First Division.—A point on which there can be no dispute, however, is that the first division is rarely affected first (2 per cent. of my series). Many patients with severe neuralgia complain of pain passing up through the eye to the forehead from the second division, but this is a pure overflow and stops when the pain in the lower part of the face is put to an end. Further, pain high in the temple may originate in the auriculo-temporal branch of the third division and from the zygomatico-temporal branch of the second, and may be described by our patients, who are not anatomists, as "pain in the forehead."

The freedom of the ophthalmic division from primary neuralgia is of immense importance when we come to treatment, for it is evident that we can, if we choose, omit this part of its predestined deep fibres from destruction without fear of subsequent pain in this area. The recognition that this branch can be spared has profoundly modified the surgical treatment of neuralgia.

Second Division.—The pain is generally in the soft parts of the face, and especially in the upper lip and beside the nose. It may also affect the gum about the canine fossa, but does not often spread to the palate. Second division neuralgia tends, therefore, to be seated in the soft parts, and is, if one may call it so, an anterior pain. It may, of course, extend into the full distribution of the second division, including the posterior palatine nerves, the upper gum, the temple, as already mentioned, the side of the nose and the eye, and overflow beyond these confines on the same side but not to the other. On the whole, however, the presence of pain in the upper lip and the fear of the lip and cheek being touched are the striking features of neuralgia in this division.

Third Division.—The pain in neuralgias of this division is chiefly in the bone and gum, going back to the tragus but not so often into the lip or tongue. Indeed, the freedom from severe pain in the tongue is notable; even in well-established cases it may dart there only occasionally. Third division pain is therefore more often a deep-seated pain and is not so tiresome in the lip as with the second division, though the pain is in itself no less severe.

"Atypical Neuralgias."—In very marked contradistinction to the above is a most difficult group, which is at present classified under the non-committal title of "atypical neuralgias." The outstanding feature of these cases is the non-paroxysmal nature of the pain, which is continuous night and day, often for years. It is described as a burning and often as of a pulling kind, and is generally felt all over the trigeminal field and often beyond it from the start. The pain is equally described as unbearable, but it never remits and it never causes those sudden eloquent silences, those postures and anguished expressions which are diagnostic of true neuralgia. In fact, when the patient, in the course of describing the pain which has been so continuously present, is interrupted and asked whether the pain is there now, the invariable answer is, "Yes, of course, it never ceases," although his demeanour has given the clinician no special inkling that he was indeed in pain at the time. This is quite diagnostic. These patients are generally younger than the true "neuralgias," often in the twenties or thirties, an

age at which paroxysmal neuralgia is rare, though not unknown. Further, the pain is often centred about the eye and affects the forehead and scalp more than the rest of the face, radiating to the neck, shoulder, and arm. Both sides are commonly affected, though one may be much less troublesome than the other. It is important that we should differentiate this type because surgical treatment is entirely useless, and indeed should be avoided, as it commonly makes the condition worse since all treatment for pain is essentially destructive. Such steps give the patient a positive lesion with which to tempt some other surgeon to extend the damaged area in the hope of cure—a praiseworthy but vain aspiration. (See excellent reviews of this subject by Glaser and by Stewart and Lambert on spheno-palatine neuralgia.)

Glosso-pharyngeal Neuralgia.—This is a rare disease but one which is easily confounded with trigeminal neuralgia. There has been only one case in my series, and this will be separately reported. The pain is equally spasmodic, but is generally localized to the depths of the ear and to the fauces or to one or the other, an elective tendency; it does not spread to the face. If such a distribution is localized to the ear, it may delude the surgeon into thinking that it is a curiously local example of third division neuralgia, seated in those twigs which supply the tragus, auditory meatus, and tympanum. The pain is more localized and more deeply seated than in trigeminal neuralgia, and is brought on not by chewing and moving the jaws but by swallowing. The treatment is of course resection of the glosso-pharyngeal nerve.

Post-herpetic Neuralgia.—Brief mention must be made of post-herpetic neuralgia. It is well known that herpes is a distressingly painful affection, but in the young or moderately young it passes off without any legacies but scars. In the aged it is almost always followed by pain of the most insistent and wearying nature. This pain, like that of the atypical neuralgias of younger persons, is continuous and unremitting, and it is my experience, and the experience, I think, of all of us, that long-standing continuous pain is intractable to surgery unless it is secondary to chronic malignant disease. Occasionally post-herpetic neuralgia is spasmodic, and in such cases there is more hope of surgical cure.

Tumours and Gummata of the Gasserian Ganglion.—Lastly we have those cases in which pain in the trigeminal field is caused by tumours or focal leptomeningitides affecting the root of the nerve in the posterior fossa, by tumours or aneurysms involving the ganglion in the middle fossa, or by nasopharyngeal or other growths infiltrating the issuing divisions. The differentiation of neuralgias due to these conditions is a fascinating subject, but it has no place in the present discussion. Fortunately we have a touchstone by which these causes can generally be eliminated immediately—namely, anaesthesia or analgesia of the affected area. In true paroxysmal neuralgia there is never any alteration in the sensory perception of the skin, unless it be a certain hypersensitiveness. This is a fundamental point in diagnosis. If cutaneous anaesthesia is found on testing (anaesthesia dolorosa), then we know at once that the case is one in which there is some definite lesion of the trigeminal or its branches, and a full study must be undertaken to discover the cause. I am aware that in rare cases acoustic neuromata have been discovered unexpectedly in the course of an attack through the posterior fossa on the sensory root of a patient thought to be the subject of a plain neuralgia. But such isolated experiences do not in my opinion imply, as some would have us believe, that this is a sufficient reason for the routine adoption of this approach. The fact is that neuralgia is a really rare

leading symptom in lateral recess tumours. Severe neuralgic pain sometimes occurs in malignant disease of the mouth, and no little success may be obtained by a well-planned nerve section, which gives astonishing relief. Mention must be made of trigeminal neuralgia, often with a large overflow, occurring in disseminated sclerosis. These cases can be helped by surgery. The causal lesion seems to be a patch of sclerosis in the pons in the area of entry of the sensory root (Parker).

DIFFERENTIAL DIAGNOSIS OF PAROXYSMAL NEURALGIA

The foregoing account should make clear enough the characteristics of major or paroxysmal neuralgia, and also those states most likely to incite it as an auxiliary phenomenon. To recapitulate: the patient is usually female, the pain is limited to one or two divisions save in the worst cases, the attacks are of brief duration, and between the pains the patient is moderately comfortable though apprehensive. There is a strong tendency towards remissions in which the pain is negligible or entirely absent for weeks or months. There is no alteration in the sensibility of the skin, certainly no diminution. The pain of secondary neuralgia is very similar, and when due to dental causes it is easily remedied. Antral infection, however, is more insidious. It might be assumed that if pain corresponding to the paroxysmal type has been present for a long period, possibly for two or three years, and there is no history obtainable of disease in the nose or of purulent discharge, paranasal infection cannot be present. This would be an unsafe conclusion, for I have known of such cases, and I firmly believe that x-ray studies of the sinuses should be made as a routine if error and disappointments are to be avoided. Generally these infected patients admit having had occasional swelling of the face or, what is more important, of the lids, an oedema lasting for a day or two. Flushing and a sensation of subjective feeling of swelling occurs in some paroxysmal cases due to disturbance of sympathetic control during the attack. This is of short duration and may usually be distinguished from a real oedema.

TREATMENT

It is just forty years since Horsley published a paper entitled "Remarks on the various surgical procedures devised for the relief or cure of trigeminal neuralgia." All the operations there reported, thirty-one on nineteen patients, were peripheral save one—a "section of the fifth nerve from the bulb" by a transtentorial approach ending fatally. The years that have passed have seen many further stages in the struggle between the surgeon and a nerve which shows the most astonishing and obstinate powers of regeneration, powers which would delight us if they were reproduced by, say, the median or ulnar at the wrist. The story of the surgery of neuralgia is one of a slowly advancing attack from the periphery inwards to the ganglion and finally to the sensory root. The names of many famous men are associated with this movement, especially Rose, Horsley, Krause, and in particular Jonathan Hutchinson junior.

Operations on Gasserian Ganglion

Hutchinson perfected the removal of the lower two-thirds of the ganglion with the preservation of the ophthalmic division and that portion of the ganglion which is proper to it. This he did for three reasons: first, because primary neuralgia is rare in the first division and when present is generally an overflow, so that the division may usually be left intact; secondly, in order to preserve sen-

sation in the eye and to prevent the neuro-paralytic keratitis which is so apt to follow complete denervation of the cornea; and thirdly, because by leaving it he avoided the chief danger of the operation at that date, injury to the cavernous sinus. Hutchinson brought this operation to a high pitch of efficiency, for in 1919 he reported 70 partial excisions of the Gasserian ganglion with 3 deaths (4 per cent. mortality) and refers to Horsley's (1908) 120 cases with 6 deaths (5 per cent.). Horsley's final figures were never, I think, published. These English figures compared extraordinarily favourably with those from foreign clinics, and were unquestionably far better than those of surgeons with less experience in British and other centres.

The Gasserian ganglion in its upper and anterior part lies in the wall of the cavernous sinus, the ophthalmic division penetrating it. The whole ganglion cannot be removed without injuring the sinus, and misguided attempts to do so were responsible for the heavy operative mortality that years later still hangs like a cloud over the subject. Hutchinson's operation was well conceived in that he removed only such part of the ganglion as could be extirpated without opening the cavernous sinus, though even so, a small opening is almost inevitable when the cut second division is stripped back. None the less the Hutchinson operation has been abandoned in favour of another of which he was himself, curiously enough, extremely sceptical. I refer of course to section of the sensory root by the temporal extradural route, the advantage of which is that the cavernous sinus is not a hazard at all, for the operation has nothing to do with it.

Methods of Division of Sensory Root

This operation, generally credited to Spiller, has been perfected by his surgical colleague Frazier, who has recently recorded 511 cases, with 1 death in the last 269 cases (0.37 per cent.). Prior to this, Cushing in 1920 had reported 332 Gasserian operations with 2 deaths (0.6 per cent.), most being root sections, others partial gasserec-tomies. Similar figures have been recorded by Adson. Cushing's statistics certainly provided a tremendous stimulus, for ten years ago his low mortality rate in such a large number of cases seemed well-nigh incredible.

Three innovations have contributed to the improvement in operative results: the sitting posture in the dental chair, the use of illuminated retractors, and, lastly, the employment of suction as a means of keeping the field clear.

I have personally performed the operation on eighty-five occasions with two deaths, one from an unknown cause early in the series in an exceedingly bad subject, the other a death from pulmonary embolism on the ninth day in a patient making the usual smooth post-operative recovery. This mortality is higher than I would wish, but it is only common sense to admit that occasional fatalities must occur when we are dealing with persons of the age and debility of some "neuralgias." Neither of my deaths came from technical error, and indeed it is my purpose in this address to show that the factors which can bring about morbidity and fatality are so few that such deaths as occur may be attributed more to bad luck than to the inherent hazards of the operation. I have carried out fractional root sections five times under local anaesthesia (which has its special points), helped out by a little inhalation anaesthetic when the root was being worked upon. And I have twice cut the sensory root on the second side, preserving the motor root, after the ganglion on the other side had been destroyed by alcohol by the Härtel method. The precise mortality in decimal points of zero is not of great importance: what is important is

that it is clear that with good surgery and good fortune only one patient in a very large number is going to die after the operation. Yet it would seem that practitioners the world over still think of Gasserian operations as having the same mortality as in the immediate post-Listerian period, or so one would judge from their attitude towards it. It is a fact, of course, that the operation of section of the sensory root is a very difficult operation to learn and is at times an exceedingly difficult operation to perform, though at other times it may be simple.

It is clear from the low mortality of this operation in large series of cases that there must be something essentially benign, some factor of risk must be lacking, as compared with even the partial Hutchinsonian resections. This logical assumption is correct and it lies in the avoidance of any likelihood of damage to the cavernous sinus. It is true that the sinus and even the carotid artery may be damaged in this operation with fatal results, but only when the surgeon has gone astray, and these structures are not normal risks when the operation is well performed. The sensory root can be picked up at the posterior end of the cavum meckelii and the actual section carried through without the least loss of blood. It is imperative that the field should be entirely dry before the final exposure of the root in its cerebro-spinal fluid sac, and it is in effecting this that the surgeon may have to use all the skill at his command, not because of bleeding from the cavernous sinus but of bleeding from veins in the dura. Some individuals have none of any note, others may be well endowed.

Advantages of Fractional Division

Total division of the sensory root, however, is open to the dangers that come from an insensitive cornea and is in this respect inferior to the Hutchinsonian operation, though superior to it in point of mortality. Realizing this, Frazier has applied the Hutchinsonian principle to the root section, preserving those fibres in the root which supply the eye, and Adson has added a further refinement in the preservation of the motor root. This operation, which is now the accepted standard operation among neurological surgeons and the one which I perform, is referred to as "fractional division of the sensory root." It is fortunate, indeed, that the arrangement of fibres in the root of the trigeminal nerve allows, as Frazier has shown, of such manoeuvres being carried through with success. There is no doubt a slight interchange and a little interweaving of fibres, more in some individuals than in others, but in the main the fibres preserve a predestined course in the root, those to the ophthalmic division lying innermost and above, those to the third division lying below and outside, those to the second occupying an intermediate position. The parallel arrangement of the fibres is well shown in Fig. 1. The motor root lies behind the sensory, crossing from above downwards, and is free from the sensory fibres in that part of the root exposed at operation. The safest way of preserving it is, in my experience, to cut through the root from without inwards with long fine Stille scissors, until a hole is made through which the motor root, running in a slightly different direction, is clearly seen. In cases in which alcohol injections have been made into the ganglion beforehand, a satisfactory fractional section is much more difficult to carry out, for there are many more adhesions, the fibres do not separate readily one from another as they ordinarily will, and the motor root is commonly matted to the sensory part. These are serious obstacles to clean operating, and it is desirable that we should attempt to standardize our treatment of neuralgic patients and avoid making ganglionic injections in patients who are suitable subjects for fractional section.

Results of Fractional Division

Personally I make a practice of dividing the lower and outer two-thirds (lower and outer because the root lies obliquely with its flat upper surface looking upwards and outwards). The resulting anaesthesia affects the lower two-thirds of the face and half the cornea. In many cases

FIG. 1.—The sensory and motor roots of the trigeminal nerve, from below (left) and above (right). Note the parallel fibres in the sensory root.

the line of anaesthesia runs strictly through the pupil, the lower half of the cornea being insensitive, the upper normal. Figure 2, which is the work of an entirely independent observer on cases operated upon by me, shows the average result. The retained sensation in the cornea has been sufficient to prevent the neuro-paralytic keratitis of which Paton made so masterly a review. From the recurrence in a series of cases of this picture of partial ocular anaesthesia it becomes abundantly manifest that we have in this operation a method of attack on trigeminal neuralgia which seems to meet all our needs. In two or three cases I have ventured with success to cut the middle

FIG. 2.—Average area of anaesthesia after fractional section of the sensory root. (F. N. Marshall.)

portion of the sensory root alone (see Fig. 3) in patients whose chief pain was referred to the second division. The anaesthesia following partial divisions is never dense, but the operation none the less stops the neuralgic pains. This partial preservation of sensation is a great boon to the patient. We have an anomalous situation in the persistence of sensation after fractional root division, for if

anaesthesia is incomplete after an alcohol injection, the pain is almost certain to return (though there are cases in which it does not); on the other hand, if the face is not entirely numb in the painful area after root section it does not seem to matter. No doubt the explanation is to be sought in the site of the interruption of conduction in the two operations. It must be conceded that on occasions the motor root cannot be isolated with certainty, but its conservation should be aimed at. Marshall's studies on my cases show that there is no lessening in the anaesthesia when it has been spared.

FIG. 3.—Diagrammatic schema of fractional root section with preservation of motor root. Ophthalmic fibres preserved.

It has been claimed by those who urge the division of the root in the posterior fossa by a cerebellar approach that fractional section stops the pain without producing a totally anaesthetic face. This same result, in my experience, follows fractional section in the middle fossa, and is not a specific result of the immediately pre-pontine section. It is, indeed, a well-recognized result of root section and partial nerve section elsewhere that analgesia is produced over a wider area than tactile anaesthesia.

Alcohol Injections

In view of the mortality following Gasserian operations in the remote past it is not a matter for wonder that attempts were made to find some less lethal method of treatment. Peripheral alcohol injections have proved their usefulness, and in the hands of some, notably of Harris in this country, they have proved of great service. I believe that it is wise to advise a patient to have at least one peripheral injection before proceeding to operation, especially in the less severe case clearly limited to one division. The advantage is that he has a foretaste of the numbness (and that curious and frequently accompanying feeling of swelling) which will be permanent after the root section, an anaesthesia which must always come as something of a surprise. Pre-operative advice, though useful, can never adequately describe the actual feeling. However, if a fractional root division is to be performed preliminary injections are not so necessary, as some sensation is retained. As to injections into the ganglion itself, whether by Harris's lateral method of infiltration or by the anterior direct placement of alcohol into the ganglion and root as devised by Härtel, and developed by my colleagues Rayner and Morris in this country, I believe that this method should be restricted to those who are judged to be physically incapable by reason of their age or of coexisting disease to undergo operation. I have arbitrarily fixed 70 years as a limit, but I have no doubt that even some older persons could be shepherded through a root section if it were necessary. In younger persons and especially those in the forties and fifties, and those whom we term a "good" sixty or more, I have no hesitation whatever in saying that fractional root section is infinitely the better course. The matter is capable, I believe, of precise and logical statement. If operation had nothing better to offer than a complete and dense anaesthesia of the face with the risks consequent on total interruption of the ophthalmic fibres, there would be nothing

to choose between such an operation and alcohol injection—indeed, the balance would be in favour of the latter except after its repeated failure. But fractional root section offers something which alcoholic destruction of the ganglion cannot give—namely, preservation at will of such fibres as we wish to keep. This being the case there seems to me to be little room for argument as to which is the method of choice, though the exigencies of circumstances may at times induce us to use a method which is both theoretically and practically inferior. Alcohol injections are in any case more uncertain than operation. The not infrequent result of injections is that we achieve rather less than we hoped for, while at times, using a precisely similar technique, we obtain more, unfortunately, than we bargained for. We must not forget that trigeminal neuralgia is a really serious affection, and we do not make it less serious by using a method which is to the casual eye a simple and innocuous one, and yet may require much repetition and bring about an occasional catastrophe. But in the aged and the infirm we must continue to use this somewhat uncontrollable method.

BIBLIOGRAPHY

- Horsley: *British Medical Journal*, December, 1891.
 Rose: *Surgical Treatment of Neuralgia*, London, 1892.
 Hutchinson: *Facial Neuralgia and its Treatment*, London, 1919.
 Harris: *Neuritis and Neuralgia*, London, 1926.
 Cushing: *Amer. Journ. Med. Sci.*, 1920, clx, 157.
 Frazier: *Annals of Surgery*, September, 1928, 534.
 Adson: *Surg. Gynecol. and Obstet.*, September, 1922, 352.
 Parker: *Brain*, March, 1928.
 Glaser: *Arch. Neurol. and Psychiat.*, 1928, xx, 537.
 Stewart and Lambert: *Journ. of Laryngol. and Otol.*, November, 1930, 753.
 Heymann: *Deut. Zeit. f. Chir.*, April, 1929, i.
 Marshall: Thesis, Manchester University, 1931.
 Paton: *Brit. Journ. of Ophthalmol.*, June, 1926, 305.
 Rayner: *Brit. Journ. of Surg.*, 1920, vii, 516.
 Morris: *British Medical Journal*, January, 1931, 122.

CHRONIC MICROCYTIC ANAEMIA*

BY

L. J. WITTS, M.D., F.R.C.P.

ASSISTANT PHYSICIAN, GUY'S HOSPITAL; WILL EDMONDS CLINICAL RESEARCH FELLOW

In opening a discussion on the classification and treatment of the secondary anaemias one is at once faced with the difficulties of the nomenclature. The adjective *secondary* was originally applied to anaemia whose pathology was well understood, in contrast to *primary* anaemias such as pernicious anaemia, whose intrinsic nature was obscure. Recent advances in knowledge have done much to obliterate this distinction. We now have more information about the etiology of pernicious anaemia than of many anaemias called secondary. An additional meaning has, however, become attached to the term secondary. In anaemia from simple causes, such as haemorrhage, infection, or toxæmia, the haemoglobin is decreased out of proportion to the red cells, which are often smaller than normal, and the colour index is low. When we say that an anaemia is of the secondary type, we usually mean only that the colour index is low. It is in this sense that I shall interpret the title of our discussion. I hope to show that very little is known of the pathology of many cases of anaemia of low colour index, and that there is a large and coherent group which forms an independent malady, a disease *sui generis*.

The adjective *chlorotic* has similarly been so devalORIZED as to have lost all meaning. True chlorosis has practically disappeared.⁷ There can be few medical men under the age of forty who have ever seen a case of "green-sickness." To-day anaemia is uncommon in adolescent

* Read in opening a discussion in the Section of Medicine at the Annual Meeting of the British Medical Association, Eastbourne, 1931.