to a direct injury, would have been very much worse. As this haemorrhage is absorbed so will the function of the pituitary increase, and the patient already shows signs of improvement in that less pituitary extract is required to control the thirst and polyuria. The necessity for early recognition of diabetes insipidus cannot be too strongly emphasized, as the polyuria in time has a deleterious effect upon the kidneys and bladder. Pituitary extract hypodermically is without doubt the treatment of choice, but it must be remembered that it is not a cure. The treatment must be maintained and the co-operation of the patient sought.

In conclusion, I desire to express my indebtedness to Mr. Blair for the facilities he has afforded me in observing this case.

BIBLIOGRAPHY

- Eaves: *Brain*, April, 1930, liii, 47.
 Moorhead: *Irish Journ. Med. Sci.*, March, 1930, p. 109.
 Williams: *Endocrinology and Metabolism*, edited by Barker, 1922, iv, 861.
 Schäfer: *Essentials of Physiology*.
 Poulton: *Practice of Medicine*, p. 595.

CONGENITAL PYLORIC STENOSIS*

BY

C. MAX PAGE, D.S.O., F.R.C.S.

SURGEON TO ST. THOMAS'S HOSPITAL AND THE VICTORIA HOSPITAL FOR CHILDREN

Although the morbid anatomy and clinical features of congenital hypertrophic stenosis were clearly defined by Hirschsprung in 1888, no striking advance in the treatment of the condition was effected till the introduction in 1913 by Rammstedt of the simple operation of section of the sphincter. Till then treatment was essentially medical. Surgical intervention was only given a chance in obstinate cases and was associated in general with a mortality rate of about 50 per cent., although a few operators using Loreta's method of dilatation quoted better results. With Rammstedt's operation the reported mortality varies from 5 to 25 per cent., and it has become almost the standard method of treatment.

It is noticeable in the literature, however, that the swing-over to surgical treatment has by no means been general, particularly in Germany and Denmark. As late as 1927 Professor Monrad of Copenhagen reported 228 cases of the condition all treated medically and with an over-all death rate of only 14.4 per cent. It may therefore be a matter of dispute how far the extended use of the operation has reduced the mortality. One undoubted advantage it has by common acceptance—namely, the shortening of the period of treatment. The wide variation in the mortality reported by various observers shows that the problem is affected by several factors. Both pre-operative and post-operative treatment influence the result as much perhaps as the method of operation.

The fact that I am allowed to open this discussion is, I think, evidence of how far in this country radical surgical treatment has been accepted. I would emphasize, however, that the surgeon plays only a small, if important, part in bringing these cases to a successful issue. The direction of the pre-operative and post-operative treatment requires the greatest judgement and care of a skilled paediatrician if satisfactory results are to be obtained. As opener I propose to outline the general points in relation to diagnosis. I shall consider in some detail the operative

technique and its bearing on results, leaving it to those who follow me to consider in detail the more difficult medical and biochemical problems on which they are so much better qualified to speak.

SYMPTOMS AND DIAGNOSIS

The cause of the pyloric hypertrophy remains a matter of speculation. The fact that it is commoner in males than in females (4:1) and in the first-born is accepted. Although obstructive symptoms are seldom in evidence till the second week of life, it would seem probable from the size of the tumour even at the earliest stage that the hypertrophy develops during intrauterine life. The youngest patient I have operated on for the condition was 18 days old, and the oldest was 6 months. It does not seem to me to be necessary to discuss whether spasm and stenosis of the pylorus should be distinguished. In my experience some degree of hypertrophy is present in all cases, and even in its fully developed form symptoms are produced by spasm and not stenosis in its true sense.

The points determining diagnosis are, I think, well established, and do not provoke controversy. It is based on the presence of repeated projectile vomiting, visible peristalsis of an enlarged stomach, coupled with wasting and constipation developing about the second week of life in an infant till then apparently normal. A tumour may be felt at the pylorus, but evidence of this is certainly not necessary to establish the diagnosis. I have noticed at operation wide variation in the size and consistence of the tumour and also in the depth at which it lies under the liver. Radiography gives interesting pictures for study and definite evidence of the degree of obstruction, but it is not essential as a routine diagnostic method.

WHEN TO OPERATE

The decision as to when a case should be operated on is no doubt difficult, and will vary with the facilities available. My own impression is that once the full range of symptoms above described is established, the sooner operation is carried out the better will be the results. I think my colleague, Dr. Jewesbury, who has passed on to me the majority of the 150 cases I have operated on, takes this view. If serious emaciation has developed, the risk of operation is certainly gravely increased.

TREATMENT BEFORE OPERATION

Pre-operative preparation should, I think, consist in regular washing out of the stomach for a few days and the administration of subcutaneous glucose saline for a similar period. Regular small feeds of a suitably diluted milk given at two-hour intervals are continued. I do not know how far the administration of atropine is useful, and I shall be interested to hear the views of those who follow me on this subject. To them I shall leave the fuller discussion of this problem and the strictly medical treatment, as they are better qualified to express an opinion.

OPERATIVE TECHNIQUE

It is clear that the choice and administration of the anaesthetic in these infants with so frail a hold on life is a matter of importance. Gray and other surgeons at the Great Ormond Street Hospital consider gas and oxygen to be the best. In Germany chloroform seems to be popular. It was used by Kirschner, who records 15 cases without any deaths. Pauchet prefers local anaesthesia. In my own experience light ether anaesthesia given on an open mask has been satisfactory. It is certainly simple to give and readily controllable. In two cases in which death

* Read in opening a discussion in the Section of Diseases of Children at the Annual Meeting of the British Medical Association at Eastbourne, 1931.

followed closely on operation it was observed that the anaesthesia had been carried to the full degree of relaxation. This, I think, is to be avoided. The operation can be easily performed without full muscular relaxation, although irregular respiratory movements are undesirable during the few minutes the peritoneum need be open. I have carried out the operation under local anaesthesia in three cases, but I think that the movements of the infant and the tendency to the extrusion of the fine omentum and viscera add unnecessary shock and difficulties to the procedure.

I make a vertical incision about one inch in length, half an inch to the right of the mid-line situated half-way between the xiphisternum and the umbilicus. The rectus muscle is split and the underlying peritoneum opened to the full extent of the wound. This exposes the right lobe of the liver. An incision lower down may make delivery of the pylorus easier, but it does not control the other viscera so well either during operation or during the healing period. The lower angle of the wound is then drawn down with a retractor, the distended pyloric antrum thus coming into view. This is drawn out of the wound with forceps and will lead to the pyloric tumour. The delivery of the latter is generally easy with gentle traction by a rat-toothed forceps, but sometimes it is less mobile, and a special ring forceps may be convenient in order to bring it to the surface. The section of the peritoneum and superficial muscle fibres of the tumour is made with a knife, the pylorus being held between the thumb and finger of the left hand. The division of the sphincter is completed with a blunt dissector. There can be no doubt when the section is complete, as the white mucous membrane prolapses into the wound. Care should be taken at the duodenal extremity of the swelling not to puncture the mucous membrane, as at this situation the muscle tumour stops abruptly. On its stomach side the line of demarcation from the normal musculature is gradual and no harm is done by extending the incision well on to the pyloric antrum. The section is made along a line on the pyloric tumour where the surface vessels cannot be seen. If this line is chosen and if the division is completed with a blunt dissector, I do not think there is any need to control the haemorrhage by ligature. If a bleeding point of any size is seen it can be under-run with fine gut. In my experience the necessity for this is very unusual. The pyloric tumour is then slipped back into the abdomen and the wound sewn up in two layers with continuous fine catgut. The section is sutured with interrupted salmon-gut sutures. The operation performed in this way takes about five minutes from start to finish. It should not involve exposure of any viscus other than the pyloric end of the stomach.

AFTER-TREATMENT

In after-management the feeding is important. It must be remembered that, though the stricture has been divided, a dilated and hypertrophied stomach is present, and there may be some mild infection of the gastro-intestinal tract. Regular and dilute feeds are given from within an hour or two of the operation. A little vomiting is not uncommon in the first day or two, though it seldom reaches the projectile type.

COMPLICATIONS

The complications from the surgical standpoint which may be met with are: wound infection, prolapse of viscera, recurrent vomiting, hyperpyrexia, broncho-pneumonia, and gastro-enteritis.

A mild degree of infection of the wound is more common than one would expect. It may take the form of a mild

red infiltration. I think this must be due to the irritation of the catgut. I shall be interested to hear the views of other surgeons on the question of suture material for these cases. I have observed prolapse of omentum in one case and prolapse of gut in another. In the first instance, through-and-through sutures only had been employed to enclose the abdominal wound, and a fragment of omentum prolapsed between two of them. The omentum was ligatured and cut off flush with the wound, and no fresh complications were observed, the patient making a normal recovery. In the other instance the wound suppurated and broke down about the fifth day, allowing prolapse of stomach and intestines. Re-suture was promptly carried out, but the child died of general peritonitis three days later. I notice in the literature that recurrent vomiting of a projectile type has been recorded with some frequency. In my experience it has only once been necessary to reopen the abdomen and carry out a secondary operation. In one fatal case relief of stenosis was incomplete; no secondary operation had been performed. As I mentioned above, a certain amount of forcible vomiting may occur for a day or two, and need not cause alarm. Some measure of hyperpyrexia is observed in about 10 per cent. of cases operated on. I am not quite clear about its significance. In one case the temperature rose to about 108°, and the baby died on the second day. In other instances it has risen to 106° and 107°, and has been brought down by sponging; it does not appear to have interfered with a successful result. Broncho-pneumonia was responsible for two deaths at a fairly early stage subsequent to operation. I do not think it common enough to contraindicate the use of ether, though no doubt these fatalities may be quoted as an argument against its use. Gastro-enteritis was the cause of death in two cases. In regard to this latter complication, it has been very much less in evidence since operation has been undertaken at an earlier stage and the after-treatment has been somewhat modified.

In regard to the function of the stomach and the condition of the children after this operation—as far as cases under Dr. Jewesbury have been observed—it appears that the child becomes normal both in physique and in its digestive functions.

MORTALITY

The 150 cases in which I have operated, from 1922 up to date, may be divided into two groups, namely, 60 carried out at the Victoria Hospital and Princess Mary Home, and 90 cases at St. Thomas's Hospital. It is interesting to note that the mortality in the first series averages 6.6 per cent.; in the St. Thomas's cases it is 12.2 per cent. It is not altogether easy to explain the difference in results, as the surgical and medical treatment was carried out on the same lines, though it is perhaps true that a larger number of cases at St. Thomas's came under care at a late stage of the condition, but it brings home to one the difficulty in drawing conclusions from a small group of figures. I would quote as a further example of this the fact that in 1924-1925 at St. Thomas's 14 successive cases were operated on without any fatality.

CONCLUSION

To sum up, I think one may say that the modern surgical treatment, if it is applied early and associated with vigilant medical supervision and skilled nursing, may be expected to reduce the mortality of this condition to 5 to 10 per cent. This estimate may not appear to be very optimistic, but it should also be remembered that the treatment curtails very greatly the period of hospitalization.