measures were directed to careful suture of skin and mucous membrane, and prevention of adhesions. One week later he presented the appearance shown (Fig. 7 on Plate); even in this case the tip of the nose remains central. The fragments of the nasal bones, maxillae, and ethmoid were carefully manipulated into place under general anaesthesia and retained on the triangle splint. The nasal airway is perfect, and the cosmetic result impaired only by the scar (Fig. 8 on Plate).

SUMMARY

The common type of "broken nose" is due to lateral (but not backward) displacement of both nasal bones with imbrication. Treatment consists of (a) disimpaction, (b) free mobilization of both bones, and (c) symmetrical moulding together. No splints are needed.

When the nose has been crushed, internal splinting is required to restore the profile; a convenient form of splint is described. Resection of the septum may be needed. Where a portion of the maxilla is impacted in the antrum access is obtained by the canine fossa route. If much damage has been caused to skin and soft parts, every effort must be made to conserve tissue, limit sepsis, and prevent adhesions.

I am indebted to Dr. F. J. A. Mayes for the skiagrams and photographs.

THE ROLE OF THE PITUITARY IN THE ETIOLOGY OF CANCER

[PRELIMINARY PAPER]

BY

WILLIAM SUSMAN, B.A., M.D., C.M.

LECTURER IN PATHOLOGY, MANCHESTER UNIVERSITY; HONORARY
PATHOLOGIST, VICTORIA MEMORIAL JEWISH HOSPITAL

(With Special Plate)

Although many of the functions of the ductless glands remain obscure, it is known that the anterior pituitary concerns itself with promoting growth. However, the prevailing conception of this function of the anterior pituitary is limited to the organism as a whole and not to cell growth, or tissue growth, as seen in repair and disease. Zondek (1930) has shown that in 15 per cent. of 118 cases of malignant disease the anterior pituitary hormone was present in the urine in sufficient quantity to give a positive Zondek-Aschheim reaction in mice. In other words, in neoplastic disease anterior pituitary activity is increased to an appreciable degree in a significant proportion of cases.

That this activity of the anterior pituitary does not in itself cause cancer is evident from the fact that in 98 per cent. of healthy pregnant females the Zondek-Aschheim reaction is positive. The relationship of malignant disease with pregnancy, in that both types of cases may give a positive Zondek-Aschheim reaction, is interesting in two ways. In both there is a tumour present, but the obvious point of difference between the two is that in pregnancy the growth is controlled—a condition lacking in the malignant growth.

GROWTH-RETARDING HORMONE IN POSTERIOR LOBE

What is responsible for this controlling of growth, whether in the case of a foetus, an organ, a tissue, or a cell? A hypothesis suggested itself in the examination of the pituitary from a male, aged 18, showing all the features of renal infantilism, in whom the posterior pituitary was from two to three times as large as the anterior lobe; whereas, in the normal, the posterior lobe is about one-third the size of the anterior. With a posterior lobe about six times as large as the normal, one can expect some positive feature in the body indicative

of this exaggerated function of the part, and in the case referred to the exaggerated feature was in all probability infantilism. This suspicion is, to some degree, confirmed by Sharpey-Schafer, who states that if posterior pituitary is grafted into young tadpoles they subsequently show definite retardation of growth. Can it be that the function of the posterior pituitary is to check growth? This would be in accordance with the older view that the ductless glands (considering the pituitary and the adrenal each as two glands) work in pairs, one giving rise to a stimulating impulse, and the other checking the former by acting in the opposite manner.

Although the pituitary is the seat of the growth-promoting hormone, and, according to the above hypothesis, of the growth-checking hormone, metabolism is almost as important in the growth process. Hence, in malignant disease, the glands concerned in metabolism must also be considered. That the endocrines might be disturbed in cancer seemed probable. Could the course of the disease be influenced by adjusting the workings of any or all of the glands?

HISTOLOGICAL OBSERVATIONS

In order to appreciate fully the character of this investigation it would be as well to state now that the ideas expressed in this work gradually evolved during a histological survey of the whole endocrine system—on the pituitaries, thyroids, and adrenals, of over 200 cases, and on the parathyroids, thymuses and pancreases, of about 100 cases or over. The selection of the material depended entirely on the state of preservation of the tissue, and not on the type of the case. Hence the number of glands from cancer cases was limited.

The following data on the pituitary were based on an examination of 230 glands, of which 30 were from cases of malignant disease—including 13 of glioma cerebri.

Working on Stewart's hypothesis (1921) that the cells of the anterior lobe are merely in different stages of function, and that the acidophilic cells contain the active secretion, I found that the anterior pituitary in the tumour cases did show a degree of activity appreciably higher than in the normal. Histologically, Zondek's conclusion that 15 per cent. of cases of malignant disease give a positive Zondek-Aschheim reaction was confirmed. In many, or perhaps all, cases of malignant disease, then, the anterior pituitary might be considered as overactive.

The Posterior Lobe

The posterior lobe of the hypophysis offered a more difficult task. The belief that the posterior part of this gland was essentially neuroglial was found to be untenable. Preparations stained for neuroglia showed the neuroglia distributed as strands about circular or elliptical masses of tissue, and in well-preserved sections stained by haemalum and eosin the parenchymatous character of the lobules was evident (Fig. 1 on Plate). The posterior pituitary, then, consists of lobules of roundish epithelial cells; these lobules were well vascularized and surrounded by a neuroglial stroma. The colloid of this section of the pituitary was taken to be its effective secretion, and appeared to emanate from the epithelial lobules, from whence it passed through the veins of the posterior pituitary, to the vessels of the pars intermedia, where the secretions of both anterior and posterior lobes intermixed. If, as already suggested, the secretion of the posterior pituitary is concerned with checking growth, then this lobe of the gland should show changes, with an incidence parallel with that of cancer, in the various age groups; while in cancer cases it should show these and corresponding changes in a greater degree and more frequently.

In this study the brown pigment of the posterior pituitary was frequently found associated with, or in,

masses of colloid. Often, too, the associated colloid was of a greyish colour when stained with haemalum and eosin. The pigment tended to migrate to the perivascular channels. In general, the impression was that the brown pigment represented an altered and perhaps ineffective secretion. For purposes of comparison the pigment in each gland was gauged to be scanty, of moderate amount, or abundant; and in order to produce a graph these grades were evaluated as 1, 2, and 3. The arithmetical average for 180 cases is shown in the graph (reproduced below),

plotted against the age groups. The results are of necessity very crude, but the curve is sufficiently characteristic to indicate that this pigment increases quite rapidly up to the age of fifty years, and probably slowly from then onwards. This corresponds with the frequency of cancer.

The graph shows that brown pigment in the posterior pituitary tends to increase with age.

The numbers in each age group are:

Age group	Male	Female	Total
0-9	9	3	12
10-19	10	8	18
20-29	16	8	24
30-39	15	8	23
40-49	23	16	39
50-59	27	9	36
60-69	8	11	19
70 and over	6	3	9

In the thirty glands from tumour cases there were nine which showed a considerable amount of pigment in the posterior pituitary; twenty-one showed enough sclerosis to be recognizable under the low power of the microscope; nine had both sclerosis and pigment present; and in two cases the posterior section was considerably compressed by a large cyst originating in the pars intermedia; in six cases, however, no obvious change could be made out.

Pituitary Changes in Cancer

The histological examination of the pituitary in cancer cases suggested: (1) that the anterior pituitary was overactive and was stimulating growth; and (2) that the lesions in the posterior pituitary probably affected the quality and quantity of the posterior secretion, and by so doing, in cancer cases, the growth-restraining influence was inadequate.

As already indicated the growth mechanism was closely associated with metabolism. The thyroid and parathyroid glands in cancer cases showed no change which could be definitely associated with the disease in question. However, in the pancreas, an important and very significant abnormality was noted.

Changes in the Pancreas

The pancreas was examined in 170 cases, including fourteen cases of cancer apart from carcinoma of the pancreas. The islets in six cases were very numerous, in six numerous, and only in two cases of normal incidence. In all except two cases the islets were enlarged, if the normal islet is taken to be 80 by 110 μ (Fig. 2). The groups of cases were certainly small, but these results can be considered to indicate that in cancer cases there is an increased demand for carbohydrate.

Warburg (1923 and 1924) found that, with the same amount of oxygen, cancerous tissue would utilize more carbohydrate, and produce three to four times as much lactic acid as normal tissue. Handel and Tadenuma (1924) showed experimentally that transplanted rat tumours grew more rapidly when the recipient animals were fed on glucose and injected with insulin. Borrel (1922) states that injections of glycogen had an accelerating effect on experimentally produced tumours. Thus the experimental work of others confirms the view obtained from the histological examination of the pancreas in cancer cases. It might be argued that cancer, in these circumstances, should be common in diabetes mellitus. In this disease, however, although the blood sugar is high, there need not necessarily coexist an overactive anterior and a deficient posterior pituitary.

To summarize briefly: in cancer cases there probably exists an unbalanced state in several of the ductless glands which permits the development of the new growth; the following points appear to play an important part—(1) overactivity of the anterior pituitary; (2) underactivity of the posterior pituitary; and (3) an increased demand for carbohydrate, giving rise to abundant and enlarged islets of Langerhans.

EXPERIMENTAL OBSERVATIONS

The experimental phase of this investigation can be divided into two parts: (a) to determine whether tumour growth is influenced in the presence of abundant carbohydrate, and (b) to determine whether injections of pituitrin would have any effect on checking the growth of epitheliomata in mice.

Effect of Excessive Carbohydrate

One hundred mice were divided into three groups—I of 30, II of 30, and III of 40. All the animals were painted with shale oil five times a week. Group I was fed on oats and cheese; Group II on oats, bread soaked in simple syrup, and simple syrup in the drinking tube. Group III was fed similarly to Group II, but the animals received, in addition, injections of 0.1 c.cm. of a 10 per cent. sterile solution of glucose. These injections varied from three times in the first week to once a fortnight during the first two months. When tumours began to appear, they again received similar injections once a week. At sixteen weeks 30 per cent. of group I, 40 per cent. of Group II, and 56 per cent. of Group III had developed warts. By the twentieth week the first group had 59 per cent., the second group 66 per cent., and the third group 100 per cent. showing tumours. In all, at the thirty-eighth week eighteen epitheliomata had been produced, and of these two occurred in Group I, eight in Group II, and eight in Group III. The incidence of warts in the various groups is not very significant, because the number of animals was too small; but, in view of the results of other workers, these data can be considered as suggestive. The incidence of epitheliomata is much more significant, and appears to show quite definitely that the presence of an abundant supply of glucose does promote the development of epitheliomata in mice.

Effect of Pituitrin

As the epitheliomatous mice became available they were segregated, and the oil applications ceased. In all, thirteen of the eighteen epitheliomata were clinically evident, while the remaining five were recognized only after microscopical examination. Of the group of thirteen, five were used as controls, while the remaining eight were treated. The five controls had the tumours of lesser severity, while the eight animals that were to receive treatment all had severe, extensive, and very obvious epitheliomata. The purpose of the controls was to find

out if any natural regression occurred in the tumours on a basis similar to that which took place in the treated animals. In only one of the control animals did the tumour drop off, and that occurred in a pregnant mouse; but with the termination of pregnancy the growth returned very rapidly.

In the group for treatment, 0.1 to 0.3 c.cm. of a solution of pituitrin (P., D. and Co.) (diluted to 0.1 c.cm. in 35 c.cm. of sterile normal saline) was injected twice daily. The injections were made subcutaneously over the abdomen, or at some point distant to the tumour. The general reactions noted were as follows. Within three days the growth contracted noticeably and became covered with a scab. Gradually the growth became sharply demarcated from the skin. In ten to fourteen days the base became prominent and fluctuant, and from fourteen days onwards necrotic material could be expressed from this region. From a month to six weeks or so, the growth separated, leaving a simple ulcer (Fig. 3), in some cases a few malignant fragments, and in others a thin epitheliomatous layer at the base. Most of the animals died through the wound having become infected, or because of the presence in the body of large numbers of encysted parasites.

EXPERIMENTAL CONCLUSIONS

The ultimate results were: (1) complete disappearance of the epithelioma—two cases, four weeks and eight weeks; (2) one or several minute epitheliomatous nodules found on histological examination—four cases, three weeks, five weeks, six weeks, and six weeks; (3) growth reduced to a thin layer at base—two cases, four weeks and nine weeks. It was thus shown, experimentally, that, in mice, the presence of glucose in abundance has a stimulating effect in producing epitheliomata, and that, within limits, pituitrin has a checking influence on malignant tumours. The experimental field was obviously limited; the results obtained, however, were in accordance with the theory enunciated on the basis of the histological findings. Consequently the application of the theory in the therapy of human cases was justifiable. The investigation in its subsequent stage was therefore of a clinical nature.

REMARKS ON THE FIRST GROUP

Up to the present stage no means were available by which the anterior pituitary could be controlled, and hence the only therapeutic means available were: (1) a diet low in carbohydrate to starve the tumour, and (2) pituitrin (P., D. and Co.) to check the malignant growth. The cases in this group were four in number, and when treatment commenced the average expectation of life was only a few weeks at most, for the growths were all very advanced.

CASE I

Male, aged 49. Primary carcinoma of tongue with bilateral involvement of submaxillary gland and extensive infiltration into the neck on both sides. There was a large tumour-like swelling on both sides of the chin, and in each tumour there was a small sinus. General condition was fair and cachexia moderate.

Treatment.—Low carbohydrate diet; 0.5 c.cm. to 2 c.cm. of pituitrin (P., D. and Co.) twice daily. At the end of three weeks the patient died. After the first week the discharge from both sinuses progressively increased, and the sinuses enlarged. After the second week, the submaxillary tumours had visibly diminished in size. At necropsy no growth was found in the tongue, while the submaxillary growths were soft, crumbly, and necrotic; but firm tumour tissue was present in the peripheral shell (Fig. 4). A mediastinal secondary was in a similar state. The cause of death was fatty infiltration of the heart and cirrhosis of the liver.

CASE II

Female, aged 59. Epithelioma of right cheek about 12 cm. in diameter, of the cauliflower type, and with an extension down the lateral side of the neck. General condition bad. Cachexia advanced. Treatment similar. After five and a half weeks of treatment, this patient died in a highly toxic state. At necropsy the cauliflower growth had been reduced to a nodule 2 cm. in diameter, but the original base was still malignant. The mass on the side of the neck was hollow and contained clear fluid. Extending below and behind the right eye there was a mass of necrotic tumour tissue. Cause of death—fatty degeneration of the myocardium.

CASE III

Female, aged 61. Recurrence of carcinoma of the left breast in surgical scar and in adjacent area, after radical amputation. The area involved was 12 cm. by 18 cm. over the lateral and anterior aspect of the left breast. Condition poor, cachexia moderate. Treatment similar for the first six and a half weeks. At six weeks the malignant edge had begun to disappear, and the centre of the wound seemed to be clearing of growth. The involved area became irregular through some healing having occurred from the sides.

CASE IV

Female, aged 44. Carcinoma of both breasts. Both breasts were very prominent, and growth had ulcerated through the right breast in the region of the nipple. Condition poor. Cachexia advanced. Malignant growth advanced. Treatment similar for the first six and a half weeks. During the first week the area about both breasts became intensely congested and denuded of skin. The necrosis in the right breast progressively increased, and at three weeks both breasts appeared to have decreased in size; at one month after treatment the left breast was no longer fixed.

Commentary

In all the four cases in this group there was intense pain in the tumour three to ten minutes after each injection, which lasted for about half an hour. This began at the growing margin and spread over the whole tumour. In all cases, too, the diastolic pressure increased, leaving the pulse pressure at twenty minutes after an injection as low as from 16 to 20. This, however, was transient. More troublesome was the small amount of urine which was periodically recorded, but the amount returned to normal when the dose of pituitrin was reduced, or, if necessary, omitted for that day. Signs of an overdose of pituitrin occurred occasionally, and in the form of spasm of the occipital, brachial, or femoral arteries. One case showed dilatation of the pupils, and at another time the systolic pressure rose by 100 points. Fortunately, these were all transient.

During the six and a half weeks included in this period, when the treatment consisted only of injections of pituitrin and of a diet low in carbohydrate, all the growths showed definite regression. In all cases, however, this regression appeared to come to a standstill, and for a time it seemed that this line of approach to the treatment of malignant disease would be of a very limited character. It was at this point that the Zondek-Aschheim reaction in its relationship to malignant disease was considered, and by this means a method for restraining the activity of the anterior pituitary presented itself.

REMARKS ON THE SECOND GROUP

Zondek showed that in 15 per cent. of 118 cases of malignant disease the Zondek-Aschheim reaction was positive. The unpublished results of eighteen cases investigated by Susman and Nuttall showed that fourteen gave a positive reaction; these included four of the six cases in both groups. The remaining two have not been tested. In short, it has been a working rule to suppose

that every case of cancer has a hyperactive anterior pituitary, for it was found that several of the fourteen cases which had previously given a negative reaction became positive if the test was repeated several times. Case II presented a further interesting point. The patient was obviously in a highly toxic state. Her urine, when injected into mice, was lethal. When the urine was diluted twenty times the Zondek-Aschheim reaction was very definitely positive. In other words, the reacting hormone was present in very great amount in the undiluted urine, and it was just possible that it was the excess of anterior pituitary secretion which caused the profound toxæmia. Zondek (1930) found that, in cases in which castration had been performed by operation, 75 per cent. gave a positive reaction, and this included bilateral oöphorectomy. This suggested that the genital glands had a restraining influence on the anterior pituitary, and that perhaps, by the administration of an ovarian extract (theelin, P., D. and Co.), the overactivity of the anterior pituitary in malignant disease might be controlled. The therapeutic agents utilized in Group II were, then: (a) a diet low in carbohydrate, to starve the growth; (b) pituitrin (P., D. and Co.), to reinforce the secretion of the posterior pituitary; and (c) theelin (P., D. and Co.), to restrain the overactivity of the anterior pituitary.

CASE III

Treatment continued as in Group I with the addition of 0.25 c.cm. of theelin daily for three weeks, and then once or twice weekly. The malignant area continued to clear. In two months the central area was largely cleared, but there probably remained a thin base of malignant tissue. In three months the malignant edge was no longer apparent, and the surface was flat and clean. The appearance was good, but the area was probably not entirely free from growth.

CASE IV

Treatment as in Group I, with the addition of 0.25 c.cm. to 0.5 c.cm. theelin once daily. Necrosis of the right breast continued, and by the tenth week the right breast was reduced to a layer of tumour tissue on the chest wall. The left had become much reduced in size; progress continues.

CASE V

Male, aged 55. Epithelioma of tongue with involvement of floor of the mouth and left submaxillary gland. Tongue was fixed, and patient was unable to speak.

Treatment.—Diet low in carbohydrate, pituitrin 1 c.cm. to 2 c.cm. twice daily, and theelin 0.25 c.cm. to 0.5 c.cm. daily. Within two days his speech was understandable. At the end of one month the tongue was cleaner, not so large, and slightly mobile. Pain in the tongue was severe after injections. The patient looked well and felt well.

CASE VI

Female, aged 76. Slow-growing epithelioma (Fig. 5) on the dorsum of the foot, 2.6 cm. by 3 cm., with an actively infiltrating edge on the right side and hyperkeratosis on the left (Fig. 7).

Treatment.—Diet low in carbohydrate. Pituitrin from 0.5 c.cm. to 2 c.cm. twice daily, and theelin from 0.25 c.cm. to 0.5 c.cm. daily, with occasional lapses for several days. For the first two weeks theelin was not available.

In five days the advancing edge disappeared. At two weeks the tumour was sharply demarcated from the skin, and the crescent of tumour tissue at the upper pole was necrotic (Fig. 8). Sections at this stage showed that the tumour was well differentiated from the epidermis (Fig. 6), and was being surrounded by cellular connective tissue. The tumour became very congested, and much necrotic material came away in the dressings. At the end of one month the tumour edge was separating, the skin was growing in underneath, and the crescent at the upper pole had disappeared. Since treatment began the tumour was no longer painful. At six weeks a probe could be passed in under the growth over a considerable part of its circumference. At the end of seven weeks the tumour was removed completely and easily by enucleation

with a very blunt instrument. When the tumour was removed there could be seen a central area where the tumour had been attached, about 2 cm. in diameter, and an outer rim of skin, about 0.4 cm. in width, signifying that separation had occurred to the extent of about 0.8 cm. of the diameter of the growth before the operation. Since the operation the patient has been receiving 0.5 c.cm. to 1 c.cm. of pituitrin daily, with occasional injections of 0.25 c.cm. theelin. Three weeks after the tumour had been enucleated the wound had healed completely, and the patient was discharged from hospital. For the last month the patient looked very well, and felt much younger.

CASE VII

Female, aged 54. Carcinoma right bronchus. Before treatment she complained of a constricting feeling about the trachea, and had pleural effusion, which caused dyspnoea.

Treatment.—Diet low in carbohydrate, 0.5 c.cm. to 1 c.cm. pituitrin twice daily, and 0.25 c.cm. to 0.5 c.cm. theelin once daily. In fourteen days the constricting feeling about the trachea disappeared. She looked and felt quite well, but respirations continued between 35 and 40. This would be largely due to the intense congestion induced in the affected lung by the treatment. Pain occurred in the right chest after each injection.

DISCUSSION

The crucial test for this form of treatment was in Case VI. There we find that the growing edge disappeared in five days, that the tumour began to separate at fourteen days (at which time the associated hyperkeratosis had disappeared), that in six weeks a probe could be inserted some distance under the growth at most points along the circumference of the tumour, and that at seven weeks the tumour, an epithelioma proved by microscopical examination, was successfully enucleated without any obvious cutting. This case alone showed that the treatment, based on a definite hypothesis, was successful. There is nothing, however, to indicate that recurrences may not happen at some time or other. It is hoped that the patient will continue receiving periodic injections of pituitrin and theelin, for at present there is no way of establishing the fact that the normal endocrine balance has been restored.

As for the other cases, all the tumours appear to show some regression; life has been prolonged; and all the patients look and feel better than before. After theelin was employed there has been no serious set-back in the progress of the cases and no deaths.

That pituitrin can affect tumour growth was shown by Norgate (1921) when he treated thirty-six cases with injections up to 3 c.cm. a week, but these injections were made into the tumours. He found that it caused an improvement in appetite, in weight, and regression of the tumour. Secondaries were rare. On the whole, life was prolonged for a year or more. He concluded that pituitrin only delayed growth by cutting off the blood supply. Carse (1918) injected pituitrin in cancer cases to raise the blood pressure, and he found a noticeable improvement in the cases so treated. Neither of these inquiries was based on a belief that the ductless glands might play an important part in cancer.

The pain which followed after the injection of pituitrin in cancer cases was investigated by Reding and Slosse (1926), who found that this phenomenon occurred in 80 per cent. of eighty cases. No pain was complained of by patients suffering from a non-malignant ulcer, tuberculosis, or syphilis. They concluded that the pain was not due to vaso-constriction, and that this phenomenon was neither restricted to any particular type of growth nor dependent on the amount of smooth muscle in the part.

In conclusion, only a few more points need be dealt with. Pain was controlled by morphine or its derivatives. When the output of urine came below 20 ounces the dose of pituitrin was reduced, and if necessary stopped. The output of urine invariably returned to normal in a day or

two. The low pulse pressure which frequently occurred after an injection of pituitrin merely caused a transient faintness. Finally, the effectiveness of the dose was gauged by the temperature, for when the temperature varied between 97° and 99° it was judged that the regressive changes were sufficient. When toxic absorption caused the temperature to rise above 99°, the usual measures were employed to bring it down.

SUMMARY

1. Based on the hypothesis that in cancer cases the anterior pituitary was overactive, and the posterior pituitary underactive, two cases of advanced cancer were treated with pituitrin alone, and five cases with pituitrin and theelin. All were placed on a diet low in carbohydrate.

2. An epithelioma began to separate off and was enucleated after seven weeks of this treatment. The growing edge disappeared in five days.

3. All the cases showed regressive changes in the tumours, and life appeared to be definitely prolonged.

I wish to thank the honorary, medical, and nursing staff of the Christie Cancer Hospital, and Dr. Stanley White of Messrs. Parke, Davis and Co., for their valuable co-operation in the clinical work. I am indebted to the Manchester Committee on Cancer, and the Manchester University Grants Committee for grants to defray the expenses of this investigation. For the photographs and photomicrographs I have to thank Mr. H. C. Taylor of the pathological department.

BIBLIOGRAPHY

- Borrel, A., and de Coulon, A.: *Compt. rend. Soc. de Biol.*, 1922, lxxxvi, 1096.
 Carse, G.: *Med. World*, 1918, x, No. 21.
 Hande, M., and Tadenuma, K.: *Zeit. f. Krebsforsch.*, 1924, xxi, 288.
 Norgate, J. H.: *Brit. Journ. Surg.*, 1921, ix, 495.
 Reding, R., and Slosse, A.: *Compt. rend. Soc. de Biol.*, 1926, xcv, 1442.
 Sharpey-Schafer, F.: *The Endocrine Organs*, Part II, p. 250.
 Stewart, F. W.: *Compt. rend. Soc. de Biol.*, 1921, lxxxiv, 49.
 Warburg, O.: *Biochem. Zeit.*, 1923, cxlii, 317, 334; *Ibid.*, 1924, cxlvi, 309; *Klin. Woch.*, 1924, iii, 1062.
 Zondek, B.: *Klin. Woch.*, 1930, ix, 679.

TORSION OF OVARIAN CYST IN CHILDREN

BY

W. ARKLEY STEEL, M.D.St. AND., M.R.C.P.LOND.

REDHILL HOSPITAL, EDGWARE

(With Special Plate)

The age incidence of ovarian tumours has been noted in several valuable series of recorded cases. Statistics indicate the occurrence of ovarian tumours at the extremes of age and their comparative infrequency in childhood. The proportion of cases met with in children does not exceed 3 per cent. Illustrative of the early age at which ovarian tumours may appear is the case reported by Doran of an ovarian tumour in a premature child, aged 7 months. Another instance is given by Chiene, who performed ovariectomy on a child aged 3 months. The majority, however, of these tumours in children are met with between the ages of 10 and 14. Both the solid and the cystic types of tumour of the ovary are found in children. Of the simple variety, cystadenomata and dermoid cysts are equally common, and constitute about two-thirds of all tumours. The malignant growths form the remainder, and, of these, sarcomata are rather more common than carcinomata.

Cases of ovarian tumours are brought to our notice by reason of: (1) their increasing growth giving rise to abdominal deformity; (2) constitutional disturbances; (3) mechanical effects, causing pressure on the surrounding viscera; (4) complications. In children the complications of rupture, or of suppurative of an ovarian cyst,

are exceedingly rare, and torsion, although the commonest complication, is relatively uncommon. The rarity of the condition justifies the detailed description of the following case, which was under my care.

ILLUSTRATIVE CASE

J. L. M. J., female, aged 9 years, was admitted to hospital on October 13th, 1930, as a surgical emergency. The parents were both healthy, but somewhat dull intellectually. There was one other sister alive and well.

Previous History.—The patient was born after a short and normal labour, and, except for measles and whooping-cough, she had always been very healthy. When about 5 years of age she had an attack of pain, which commenced on the left side and extended across the upper part of the abdomen. This was accompanied by retching, and "trouble with the bowels." These attacks recurred each year, but were never severe enough to confine the child to bed. The mother was emphatic that the onset of each attack synchronized with the approach of the colder weather, taking place usually in October. The pain lasted as a rule for fourteen days, but eventually responded to doses of castor oil. She had had attacks of vague abdominal pain frequently throughout the last year.

Present Illness.—At 9 p.m., on October 5th, 1930, the patient had a sudden attack of vomiting. There was no pain at first, but the next day she complained of pain in the left side of the abdomen. This pain extended across the upper part of the abdomen, and in no way differed from that experienced in previous attacks. She attended school from October 7th to 10th, but at 11 a.m. on the latter date she had a sudden attack of pain, which "doubled her up." The pain started in the left side, and passed down to the left groin and across the abdomen. She vomited several times. None of the measures which had been successful in previous attacks relieved the pain. The child had a peculiar taste for castor oil, and of her own accord frequently took copious draughts straight from the bottle. There was no action of the bowels from October 9th until the 13th, when she passed a normal motion. Micturition was normal up to the day of admission, when she complained of burning pain on passing urine. The abdominal pain continued, and her general condition became gradually worse. She was sent to hospital as a case of acute appendicitis.

On later interrogation, the mother said that for the past three years she had noticed her child's abdomen becoming more prominent, and that it "seemed to be getting hard on the left side." The patient often acted in a peculiar manner, and the mother was inclined to regard her as somewhat "simple-minded." She had never menstruated, nor was there any precocious sexual development.

Examination.—On admission the patient was extremely ill and dehydrated, crying out from time to time owing to spasms of pain. The temperature was 100.8° F., pulse rate 124, respirations 24. The tongue was furred and dry. The upper part of the abdomen moved slightly with respiration, the lower part not at all. A globular swelling, extending from the pubis to the umbilicus, was apparent, filling the hypogastrium and extending to the left side. Palpation confirmed the presence of this swelling, which was felt to be tense, tender, and cystic. It appeared to rise from the pelvis, and was rather more prominent on the left side. It was so centrally placed, however, that the possibility of its being bladder was considered. Catheterization was performed, but had no effect in diminishing the size of the tumour. There was generalized tenderness over the abdomen and in the loins. Rigidity was present over the whole abdomen, but was most marked in the muscles overlying the tumour, especially on the left side. On rectal examination a hard, tense, tender, and cystic swelling was felt filling the pelvis, and so bulging the anterior wall of the rectum as practically to occlude the lumen of the bowel. A slight degree of vulvitis was present, with a scanty yellowish discharge. The examination of the urine showed the presence of ketones, but was otherwise normal. The other systems revealed nothing abnormal. The diagnosis of acute appendicitis was excluded. The condition was thought to be an abscess, circumscribed in the pelvis, due possibly to an extension from the vulvitis. An operation was decided upon, and was performed at 11:50 p.m. on October 13th.

WILLIAM SUSMAN: ROLE OF THE PITUITARY IN THE ETIOLOGY OF CANCER

FIG. 1.—Posterior pituitary—haemalum and eosin. The photograph shows a definite parenchymatous portion with a fairly pronounced stroma. The lobule is of an epithelial character, and the stroma glial. In the lower right-hand corner a glial sclerosis is evident. $\times 100$.

FIG. 2.—Abundant and enlarged islets in a case of carcinoma of rectum. $\times 50$.

FIG. 3.—A simple ulcer on the back of a mouse after the epithelioma had come away a month after treatment. Section at two months. $\times 50$.

FIG. 4.—Case I. The growth had become hyalinized, and was separating from the thin floor of malignant tissue. $\times 60$.

WILLIAM SUSMAN: ROLE OF THE PITUITARY IN THE ETIOLOGY OF CANCER

FIG. 5.—Case VI. Histological features of the epithelioma of the foot before treatment. $\times 70$.

FIG. 6.—Epidermal layer has returned to the normal thickness, and the tumour is sharply differentiated from the epidermis. The deeper portion of the growth was bounded by cellular connective tissue. Section taken after fourteen days' treatment. $\times 80$.

FIG. 7.—Case VI. Macroscopic features of the foot and tumour before treatment. Note the hyperkeratosis on the left and the invading margin on the right of the tumour.

FIG. 8.—Case VI. After fourteen days' treatment the keratosis had disappeared and the tumour is sharply outlined. The advancing edge had disappeared after five days.