personal, familial, and community hygiene; and what vistas of possibility are opened up by Professor Mellanby's lecture on the prevention of disease by the correct feeding of children. It is obvious that we must not try to escape infection for our children, for, dangerous though infection is, it does bring with it protection, and we must avoid the state of affairs that Heimbeck deplores in his own country in these words:

"Thanks to tuberculosis work, infection in the home is not so common and is postponed until the individual goes out to work. Thus the endeavour to avoid tuberculosis infection in childhood has resulted in the transference of a high morbidity and mortality to just those years when the individual should begin to work for society in return for what society has done for him in the matter of education in his earlier years."

Really, it is Nature's problem. She will find her own solution, and I think she is solving it, as Dudley Sheldon suggests when he says:

"Morbidity equals infection pressure versus herd immunity. Herd immunity increases with the herd's past experience of the bacterial cause of infectious disease, and although infection pressure must be greater with the increase of urbanization, herd immunity outstrips it."

THE BROKEN NOSE*

BY

E. WATSON-WILLIAMS, M.C., CH.M.

SURGEON IN CHARGE, EAR, NOSE, AND THROAT DEPARTMENT,
BRISTOL ROYAL INFIRMARY

(With Special Plate)

A broken nose is common enough and regarded often, except perhaps by the victim, as matter for mirth rather than for sympathy. Fortunately, even in these motoring days, the majority of our cases are of a mild type: when there is no subsequent external deformity, any distortion of the nasal septum can be readily dealt with by sub-mucous resection. Omitting those in which there was no manifest displacement of the nasal bones, the etiology in twenty-seven cases treated personally during the four months March to June, 1931, was as follows:

	Mild	Severe
Football	3	—
Cricket	4	—
Boxing, etc.	1	1
Motoring	2	4
Cycling	1	1*
Falls, etc.	3	1
No record	6	—
	20	7

* Collision with motor lorry

But the deformity resulting even in the milder cases may be very annoying—to a girl possibly really distressing. If it is to be avoided it is important that treatment should be undertaken *early*, since the processes of repair lead to rapid absorption of bone edges, and callus formation may produce an unsightly bossing along the margin of the frontal process of the maxilla; so that a good result is often not obtained by late treatment, even though the bones be re-fractured and set. Still more important is it that treatment should be intelligent—that is, guided by a proper understanding of the exact nature of the deformity, and the best method of reduction. Though it is often easy to see the line of fracture in an ordinary skiagram, the actual displacement in a mild case is not well shown unless resort is had to a stereogram. Gross damage is, of course, sufficiently obvious in any good film (Figs. 1 and 2 on Plate).

* A Bristol University Post-Graduate Lecture.

In a typical mild case the accident is followed immediately by epistaxis, and very soon by nasal swelling, internal and external, probably with a black eye in addition; it is impossible in these circumstances to estimate damage. Local treatment is confined to suture of lacerations, etc. In four to eight days the swelling has gone down, nasal obstruction is slight or absent, and one can make out the characteristic deformity (Fig. 3 on Plate). The bony nose is bent definitely over to one side, the septum and tip being little affected. If now one attempts simply to push the nose back into place, in all but the most trivial cases this is found unexpectedly troublesome, and a symmetrical appearance is not achieved (Fig. 11). Also, the deformity tends obstinately to recur; internal splints effect nothing, while external splints or padding are often impossible to retain in place. If the true mechanism of the fracture-dislocation is appreciated, reduction is perfectly simple, and one can almost guarantee that there will be no subsequent deformity.

The actual displacement of bone presents three features (Figs. 9 and 10).

1. On the side from which the blow fell the lower two-thirds of the nasal bone, and often the anterior sharp margin of the frontal process of the maxilla, are displaced inward toward the mid-line, mainly by rotation round the posterior margin of the fragment as a hinge, with very little backward displacement.

2. On the opposite side a corresponding piece of bone, or rather less, is displaced in the same direction—that is, outward—again hinging on its posterior margin.

3. In the middle the first fragment has been driven in beneath the second. It is lack of appreciation of this imbrication-impaction that is the source of difficulty and failure in reduction. Figs. 9 to 13 are diagrammatic sections at right angles to the plane of the nasal profile, to explain the common types of displacement and the steps in reduction.

TREATMENT

A general anaesthetic is necessary, and oxygen-ether will be found convenient. Bleeding is usually slight; if desired, the nose may be lightly sprayed just before induction with 5 per cent. cocaine solution and packed anteriorly with gauze wrung out of 1 in 10,000 adrenaline. When anaesthesia is complete a post-nasal sponge can be introduced. The only instrument needed is a strong forceps, the blades 2 to 2½ inches long and slightly bowed so that only the tips touch. Jansen's longest bone forceps may be used; the common septum forceps is too thick for most noses. Before use the tips are well padded by winding on three or four turns of half-inch strapping.

Operative Procedure

A.—Into the nostril of the side *toward which* the nose is bent pass one blade of the forceps, pushing it up until the other blade outside the nose is just over the nasal bone. Grasp the bone firmly and twist it yet further outward, still round the posterior margin as an axis (Fig. 12). (This step is not always essential, but greatly facilitates disimpaction.)

B.—In exactly the same manner grasp the other or in-driven bone, and rotate it also well outward to a position corresponding with the first (Fig. 13). Sometimes a slight lifting movement is required in addition.

C.—A finger placed externally can now feel that both bones are freely movable, and that there is a definite gap between them anteriorly. With the finger mould the two bones symmetrically together, like the leaves of a double drawbridge, so that they lock in the mid-line. If necessary, the forceps may be used with one blade up each

nostril during this process to support the bones or to correct any anterior deviation of the septum. Once locked there is no tendency to displacement, and indeed considerable force may be needed for this; no splints are required, and the patient can return home forthwith. If during the ensuing week he thinks the nose is not quite symmetrical he may massage it forcibly into place with the thumb twice daily: this is rarely necessary. In this type of case there is seldom much alteration in the profile, and bilateral symmetry supplies the best test of treatment.

CASE I

Nose broken at football, leading to typical deformity of the simple type (Fig. 3 on Plate). The second photograph (Fig. 4 on Plate) shows the result four days after reduction by the method described, no splints being required.

When the violence has been great the problem is frequently less simple. In addition to laceration of the soft parts the bones may be comminuted and driven backward; the whole bridge of the nose may be knocked flat

upright, bend each leg directly backward about half an inch above bend P (Fig. 16). Bend R: at R, two inches back from Q, or somewhat less, bend the wire up and then forward parallel and close to itself (Fig. 17). Bend S: at S, a point slightly in front of Q, bend the wire down at right angles, shaping it to the contour of the nasal profile in the region where support is wanted. Bend T: bend the wire backward at a point that will correspond to the upper part of the vestibule, and then again downward at T (Fig. 18). The distance TR = the distance QR; ST is equal in length to the distance from the upper end of the fracture to the lower margin of the nasal aperture.

The splint is now ready for insertion. Pass the portion QRS into the nose, one "leg" on each side of the septum, until the central part PQ presses firmly against the philtrum. With septum or other forceps hold the part QR in close contact with the floor of the nose, and thrust the portion ST directly upward in the direction indicated by the arrow (Fig. 18); in doing this

TRANSVERSE SECTIONS THROUGH NASAL BONES AT RIGHT ANGLES TO LINE OF PROFILE

FIG. 9.—Typical "mild" deformity.

FIG. 10.—Another common type of fracture.

FIG. 11.—Result of replacement without disimpaction.

FIG. 12.—First movement of disimpaction.

FIG. 13.—Second movement of disimpaction.

FIG. 14.—Crushing of right side of nose, as in Case III.

FIG. 15.—Heavy crushing blow involving antrum, as in Cases IV and V.

(Fig. 15). Further, the nasal septum is broken or bent, and occlusion of one or both passages common. Such accidents often lead to serious late deformity, calling for special reparative operative procedures; as, for example, by cartilage grafting, one method of which was described in my article in the *British Medical Journal* of November 28th, 1925.

The need for such measures can be largely avoided by timely reposition; a photograph showing the normal profile, if available, is of considerable value. In dealing with the external deformity one is not troubled by impaction. The fragments are only too mobile; with the forceps they are readily restored to the normal position, but can seldom be retained without support. For this purpose the ordinary nasal splints leave much to be desired; they are difficult to fit, more difficult to fix, and if large enough to be effective still more difficult to withdraw.

To meet this situation I have devised a wire splint which supports the bridge of the nose by counter-pressure on the nasal floor. It is prepared as follows. With narrow rubber tubing, such as that used in the Carrel technique, cover about 16 inches of malleable silver wire of 1/20 inch diameter (= 18 s.w.g.) Bend P: bend the wire in the centre so that it forms a long narrow U, the legs about 1/4 inch apart. Bend Q: holding the U

the portion below S is brought up beneath the depressed part of the nose, and the angle QRS is opened (Figs. 19 and 20). Make sure that the splint really does support the front of the nose when the forceps are withdrawn, that it is securely fixed, and that the nose is symmetrical. Tie the splint below T to the portion between P and Q which rests on the philtrum. Turn the free end outward round the base of the ala nasi to prevent upward displacement (Fig. 21), and cut off any portion that is redundant.

A piece of strapping may be used across the upper lip and projecting part of the splint if additional security is desired. But the splint is genuinely self-retaining, is well tolerated for eight to ten days, does not get foul, and permits of nasal lavage and at least fair nasal respiration. When removal is desired the steps just described are reversed. In these cases, provided antero-posterior reposition is good, there is seldom much difficulty in securing bilateral symmetry; the profile affords the best criterion of success or failure.

CASE II

This patient was attacked by a man armed with a knuckle-duster or a metal bar, and the bridge of his nose was beaten flat. The triangle splint was used for seven days, with the satisfactory result shown (Fig. 5 on Plate).

CASE III

A girl was involved in a collision between her cycle and a motor lorry. The deformity resembled the simple type, but skiagraphic examination showed that the right nasal bone had been crushed in against the septum, and the frontal process of the maxilla fractured off, though the left side of the nose was comparatively little affected (Fig. 1 on Plate; compare Fig. 14). The triangle splint was employed for seven days and a perfect restoration achieved.

The nasal septum can sometimes be reduced into place by forceps pressure and manipulation. Should this fail, it is best at once to resect the damaged portion by the submucous route; for the profile of the nose beneath the nasal bones (except that of the tip) is supported by the anterior edge of the quadrilateral cartilage; and while that is distorted, restoration of symmetry may prove impossible.

Another complication of severe crushing is fracture of the body of the maxilla, the line of cleavage often running from the orbital margin at the maxillary-malar suture

supports; the septal cartilage was resected. The cosmetic result was perfect.

CASE V

This lady fell from a ladder, and her nose, coming in contact with the edge of a dresser, was crushed level with the cheeks. A skiagram showed that in addition to comminution of the nasal bones and fracture of the vertical plate of the ethmoid a large piece of the maxilla had been driven into the antrum. In order to reduce the latter it was necessary to approach it via the canine fossa route, to cut off an angle that defied all efforts at reduction, and then to lever it up with a hook passed through the oral incision and an elevator introduced up the nostril. The nose was then restored, and the triangle splint used for ten days. A perfectly straight profile was obtained, the symmetry being well shown in the photograph. (Fig. 6 on Plate.)

Finally, in the most serious cases there may be extensive damage to skin and mucous membrane, or the nose may be almost torn off. Where considerable loss of substance occurs one may have later to undertake elaborate reparative operations—for example, the now well-known tube-graft rhinoplasty that we owe to Sir Harold Gillies. In

THE SELF-RETAINING TRIANGLE NASAL SPLINT

FIG. 16.—Bends P and Q.

FIG. 17.—Bend R, right side only.

FIG. 19.—The splint expanded and tied.

FIG. 20.—Section through left nasal chamber, showing position of splint.

FIG. 18.—The splint ready for insertion.

FIG. 21.—The upper lip with splint in position.

downward, inward, and backward (Fig. 15). The large fragment thus detached may become impacted in the antrum which the fracture has opened, producing by distortion of the orbital margin a most peculiar appearance. It may be possible to lever the fragment up into place by passing a blunt instrument up the nostril, taking care that this does not slip upward toward the roof of the nose. If that fails, convenient access may be obtained by an incision in the mucous membrane of the canine fossa, reflecting the periosteum as if for a Scanes-Spicer antrostomy. Where an angle or spike of bone prevents restoration it may have to be clipped off with bone-cutting forceps.

CASE IV (FIG. 2 ON PLATE)

Miss L. was involved in a serious motor accident. I saw her ten days later and found the whole nose had been pushed over to the left, only the tip remaining in the mid-line. The left nostril was completely blocked by the crumpled septal cartilage. The right side of the nose and the adjacent part of the cheek were markedly depressed, with a manifest gap in the orbital margin; the cheek and upper lip were anaesthetic. The skiagram shows the extensive damage to the maxilla, and the opacity produced by blood in the right antrum. Under general anaesthesia the large fragment was levered up into place, where it remained while the second large fragment was elevated; the nasal bones were replaced, and the whole nose now remained firmly set without

order to minimize the task of cosmetic restoration we must in the early treatment be guided by two principles. First, preserve every scrap of viable tissue, especially skin and mucous membrane. Secondly, close all wounds, even stitching skin to mucous membrane when necessary for this end, so that healing by granulation with its attendant cicatricial contraction is reduced to the unavoidable minimum. Within the nose we must prevent adhesions between the sides and the septum. A fairly stiff non-absorbent splint is needed, thin enough to be retained without discomfort for days. It can be improvised from a sheet of celluloid, obtained by soaking off the coating of an x-ray film in hot water; the thicker celluloid sold for repair of car windows is even better. A very useful splint can be made by folding a piece of strapping "sticky sides in" and dipping in melted hard paraffin wax—for example, ambrine or a good quality candle. It is worth some trouble in the initial stages to avoid extensive intranasal adhesions. At the earliest possible moment such cases should be brought under the care of a surgeon with experience of rhinoplastic technique.

CASE VI

This boy was thrown from his motor cycle and brought in unconscious. His nose was crushed, and split in the mid-line, so that the wound gave a good view of the interior. Local

measures were directed to careful suture of skin and mucous membrane, and prevention of adhesions. One week later he presented the appearance shown (Fig. 7 on Plate); even in this case the tip of the nose remains central. The fragments of the nasal bones, maxillae, and ethmoid were carefully manipulated into place under general anaesthesia and retained on the triangle splint. The nasal airway is perfect, and the cosmetic result impaired only by the scar (Fig. 8 on Plate).

SUMMARY

The common type of "broken nose" is due to lateral (but not backward) displacement of both nasal bones with imbrication. Treatment consists of (a) disimpaction, (b) free mobilization of both bones, and (c) symmetrical moulding together. No splints are needed.

When the nose has been crushed, internal splinting is required to restore the profile; a convenient form of splint is described. Resection of the septum may be needed. Where a portion of the maxilla is impacted in the antrum access is obtained by the canine fossa route. If much damage has been caused to skin and soft parts, every effort must be made to conserve tissue, limit sepsis, and prevent adhesions.

I am indebted to Dr. F. J. A. Mayes for the skiagrams and photographs.

THE ROLE OF THE PITUITARY IN THE ETIOLOGY OF CANCER

[PRELIMINARY PAPER]

BY

WILLIAM SUSMAN, B.A., M.D., C.M.

LECTURER IN PATHOLOGY, MANCHESTER UNIVERSITY; HONORARY
PATHOLOGIST, VICTORIA MEMORIAL JEWISH HOSPITAL

(With Special Plate)

Although many of the functions of the ductless glands remain obscure, it is known that the anterior pituitary concerns itself with promoting growth. However, the prevailing conception of this function of the anterior pituitary is limited to the organism as a whole and not to cell growth, or tissue growth, as seen in repair and disease. Zondek (1930) has shown that in 15 per cent. of 118 cases of malignant disease the anterior pituitary hormone was present in the urine in sufficient quantity to give a positive Zondek-Aschheim reaction in mice. In other words, in neoplastic disease anterior pituitary activity is increased to an appreciable degree in a significant proportion of cases.

That this activity of the anterior pituitary does not in itself cause cancer is evident from the fact that in 98 per cent. of healthy pregnant females the Zondek-Aschheim reaction is positive. The relationship of malignant disease with pregnancy, in that both types of cases may give a positive Zondek-Aschheim reaction, is interesting in two ways. In both there is a tumour present, but the obvious point of difference between the two is that in pregnancy the growth is controlled—a condition lacking in the malignant growth.

GROWTH-RETARDING HORMONE IN POSTERIOR LOBE

What is responsible for this controlling of growth, whether in the case of a foetus, an organ, a tissue, or a cell? A hypothesis suggested itself in the examination of the pituitary from a male, aged 18, showing all the features of renal infantilism, in whom the posterior pituitary was from two to three times as large as the anterior lobe; whereas, in the normal, the posterior lobe is about one-third the size of the anterior. With a posterior lobe about six times as large as the normal, one can expect some positive feature in the body indicative

of this exaggerated function of the part, and in the case referred to the exaggerated feature was in all probability infantilism. This suspicion is, to some degree, confirmed by Sharpey-Schafer, who states that if posterior pituitary is grafted into young tadpoles they subsequently show definite retardation of growth. Can it be that the function of the posterior pituitary is to check growth? This would be in accordance with the older view that the ductless glands (considering the pituitary and the adrenal each as two glands) work in pairs, one giving rise to a stimulating impulse, and the other checking the former by acting in the opposite manner.

Although the pituitary is the seat of the growth-promoting hormone, and, according to the above hypothesis, of the growth-checking hormone, metabolism is almost as important in the growth process. Hence, in malignant disease, the glands concerned in metabolism must also be considered. That the endocrines might be disturbed in cancer seemed probable. Could the course of the disease be influenced by adjusting the workings of any or all of the glands?

HISTOLOGICAL OBSERVATIONS

In order to appreciate fully the character of this investigation it would be as well to state now that the ideas expressed in this work gradually evolved during a histological survey of the whole endocrine system—on the pituitaries, thyroids, and adrenals, of over 200 cases, and on the parathyroids, thymuses and pancreases, of about 100 cases or over. The selection of the material depended entirely on the state of preservation of the tissue, and not on the type of the case. Hence the number of glands from cancer cases was limited.

The following data on the pituitary were based on an examination of 230 glands, of which 30 were from cases of malignant disease—including 13 of glioma cerebri.

Working on Stewart's hypothesis (1921) that the cells of the anterior lobe are merely in different stages of function, and that the acidophilic cells contain the active secretion, I found that the anterior pituitary in the tumour cases did show a degree of activity appreciably higher than in the normal. Histologically, Zondek's conclusion that 15 per cent. of cases of malignant disease give a positive Zondek-Aschheim reaction was confirmed. In many, or perhaps all, cases of malignant disease, then, the anterior pituitary might be considered as overactive.

The Posterior Lobe

The posterior lobe of the hypophysis offered a more difficult task. The belief that the posterior part of this gland was essentially neuroglial was found to be untenable. Preparations stained for neuroglia showed the neuroglia distributed as strands about circular or elliptical masses of tissue, and in well-preserved sections stained by haemalum and eosin the parenchymatous character of the lobules was evident (Fig. 1 on Plate). The posterior pituitary, then, consists of lobules of roundish epithelial cells; these lobules were well vascularized and surrounded by a neuroglial stroma. The colloid of this section of the pituitary was taken to be its effective secretion, and appeared to emanate from the epithelial lobules, from whence it passed through the veins of the posterior pituitary, to the vessels of the pars intermedia, where the secretions of both anterior and posterior lobes intermixed. If, as already suggested, the secretion of the posterior pituitary is concerned with checking growth, then this lobe of the gland should show changes, with an incidence parallel with that of cancer, in the various age groups; while in cancer cases it should show these and corresponding changes in a greater degree and more frequently.

In this study the brown pigment of the posterior pituitary was frequently found associated with, or in,

E. WATSON-WILLIAMS: THE BROKEN NOSE

FIG. 1.—Case III. Crushing of right side of nose.

FIG. 2.—Case IV. Crushing of nose with fracture involving antrum.

FIG. 3.—Case I. Characteristic deformity of "mild" type.

FIG. 5.—Case II. Severe crushing; profile restored, triangle splint used.

FIG. 7.—Case VI. Crushing and laceration of nose, after eight days.

FIG. 4.—Case I. Four days after reduction, no splints used.

FIG. 6.—Case V. Result after severe crushing with fracture of maxilla.

FIG. 8.—Case VI. Result after reduction, triangle splint used.