two. The low pulse pressure which frequently occurred after an injection of pituitrin merely caused a transient faintness. Finally, the effectiveness of the dose was gauged by the temperature, for when the temperature varied between 97° and 99° it was judged that the regressive changes were sufficient. When toxic absorption caused the temperature to rise above 99°, the usual measures were employed to bring it down.

SUMMARY

1. Based on the hypothesis that in cancer cases the anterior pituitary was overactive, and the posterior pituitary underactive, two cases of advanced cancer were treated with pituitrin alone, and five cases with pituitrin and theelin. All were placed on a diet low in carbohydrate.

2. An epithelioma began to separate off and was enucleated after seven weeks of this treatment. The growing edge disappeared in five days.

3. All the cases showed regressive changes in the tumours, and life appeared to be definitely prolonged.

I wish to thank the honorary, medical, and nursing staff of the Christie Cancer Hospital, and Dr. Stanley White of Messrs. Parke, Davis and Co., for their valuable co-operation in the clinical work. I am indebted to the Manchester Committee on Cancer, and the Manchester University Grants Committee for grants to defray the expenses of this investigation. For the photographs and photomicrographs I have to thank Mr. H. C. Taylor of the pathological department.

BIBLIOGRAPHY

- Borrel, A., and de Coulon, A.: *Compt. rend. Soc. de Biol.*, 1922, lxxxvi, 1096.
 Carse, G.: *Med. World*, 1918, x, No. 21.
 Hande, M., and Tadenuma, K.: *Zeit. f. Krebsforsch.*, 1924, xxi, 288.
 Norgate, J. H.: *Brit. Journ. Surg.*, 1921, ix, 495.
 Reding, R., and Slosse, A.: *Compt. rend. Soc. de Biol.*, 1926, xcv, 1442.
 Sharpey-Schafer, F.: *The Endocrine Organs*, Part II, p. 250.
 Stewart, F. W.: *Compt. rend. Soc. de Biol.*, 1921, lxxxiv, 49.
 Warburg, O.: *Biochem. Zeit.*, 1923, cxlii, 317, 334; *Ibid.*, 1924, cxlvi, 309; *Klin. Woch.*, 1924, iii, 1062.
 Zondek, B.: *Klin. Woch.*, 1930, ix, 679.

TORSION OF OVARIAN CYST IN CHILDREN

BY

W. ARKLEY STEEL, M.D.St. AND., M.R.C.P.LOND.

REDHILL HOSPITAL, EDGWARE

(With Special Plate)

The age incidence of ovarian tumours has been noted in several valuable series of recorded cases. Statistics indicate the occurrence of ovarian tumours at the extremes of age and their comparative infrequency in childhood. The proportion of cases met with in children does not exceed 3 per cent. Illustrative of the early age at which ovarian tumours may appear is the case reported by Doran of an ovarian tumour in a premature child, aged 7 months. Another instance is given by Chiene, who performed ovariectomy on a child aged 3 months. The majority, however, of these tumours in children are met with between the ages of 10 and 14. Both the solid and the cystic types of tumour of the ovary are found in children. Of the simple variety, cystadenomata and dermoid cysts are equally common, and constitute about two-thirds of all tumours. The malignant growths form the remainder, and, of these, sarcomata are rather more common than carcinomata.

Cases of ovarian tumours are brought to our notice by reason of: (1) their increasing growth giving rise to abdominal deformity; (2) constitutional disturbances; (3) mechanical effects, causing pressure on the surrounding viscera; (4) complications. In children the complications of rupture, or of suppuration of an ovarian cyst,

are exceedingly rare, and torsion, although the commonest complication, is relatively uncommon. The rarity of the condition justifies the detailed description of the following case, which was under my care.

ILLUSTRATIVE CASE

J. L. M. J., female, aged 9 years, was admitted to hospital on October 13th, 1930, as a surgical emergency. The parents were both healthy, but somewhat dull intellectually. There was one other sister alive and well.

Previous History.—The patient was born after a short and normal labour, and, except for measles and whooping-cough, she had always been very healthy. When about 5 years of age she had an attack of pain, which commenced on the left side and extended across the upper part of the abdomen. This was accompanied by retching, and "trouble with the bowels." These attacks recurred each year, but were never severe enough to confine the child to bed. The mother was emphatic that the onset of each attack synchronized with the approach of the colder weather, taking place usually in October. The pain lasted as a rule for fourteen days, but eventually responded to doses of castor oil. She had had attacks of vague abdominal pain frequently throughout the last year.

Present Illness.—At 9 p.m., on October 5th, 1930, the patient had a sudden attack of vomiting. There was no pain at first, but the next day she complained of pain in the left side of the abdomen. This pain extended across the upper part of the abdomen, and in no way differed from that experienced in previous attacks. She attended school from October 7th to 10th, but at 11 a.m. on the latter date she had a sudden attack of pain, which "doubled her up." The pain started in the left side, and passed down to the left groin and across the abdomen. She vomited several times. None of the measures which had been successful in previous attacks relieved the pain. The child had a peculiar taste for castor oil, and of her own accord frequently took copious draughts straight from the bottle. There was no action of the bowels from October 9th until the 13th, when she passed a normal motion. Micturition was normal up to the day of admission, when she complained of burning pain on passing urine. The abdominal pain continued, and her general condition became gradually worse. She was sent to hospital as a case of acute appendicitis.

On later interrogation, the mother said that for the past three years she had noticed her child's abdomen becoming more prominent, and that it "seemed to be getting hard on the left side." The patient often acted in a peculiar manner, and the mother was inclined to regard her as somewhat "simple-minded." She had never menstruated, nor was there any precocious sexual development.

Examination.—On admission the patient was extremely ill and dehydrated, crying out from time to time owing to spasms of pain. The temperature was 100.8° F., pulse rate 124, respirations 24. The tongue was furred and dry. The upper part of the abdomen moved slightly with respiration, the lower part not at all. A globular swelling, extending from the pubis to the umbilicus, was apparent, filling the hypogastrium and extending to the left side. Palpation confirmed the presence of this swelling, which was felt to be tense, tender, and cystic. It appeared to rise from the pelvis, and was rather more prominent on the left side. It was so centrally placed, however, that the possibility of its being bladder was considered. Catheterization was performed, but had no effect in diminishing the size of the tumour. There was generalized tenderness over the abdomen and in the loins. Rigidity was present over the whole abdomen, but was most marked in the muscles overlying the tumour, especially on the left side. On rectal examination a hard, tense, tender, and cystic swelling was felt filling the pelvis, and so bulging the anterior wall of the rectum as practically to occlude the lumen of the bowel. A slight degree of vulvitis was present, with a scanty yellowish discharge. The examination of the urine showed the presence of ketones, but was otherwise normal. The other systems revealed nothing abnormal. The diagnosis of acute appendicitis was excluded. The condition was thought to be an abscess, circumscribed in the pelvis, due possibly to an extension from the vulvitis. An operation was decided upon, and was performed at 11:50 p.m. on October 13th.

Operation.—Under chloroform and ether anaesthesia the abdomen was opened by a right paramedian incision. A large, mottled cyst presented itself in the wound, its upper limit extending above the level of the umbilicus. Further examination showed this to be a left ovarian cyst, which completely filled the pelvis. The pedicle was found to have twisted three and a half turns in a clockwise direction. Delicate adhesions were present between the cyst and the posterior aspect of the infantile uterus, and these were readily separated. The right ovary was normal. There was some clear sanguineous fluid in the pelvis. The pedicle was clamped and ligatured, and the cyst removed. The stump was peritonized, and the abdomen closed without drainage. The child made a complete recovery, and was discharged on October 29th.

Morbid Anatomy.—A photograph of the specimen (see Special Plate) shows an ovarian cyst with a broad, thinned pedicle, and also part of the Fallopian tube, which was excised with the tumour. The cyst was thin-walled, somewhat mottled in appearance, due to the deposition of fatty particles on the inner side. The contents were clear fluid, in which were floating globules of fat, and also fine hairs. The tumour is a typical dermoid cyst, and measures $4\frac{1}{2}$ by 4 by 4 inches.

FEATURES OF THE CONDITION

In a somewhat extensive review of the literature dealing with cases of torsion of an ovarian cyst in children, it is noticeable that, although cases have been reported in greater or less detail, very little attempt has been made to correlate the clinical findings of the condition. In several instances, more stress has been laid on its rarity than on the description of the actual case itself and the recognition of its possibility in diagnosis. In the above case and in those reviewed there are many common features which warrant fuller description.

Pain.—In many instances a history is reported of similar attacks, which have been of less severity, but have recurred at intervals of months or, less frequently, of years. These previous attacks are of short duration, lasting in the majority of cases but a few hours, although they may continue for several days. In the acute attack the pain is often preceded by vomiting, or vomiting may occur after the pain and be almost continuous throughout. The onset is sudden, and the pain is very severe in character, being usually constant, but sometimes assuming a spasmodic character. It is definitely localized to the side on which the tumour is situated, but radiates medially across the abdomen, so that the umbilicus may be indicated as the site of the pain. The severity of the initial excruciating pain may diminish after a varying time, leaving a dull ache.

General Condition.—The patient looks ill and may be definitely toxæmic. She is exhausted and apprehensive, often crying out from time to time, owing to spasms of pain. The tongue is furred and dry, and the bowels have usually opened naturally, or have responded to laxatives. In a few cases there may be a history of constipation of several days' duration. The pulse rate is raised, while the respiration rate is normal. The temperature is raised usually above 100° , but it varies between 99° and 103° .

Abdominal Examination.—A tumour can be recognized in the majority of cases; in twenty-one cases in which the situation has been given, the cyst has occurred thirteen times on the right side and eight times on the left. The cyst may produce only a general prominence of the abdomen, but in many instances the tumour is definitely demarcated. It may be outlined in the iliac fossa, but frequently it assumes a median position, and gives rise to the false impression that it is a distended bladder. The tumour is tense and fluctuating. Tenderness is present, and is most marked over the site of the tumour. It may be localized to a very small area in the iliac fossa, or may be general over the whole abdomen. In one case, no pain or tenderness was experienced. Rigidity occurs

and is usually well marked, especially in the muscles overlying the tumour, but it may be generalized.

Rectal Examination.—A rectal examination appears to have been made in relatively few of the cases, but when carried out a definite mass was always palpated in each case. In some instances this appeared to occupy practically the whole of the pelvis and markedly compress the walls of the rectum. The tumour is rarely movable. It is acutely tender, and gives a feeling of fluctuation to the examining finger.

Age Incidence.—The age in this series varied, the oldest patient being 13 years; the youngest was reported by M. L. Harris, who describes a case in which a twisted malignant ovarian tumour was successfully removed from a child aged 22 months.

Complications.—In a few cases a slight degree of vulvitis has been present. A blood-stained serous fluid is often found in the abdominal cavity, but it is of too small amount to be appreciated before operation. There may be small areas of localized peritonitis, and recent, or established, adhesions are usually present, which can be readily broken down and give rise to no difficulty in the operation. In several cases the appendix, which may be congested, has been found adherent to the cyst. In one case intestinal obstruction due to twisting of the ileum following removal of a cyst rendered a second operation necessary.

Diagnosis

More than twenty-five cases of torsion of an ovarian cyst in children have been investigated. Since this investigation a further case has been published and I have heard of the occurrence of yet another. Possibly, therefore, it is a commoner condition than the few published cases would lead one to believe. It is thus a striking and humiliating fact that, of all the cases investigated, only four were diagnosed prior to operation—surely an indication that ovarian cyst, quite apart from torsion, had not been considered in the differential diagnosis. Each case has been regarded—and rightly so—as an abdominal emergency requiring operation. The diagnosis most commonly made is that of a localized appendix abscess, and where there is a history of previous attacks, increased temperature and pulse rate, tenderness, rigidity, and vomiting, and especially where there is a right-sided tumour, it may be extremely difficult to differentiate the two conditions. In a case of torsion of a cyst, the onset as a rule is sudden, the bowels have been regular, and the presence of a tumour can be made out by abdominal and rectal examination from the very commencement of the attack. The diagnosis of intussusception has been suggested, but the clinical features of this condition, its greater frequency during the first year and in boys, the passage of mucus and blood in the stools, and the typical emptiness of the right iliac fossa, are absent in the ovarian lesion. Abscess formation must be excluded, and if no definite causative factor can be established to account for it, an abscess, circumscribed in the pelvis, may give rise to great difficulty in diagnosis. The knowledge that torsion of an ovarian cyst can so closely simulate an acute suppurative lesion should render it possible for a correct diagnosis to be made more easily and more frequently.

SUMMARY

1. Over twenty-five cases of torsion of ovarian cyst in children have been reviewed, and an illustrative case is given.
2. Ovarian tumour, though rare, increases in frequency with the age of the child, and is most commonly met with between the ages of 10 and 14.
3. Torsion, although very rare, is the commonest complication of these tumours.
4. A correct diagnosis of the condition is decidedly more often the exception than the rule, in spite of the fact

that a constant clinical picture is found in practically every case.

5. The typical features present are: a history of similar but less severe attacks, the sudden onset of pain, an increase of temperature and pulse rate, the palpable abdominal tumour, with overlying tenderness and rigidity, and the tender cystic mass, which can be made out per rectum.

6. The diagnosis is sometimes very difficult, the lesion for which torsion of a cyst is most often mistaken being an appendix abscess.

I have to thank Dr. Teale, who kindly examined the tumour for me and also furnished me with the photograph. The specimen is now in the Pathological Museum of University College Hospital.

BIBLIOGRAPHY

- Macrae, R. C. B.: *British Medical Journal*, 1930, i, 906.
 Angus, H. Brunton: *Ibid.*, 1906, i, 199.
 Kittell, P. B.: *St. Bartholomew's Hospital Journal*, 1919-20, xxvii, 24.
 Owen, Edmund, and Greene, Clayton: *British Medical Journal*, 1904, ii, 1517.
 Harris, M. L.: *Amer. Journ. of Obstet.*, 1904, i, 530.
 Rainey, Warren R.: *Annals of Surgery*, 1924, lxxix, 879.
 Sutcliffe, W. Greenwood: *Lancet*, 1907, ii, 772.
 Curtis, P. Murray: *Med. Journ. of Australia*, 1928, i, 497.
 Goldstine, Mark: *Surg., Gynecol. and Obstet.*, 1920, xxx, 627.
 Morris, J.: *British Medical Journal*, 1926, i, 944.
 Weil, Harry I.: *Bull. Johns Hopkins Hosp.*, 1905, xvi, 102.
 Froelich, M.: *Bull. de la Soc. d'Obstét. et de Gynéc.*, 1922, xx, 497.
 Riche, V.: *Ibid.*, 1928, xvii, 276.
 Paterson, Herbert J.: *Brit. Journ. of Children's Diseases*, 1911, viii, 295.
 Kincaid, Harvey L., and Andrews, Edmund: *Amer. Journ. Obstet. and Gynecol.*, 1928, xv, 207.
 Powell, E. M.: *Lancet*, 1923, i, 751.
 Cahill, James A.: *International Clinics*, June, 1924, ii, 212.
 Downes, William A.: *Journ. Amer. Med. Assoc.*, 1921, lxxvi, 443.
 Cavafy and Haward: *Lancet*, 1886, i, 920.

GANGRENE OF THE SKIN OF DIPHTHERITIC ORIGIN

BY

J. S. ANDERSON, M.A., M.D., D.P.H.

MEDICAL SUPERINTENDENT, LEEDS CITY HOSPITALS; LECTURER IN INFECTIOUS DISEASES, UNIVERSITY OF LEEDS

(With Special Plate)

Two types of diphtheritic infection of the skin may be described. In one a diphtheria organism is isolated, but no symptoms characteristic of the disease are present. This type, which is not uncommon, would probably be more correctly regarded as bacteriological diphtheria. In the other type the diphtheria organism is the active agent in the production of the local disease. This type is of rare occurrence. Cutaneous diphtheria is usually found to be secondary to diphtheria in some more normal site, such as the fauces, nose, or genitals, but primary diphtheria of the skin may occur. The absence of a source of origin of the infection, and the unusual site of the disease, combine to make the present case worthy of record.

The patient was a boy, aged 11 years. According to the history, he visited a swimming bath on May 20th, 1931, and sustained a slight injury of the nature of a bruise on the dorsum of the right foot. Persistent pain and discomfort led to his attendance at the Leeds Public Dispensary on May 26th. The house-surgeon described the injury then as showing the usual appearance of inflammation and the semblance of a greyish pellicle in the centre. At first sight it was taken for a burn. The lesion progressed until, by June 1st, there was a surrounding zone of cellulitis, with, in the centre, a triangular area, which was dark grey and later black, and suggested a localized form of gangrene. Incisions into the affected area revealed no pus and produced no improvement. On account of the unusual appearance of the lesion he was retained as an in-patient. Slight general disturbance was

noted from June 1st to 3rd, but the temperature did not rise above 99°. On June 4th, however, it rose to 104°, and multiple small incisions were made in the surrounding inflammatory area. At the same time a swab was taken. On June 5th and 6th the temperature hovered around 102°, and no improvement was noted. On June 6th, following the receipt of a report that diphtheria organisms were present, he was transferred to the Leeds City Hospital.

Fig. 1 (Plate) shows the appearance on admission to the Leeds City Hospital. The central area (C) consisted of a dry black eschar. Surrounding it was a greyish pellicle of semi-necrotic skin (B), and the remaining part of the dorsum of the foot showed much inflammation and oedema at (C). Neither lymphangitis nor adenitis was noted. No offensive odour was associated with the lesion. The tonsils were slightly enlarged, but there was no sign of inflammation or of exudate. There was no nasal discharge, and the anterior nares appeared healthy. The temperature was now subnormal and the pulse 90. Albuminuria was present. The Schick test was positive. Diphtheria antitoxin, 16,000 units, was administered on the day of admission, and no treatment other than dry dressings was applied to the foot for the next ten days.

Within twenty-four hours of admission the inflammatory reaction commenced to subside. The greyish pellicle ceased to extend, but became necrotic within the next few days. Eleven days after admission (Plate, Fig. 2) the lesion had taken the form of a crater with well-defined margins, and with a floor of dry, black, necrotic skin. After a further eight days, the necrotic tissue sloughed, leaving a healthy-looking granulating surface. Albuminuria persisted throughout the whole period of isolation, but no other complications were noted. By the end of July the ulcer had healed completely, and he was discharged from hospital on August 12th.

Note on Bacteriology.—The swab taken on June 4th was examined at the Bacteriological Department of the University of Leeds, and the report stated that *Corynebacterium diphtheriae* was present in pure culture. The organism gave the correct sugar reactions, and was proved to be a virulent type. On the day of admission to the Leeds City Hospital swabs taken from the nose and throat showed no diphtheria organisms, while a swab from the inflammatory area showed a pure culture of *C. diphtheriae*. On June 9th swabs from the necrotic skin and from the greyish pellicle also showed *C. diphtheriae*. On June 10th a swab taken from the greyish pellicle was again examined at the University of Leeds, and the report stated that aerobic and anaerobic cultures showed *C. diphtheriae* in pure culture. A nose swab was again negative on June 12th. On July 1st diphtheria organisms were still grown from the floor of the ulcer, but after July 5th swabs were negative.

I am indebted to the Bacteriological Department of the University of Leeds for laboratory work, and to Dr. J. F. Galpine, for helpful information regarding the early stages at the Leeds Public Dispensary.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

A CASE OF WOUND DIPHTHERIA

(With Special Plate)

The false membrane type of diphtheria of a wound and surrounding skin, once seen, cannot readily be forgotten.

A girl, aged 11 years, was admitted to the Royal Victoria and West Hants Hospital on February 2nd, 1929, on account of appendicitis. For four weeks, off and on, she had had abdominal pain of varying intensity in the right iliac region, with obstipation but without nausea and vomiting. When she was admitted the pulse rate was 104, the tongue was furred, and there was local tenderness with rigidity. On February 19th, Mr. Kinsey Morgan removed an inflamed appendix through a grid-iron incision, then closed the wound in the usual way.

W. ARKLAY STEEL: TORSION OF OVARIAN
CYST IN CHILDREN

FIG. 1.—Cyst of left ovary from a girl aged 9 years.

S. WATSON SMITH: WOUND DIPHTHERIA

FIG. 1.—Wound diphtheria of false membrane type.

J. S. ANDERSON: GANGRENE OF THE SKIN OF DIPHTHERITIC ORIGIN

FIG. 1.—Appearance on admission.

FIG. 2.—Appearance eleven days after admission.