All who have had experience of arthroplasty are agreed that the most successful cases are those in which the ankylosis is bony, with comparatively normal-shaped bones. This will include traumatic joints, and those the result of pyaemia. We may exclude tuberculous joints, which so often and for years contain foci of infection.

The operation is contraindicated in the presence of active disease, and should be delayed until at least two years after its complete cessation. It should never be performed in the presence of sinuses, nor where the soft tissues are infiltrated and tense, nor where muscles are atrophied and adherent; nor should it be performed until growth is ended. The joint which responds most favourably to an arthroplasty is the elbow; the knee, although technique has improved, mainly by the excellent work of Putti, is still in the experimental stage. It is an operation which requires the highest judgement and skill on the part of the surgeon, and fortitude on the part of the patient, for the after-treatment is very painful. The elbow is a joint where we can almost promise a satisfactory result—good movement and fair stability; but the ankle-joint, if stiff and painless, should not be mobilized; if painful it should be arthrodesed.

BIBLIOGRAPHY

- C. W. Mansell Moullin: *Sprains*, H. K. Lewis, 1887.
Wharton Hood: *Treatment of Injuries*, Macmillan, 1902.
Frank Romer: *Modern Bonesetting*, Heinemann, 1915.
Timbrell Fisher: *Manipulative Surgery*, H. K. Lewis, 1925.

AVERTIN IN GYNAECOLOGY

A RECORD OF 1,000 CASES

BY

JAMES YOUNG, D.S.O., M.D., F.R.C.S.ED., F.C.O.G.,
GYNAECOLOGIST, ROYAL INFIRMARY, EDINBURGH

In a previous paper I¹ have put on record the impressions gained after the employment of avertin in 345 unselected gynaecological operations. Our experience has now extended to over 1,000 cases, and increasing use has gone to confirm the view expressed in 1929 by Blomfield and Shipway² in their paper introducing avertin to British surgery that it is "a valuable addition to the drugs at the disposal of the anaesthetist."

The method of administration adopted is similar to that described in the previous communication and to that employed by other surgeons, and it is unnecessary here to enter into a detailed description. The procedure is very easily learned. The evening before the operation the bowel is washed out with an enema. One hour before, one-eighth of a grain of morphine is given hypodermically. Before removal from bed, the patient turns on the side and the drug in solution is slowly instilled into the rectum, the administration taking about fifteen minutes. Before the lapse of this period the patient is usually unconscious, the induction of the anaesthesia being smooth and pleasant. Throughout the induction and the period of unconsciousness the anaesthesia is strongly reminiscent of a deep natural sleep. On two or three occasions only have we found some mental excitement during the induction, while on three occasions the patients failed to pass over into complete unconsciousness. In all cases there is complete amnesia. The maximum anaesthesia is reached about thirty minutes after the commencement of the induction, and we plan that this interval will elapse before the operation is begun, although on many occasions, owing to the exigencies of the operating theatre, we have without disadvantage allowed over an hour to pass.

The drug is quickly absorbed from the rectum, but the dissolving fluid remains in the bowel for a longer

time, and in the case of operations carried out in the lithotomy position it is apt to be squeezed out during the examination or the introduction of the speculum, with a consequent soiling of the parts. To prevent this, it is our practice in such cases to withdraw the remaining fluid by means of a rubber catheter immediately before the commencement of the operation.

We employ, as a routine, a dosage of 0.1 gram per kilo of body weight. In very obese patients instead of giving the full ratio we employ a diminished dosage of 0.08 or 0.06 gram per kilo. We use the liquid preparation which contains 1 gram of avertin in each cubic centimetre, and by means of the chart supplied by the makers the respective quantities can be determined easily and rapidly. With this dosage the anaesthesia obtained is not profound, and, although sufficient often to allow of a minor procedure, such as curettage and, occasionally, of a plastic operation for vaginal prolapse, we find it an advantage as a routine to amplify it by means of ether given by the open method. The quantity of ether required is usually much less than that necessary for an unaided inhalation anaesthesia. It may now be claimed that the experience of many clinics engaged in all classes of surgical work has established that, when employed as a "basal anaesthetic" in such dosage as I have mentioned, avertin is a safe drug. During the early days fatalities were recorded, but there can be no doubt that these arose almost invariably from pushing the dosage to the stage of inducing a complete and self-sufficient anaesthesia.

ANALYSIS OF CASES

The drug has been used as the routine anaesthetic in my department during the period under review, and, except where on a few occasions a pelvic swelling—for example, fibroid tumour, pelvic abscess—has so encroached upon the rectum as to impose a mechanical obstacle to the instillation of the fluid, there has been no selection of cases. For this reason the experience can be accepted as representing that of an average gynaecological hospital practice in which, although much of the operating is of a minor nature, there is naturally at the same time a large admixture of grave and emergency material. Thus there were over 200 cases necessitating opening of the abdomen for the carrying out of hysterectomy (71), salpingo-oophorectomy (24), ovariectomy (34) and removal of cysts with twisted pedicles (6), removal of the tube in ruptured tubal pregnancy (16), myomectomy for fibromyoma (10), appendicectomy (25), etc. There were 130 cases of plastic repair for vaginal prolapse. The age of the patients has varied from 5 to 76.

In the series there were five deaths, and in no instance could the fatal issue be attributed either directly or indirectly to the avertin.

The clinical data are briefly as follows:

1. K.117, 8-para, aged 58, was admitted in a collapsed state with complete acute inversion of uterus and vaginal prolapse associated with uterine fibroid. Death occurred twelve hours after hysterectomy.
2. R.156, 4-para, aged 40, with an abdominal swelling of a year's duration, was admitted with a history of vomiting, acute abdominal pain, and fever. Death occurred twenty-eight hours after operation, involving removal of an infected and adherent ovarian tumour.
3. D.762, 1-para, aged 40, was admitted for a marked degree of vaginal prolapse. There was a history of toxic goitre for fourteen years. She developed broncho-pneumonia four days after a modified plastic repair operation, and died ten days later.
4. I.366, 9-para, aged 40, was admitted with a history of breathlessness, menorrhagia, and, recently, of retention of urine. She collapsed four hours after removal of cervical fibroid by total hysterectomy, and died three hours later. The post-mortem revealed chronic toxic changes in spleen, liver, and kidneys.

5. F. 289, 1-para, aged 60, was a very obese patient weighing 15 stone, with very advanced and generalized abdominal carcinoma, probably originating in the uterine body. She died from heart failure thirteen days after exploratory curettage.

In addition to these five deaths, there was in hospital during the period under review one death after laparotomy and drainage of an abdominal abscess, which extended deeply into the pelvis, and in which ether anaesthesia was employed.

This mortality rate of 5 in 1,000, or 0.5 per cent., is much below our previous mortality rate of 34 in 1,799 consecutive cases, or 1.9 per cent., with ether, alone or in combination with chloroform, as the routine anaesthetic. We do not claim that in this reduction the sole part has been played by the employment of avertin, for it is possible to cite ways in which the general advance in our knowledge of gynaecological therapeutics within recent years has displaced surgery in favour of conservatism, and has thus contributed towards a saving of lives. During the period under review, for example, over forty cases of acute salpingitis (both gonococcal and streptococcal) were treated palliatively without mortality, and a large number of cases of intractable uterine haemorrhage were treated by non-operative means, such as radium, x rays, etc. By curtailing the necessity for recourse to major surgery these advances have necessarily tended to a general lowering of our mortality figures. Further, the recognition of the common part played by cervical infection in the production of low abdominal and iliac fossa pain, which at times may even assume acute characters, has, by improving our diagnostic weapons, reduced abdominal exploration for suspected appendicitis, tubal and ovarian disease, etc.³ The greater insistence on a rigid pre-operative investigation and preparation, and in those cases in which post-operative infection is a special risk—for example, in vaginal prolapse—the prophylactic immunization of the patient with a streptococcal and *B. coli* vaccine, have also doubtless helped to reduce the post-operative morbidity and mortality.

After making all these allowances, however, we are convinced that the adoption of avertin as a basal anaesthetic has not only removed much of the distress formerly experienced by the patient, but has in a material way eliminated the risks incidental to all surgical procedures.

ADVANTAGES CLAIMED FOR AVERTIN

1. *Comfort of Induction.*—The importance of this factor cannot be overestimated. It is a matter of favourable comment by patients who have on previous occasions experienced other methods of anaesthesia. Apart from this softening of the memories of the operation day, however, we have the advantages accruing from the lessening of the pre-operative emotional stress. The work of Crile⁴ and others has emphasized the importance of these questions by directing attention to the profound physical changes induced in the organism by emotion. The sleep induced before the patient leaves her bed implies that with avertin we have a simple and effective method of eliminating a whole series of dread-inspiring preliminaries—the tense expectation of the event, the arrival of the attendant to herald its approach, the agitation accompanying removal to the ante-room, and the administration of the anaesthetic. To many patients these circumstances constitute the most vivid memories of the operation, and it is a fact that the more acute sensibilities of women render them especially prone to such mental suffering.

2. *Reduction of Post-operative Distress and Morbidity.*—In ordinary circumstances the patient sleeps deeply for several hours after the operation, and thereafter for a considerable time the sensations remain dulled, so that the awaking to full consciousness, with its attendant

discomfort and pain, is delayed often by many hours. Further, one of the most important of the advantages of avertin is the virtual elimination of the tendency to retching and sickness after the operation. The hospital ward following an operation day is now practically undisturbed in this respect, with the consequent removal of a grave source of distress, not only to the victims themselves, but also to the other patients, and with a consequent lessening of the work of the nursing staff. It cannot be denied that the relief brought in these ways to the patient in the immediate post-operative phase must imply, in addition, a reduction in the risk to life and health. The abolition of the straining incidental to vomiting by leaving the wounds undisturbed diminishes the chance of haemorrhage and sepsis. The importance of these advantages can hardly be overestimated in regard to laparotomy and pelvic floor wounds. Dandy⁵ has recently called attention to the same benefits which avertin confers in major brain surgery.

3. *Reduction of Post-operative Pulmonary Complications.*—Post-operative bronchitis and pneumonia, formerly a not infrequent source of anxiety and danger, have been reduced. In a total of 203 cases in which opening of the abdomen was carried out there has been an entire absence of such post-operative pulmonary troubles. Altogether, in the series of 1,000 cases, there have been three instances of mild bronchitis, one of pleurisy, and three of broncho-pneumonia. There was one death from broncho-pneumonia, which developed after a plastic operation for prolapse in a woman with toxic goitre. It is at first sight rather remarkable that, although we employ ether so consistently in conjunction with avertin, we have had only one death from broncho-pneumonia in this large series. The lessened quantity of the inhalation anaesthetic required after avertin plays, of course, an important part. We have come to the conclusion that of equal, if not of greater, moment is the fact that the smoothness of the induction and the elimination of retching, both during the course of the operation and the early post-operative phase, lead to a wholesale reduction in the risk of septic inhalation. Many other observers have referred to the infrequency of pulmonary complications after the employment of avertin. For example, Dandy⁵ recently has recorded "250 major cranial operations of every type without mortality from the anaesthetic, no post-operative pneumonia, and no deleterious effect immediate or remote." Dandy believes that with avertin there is a reduction in the risk of aspiration of gross septic material into the lungs. Wülfing,⁶ in a recent paper, based on a record of 1,200 cases, refers to the same facts, and states that in his series pulmonary troubles have been fewer and less severe than with ordinary inhalation anaesthesia. The diminished risk of pulmonary irritation makes avertin of special value where anaesthesia is contemplated in patients already suffering from bronchitis, asthma, etc.

It should further be noted that in our series there have been no cases of acute dilatation of the stomach or paralytic ileus. It is possible that the absence of these complications is to be related to the use of avertin, although on this question further data are necessary.

DISADVANTAGES OF AVERTIN

1. *Lowering of Blood Pressure.*—This constitutes one of the most striking effects of avertin. We have found that a systolic drop of 20 to 30 mm. Hg is not uncommon, and in cases of hypertension the fall may be as great as 100 mm. Hg. It commences with induction and usually reaches its maximum within fifteen to twenty minutes. Dr. W. P. Kennedy, working in my department, has subjected the blood pressure during avertin anaesthesia to a close study in 150 cases, and has published the results in the *British Journal of Anaesthesia*.⁷ This paper should

be consulted by those interested. In this place it may be stated that our conclusions are that the fall in blood pressure is of no practical significance. Even in the cases showing the maximum drop the general condition has not given rise to any anxiety. In many cases we have noted that the pressure is quickly restored spontaneously or after the administration of the ether is begun. It is quickly readjusted by the hypodermic injection of 1/2 c.cm. of a solution of ephedrine hydrochloride (50 mg. per c.cm.), and this we utilized commonly during our early experience, although, since recognizing that the condition is clinically unimportant, we are now using drugs less and less. Our findings in these respects are paralleled by those of other workers (for example Dandy).

2. *Depression of Respiration.*—Slowing and reduction in the amplitude of respiration are common, and this effect at first gave us some concern. We have, however, seen no harm arise from it. Where severe, it is at once readjusted by administering carbon dioxide. For this purpose we find it advisable that a cylinder of oxygen containing 5 per cent. CO₂ should always be at hand in the theatre.

A minor disadvantage of avertin is the headache of which a considerable proportion of patients complain on waking from the anaesthetic. It is readily controlled by pyramidon 5 grains (repeated after four hours if necessary, although one dose usually suffices).

CONCLUSIONS

1. Employed in a dosage which does not exceed 0.1 gram per kilo of body weight, avertin is a safe drug. In this dosage it may suffice for minor operative work, but a supplementary inhalation anaesthesia is usually required.

2. Its advantages are: (a) ease, comfort, and certainty of induction; (b) reduction of post-operative discomfort and sickness; (c) reduction of post-operative pulmonary complications.

3. In this series of 1,000 unselected cases there have been no deleterious effects, either immediate or remote.

4. There have been five deaths in the series, and in no instance could the fatal issue be attributed to the avertin.

REFERENCES

- ¹ Young, J.: *Lancet*, 1930, i, 1177.
- ² Blomfield, J., and Shipway, F. E.: *Ibid.*, 1929, i, 546.
- ³ Young, J.: *British Medical Journal*, March 29th, 1930.
- ⁴ Crile, G. W., and Lower, W. E.: *Anoci-Association*, W. B. Saunders Company, Philadelphia and London, 1914.
- ⁵ Dandy, W. E.: *Journ. Amer. Med. Assoc.*, May 30th, 1931, p. 1860.
- ⁶ Wülfing, M.: *Med. Klinik*, May 1st, 1931, p. 655.
- ⁷ Kennedy, W. P.: *Brit. Journ. of Anaes.*, January, 1931, viii, 2.

SOME SIALOGRAMS

BY

L. N. PYRAH, M.Sc., Ch.M., F.R.C.S.,

AND

P. R. ALLISON, B.Sc., M.B., M.R.C.S.,

(From the Department of Surgery in the University and the General Infirmary at Leeds)

(With Special Plate)

Lipiodol radiography has become an extremely useful diagnostic procedure during the last few years. Lipiodol came into its own quite early in the diagnosis of diseases of the chest, tumours of the spinal cord, and sterility. It is an oil containing about 50 per cent. of iodine, and is therefore highly opaque to x rays. It is non-toxic, and is only absorbed in small amounts, while the firm nature of its iodine union renders it very unlikely that any patient in whom it is used will develop iodism.

Sialography has been little used because the cases upon which it can throw light are very few; but in a limited

number of cases, facts of both diagnostic and pathological interest can be obtained from the study of a sialogram. During the war, sialography was frequently used for the investigation of salivary fistulae. Both parotid and submaxillary glands have easily accessible ducts, so that the technique is not difficult.

TECHNIQUE AND X-RAY METHODS

The lipiodol injection is carried out by means of a metal syringe fitted with a plunger, also of metal, as shown in the accompanying sketch. The stem of the plunger is graduated in cubic centimetres, so that the amount of lipiodol injected can be measured. The needle

is a modified lachrymal duct needle made of silver, and has a blunt, slightly bulbous end; it is malleable, and the curve can be adjusted to suit any given case. The needle is fitted to the body of the syringe by a screw attachment. There is therefore no risk of the patient's mouth being coated with lipiodol from the sudden detachment of the needle during the process of injection. With gentle usage no harm can be done to the lining membrane of the salivary duct. The plunger is fitted with a ring for the surgeon's thumb, and the body of the syringe has two rings for the index and middle fingers. The instrument is thus very easily controlled. The usual precautions of warming the syringe, needle, and lipiodol are taken; there is then no difficulty in driving the lipiodol through the rather fine-bore needle. In our hands the syringe has proved eminently satisfactory. Payne¹ has obtained good results by the use of a simple glass syringe.

Parotid Gland.—The injection is performed with the patient sitting facing the surgeon, in a good light. In injecting the right Stenson's duct, the surgeon retracts the right cheek with his left thumb and forefinger, and looks for the orifice of the duct, which is situated opposite the crown of the second upper molar tooth. The opening in most people is slit-like; in some it is situated on a slight papillary elevation. If there is difficulty in finding the orifice, the gland may be made to secrete by giving the patient a slice of lemon, or simple pressure upon the gland may expel a few drops of saliva and enable the opening to be found. The needle of the loaded syringe is brought up to the orifice of the duct and is steadied by the surgeon resting it upon his left thumb. It is slid into the duct for about an inch. The introduction of the needle causes a slight cutting pain, but not sufficiently intense to cause undue discomfort.

The injection is continued gently until the patient experiences a sensation of discomfort in the region of the parotid gland; this indicates that the finer ducts are filled with lipiodol. About 1/2 to 1 c.cm. is used. The needle is quickly but gently withdrawn, and the patient is immediately placed in the recumbent position on the x-ray table. One of us (P. R. A.) was able to inject his own Stenson's duct while standing in front of a mirror in a good light. In none of our cases has any untoward complication occurred as a result of injection.

Submaxillary Gland.—The injection is performed in a similar manner. The patient sits with the mouth widely